2016 Florida Youth Substance Abuse Survey

Gilchrist Data Tables

©2016 Florida Department of Children & Families

	Gilc	hrist	Florida S	tatewide
	N	%	Ν	%
Sex				
Female	250	51.0	31,515	47.9
Male	233	47.6	32,905	50.0
Race/Ethnic group				
African American	27	5.5	14,666	22.3
American Indian	7	1.4	568	0.9
Asian	3	0.6	743	1.1
Hispanic/Latino	25	5.1	13,174	20.0
Native Hawaiian/Pacific Islander	0	0.0	144	0.2
Other/Multiple	61	12.4	7,523	11.4
White, non-Hispanic	365	74.5	28,309	43.0
Age				
10	1	0.2	65	0.1
11	34	6.9	3,856	5.9
12	60	12.2	8,338	12.7
13	74	15.1	9,230	14.0
14	66	13.5	9,454	14.4
15	84	17.1	10,070	15.3
16	64	13.1	9,684	14.7
17	64	13.1	9,348	14.2
18	36	7.3	4,799	7.3
19 or older	3	0.6	618	0.9
Grade				
6th	73	14.9	9,301	14.1
7th	75	15.3	9,215	14.0
8th	70	14.3	9,326	14.2
9th	78	15.9	10,140	15.4
10th	82	16.7	9,834	15.0
11th	51	10.4	9,254	14.1
12th	61	12.4	8,705	13.2
Overall Middle School	218	44.5	27,678	42.1
Overall High School	272	55.5	37,765	57.4
Total	490	100.0	65,776	100.0

Table 1. Major demographic characteristics of surveyed Gilchrist and Florida Statewide, 2016

Note: Some categories do not sum to 100% of the total due to missing values (e.g., not all survey questions were answered). In addition, rounding can produce totals that do not equal 100%. "N" represents the number of valid cases. In this table, county data are unweighted while statewide data are weighted.

				Gilchrist	t					Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Alcohol	24.2	51.5	38.4	41.2	26.3	48.0	40.1	21.8	51.4	41.3	37.1	23.6	51.5	39.1
Blacking Out		21.8	22.4	20.7		22.2	21.8		15.9	16.5	15.4		16.1	15.9
Cigarettes	9.8	25.7	19.8	17.9	9.9	24.2	19.0	8.0	18.5	14.0	14.1	8.2	18.5	14.1
Vaporizer/E-Cigarette	13.0	39.0	24.4	31.0	14.7	35.5	28.0	14.6	33.9	24.4	27.1	15.7	34.5	25.8
Marijuana or Hashish	5.6	25.5	16.1	18.3	6.7	26.9	17.0	7.0	31.7	21.4	21.3	8.1	31.8	21.3
Synthetic Marijuana		8.4	6.4	10.6		9.3	8.4		4.9	4.8	5.0		4.8	4.9
Inhalants	7.8	3.7	8.6	2.4	6.7	4.6	5.5	6.4	4.6	6.1	4.7	6.4	4.7	5.4
Flakka		0.5	1.0	0.0		0.4	0.5		1.0	0.9	1.1		1.0	1.0
Club Drugs	0.0	2.6	1.4	1.7	0.0	2.0	1.5	0.7	3.0	1.9	2.2	0.8	2.9	2.1
LSD, PCP or Mushrooms	0.9	3.5	2.5	2.3	0.8	3.2	2.4	1.0	4.8	2.7	3.7	1.1	4.9	3.2
Methamphetamine	0.0	0.8	0.4	0.5	0.0	0.2	0.4	0.5	0.8	0.6	0.8	0.5	0.8	0.7
Cocaine or Crack Cocaine	0.9	4.6	2.1	3.9	0.8	3.8	3.0	0.8	2.5	1.6	1.9	0.8	2.3	1.8
Heroin	0.4	0.8	0.4	0.8	0.4	0.2	0.6	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Depressants	0.9	7.1	5.9	2.9	0.8	7.6	4.4	2.2	6.5	5.3	4.1	2.4	6.6	4.7
Prescription Pain Relievers	4.5	8.6	8.9	4.9	5.5	6.5	6.9	3.3	5.8	5.3	4.2	3.5	5.7	4.8
Prescription Amphetamines	2.5	3.4	3.6	2.4	2.3	4.1	3.0	1.3	4.7	3.3	3.2	1.4	4.6	3.2
Steroids (without a doctor's order)	0.5	0.0	0.4	0.0	0.5	0.0	0.2	0.6	0.5	0.4	0.6	0.5	0.5	0.5
Over-the-Counter Drugs	2.1	3.1	4.4	1.0	1.9	4.0	2.7	3.2	5.3	4.7	4.2	3.4	5.4	4.4
Needle to Inject Illegal Drugs		1.6	2.1	1.2		0.8	1.6		0.8	0.6	0.8		0.8	0.8
Any illicit drug	14.6	32.5	28.6	21.7	15.7	32.8	24.8	15.7	36.4	29.0	26.4	16.6	36.6	27.7
Any illicit drug other than marijuana	11.8	17.2	18.8	11.3	11.8	16.3	14.9	12.0	17.7	16.4	14.2	12.3	17.7	15.3
Alcohol only	15.0	26.2	16.2	25.8	16.8	21.7	21.5	12.5	21.0	18.1	17.0	13.4	21.0	17.5
Alcohol or any illicit drug	29.0	57.7	44.1	46.7	31.8	53.8	45.6	27.8	57.2	46.8	43.1	29.6	57.4	44.8
Any illicit drug, but no alcohol	5.0	6.4	6.5	5.2	5.0	6.5	5.8	6.5	6.1	6.0	6.5	6.5	6.1	6.3

Table 2. Percentages of Gilchrist and Florida Statewide youth who reported having used various drugs in their lifetimes, 2016

Note: The first 19 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. Respondents were asked on how many occasions in their lifetime they woke up after a night of drinking and did not remember the things they did or the places they went. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "---" indicates that data are not available.

Table 3. Percentages of Gilchrist and Florida Statewide youth who reported having used various drugs in the past 30 days,2016

				Gilchrist	t					Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Alcohol	7.4	31.8	18.1	24.5	9.0	30.5	21.5	8.3	25.5	19.9	17.0	9.1	25.2	18.3
Binge Drinking	2.4	15.1	6.6	12.6	3.3	13.5	9.8	3.2	10.9	7.9	7.7	3.4	10.8	7.7
Cigarettes	1.5	8.2	5.6	5.0	1.4	8.0	5.4	1.4	4.8	3.3	3.5	1.5	4.6	3.4
Vaporizer/E-Cigarette	6.7	18.7	9.7	17.1	6.3	17.4	13.6	5.1	12.8	8.4	10.6	5.5	12.9	9.6
Marijuana or Hashish	1.2	11.3	5.8	8.5	1.1	12.5	7.1	3.2	17.0	10.9	11.5	3.9	16.9	11.2
Synthetic Marijuana		3.4	1.0	5.9		4.3	3.4		1.0	0.9	1.2		1.1	1.0
Inhalants	2.9	1.1	3.3	0.4	2.2	1.8	1.8	2.2	1.2	1.9	1.4	2.1	1.2	1.6
Flakka		0.5	1.0	0.0		0.4	0.5		0.5	0.4	0.7		0.5	0.5
Club Drugs	0.0	0.5	0.6	0.0	0.0	0.4	0.3	0.3	0.8	0.5	0.6	0.3	0.7	0.6
LSD, PCP or Mushrooms	0.0	0.9	0.6	0.5	0.0	0.9	0.5	0.3	1.4	0.7	1.1	0.3	1.4	0.9
Methamphetamine	0.0	0.2	0.2	0.0	0.0	0.0	0.1	0.3	0.4	0.3	0.4	0.3	0.4	0.4
Cocaine or Crack Cocaine	0.0	2.0	0.7	1.6	0.0	1.7	1.1	0.3	0.8	0.6	0.6	0.3	0.8	0.6
Heroin	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.2	0.1	0.2	0.1	0.2	0.2
Depressants	0.0	3.8	2.7	1.7	0.5	4.1	2.2	0.8	2.4	2.0	1.5	0.9	2.5	1.8
Prescription Pain Relievers	1.7	3.5	3.8	1.7	1.6	4.0	2.8	1.6	2.0	2.2	1.5	1.6	2.1	1.8
Prescription Amphetamines	0.0	2.3	1.1	1.7	0.0	2.8	1.4	0.5	1.6	1.1	1.2	0.6	1.7	1.2
Steroids (without a doctor's order)	0.5	0.0	0.4	0.0	0.5	0.0	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Over-the-Counter Drugs	0.4	1.4	2.0	0.0	0.4	1.8	1.0	1.8	2.1	2.0	1.9	1.8	2.1	2.0
Any illicit drug	6.0	17.0	15.1	10.0	5.6	19.1	12.4	7.5	20.0	15.1	14.2	8.1	20.0	14.7
Any illicit drug other than marijuana	5.4	9.6	11.9	4.0	5.0	10.8	7.8	5.6	7.7	7.3	6.2	5.7	7.7	6.8
Alcohol only	5.9	21.8	11.5	18.2	7.7	19.0	15.2	5.2	13.4	11.1	9.1	5.6	13.3	10.0
Alcohol or any illicit drug	11.5	38.5	26.2	27.5	12.7	38.0	27.0	12.5	32.9	25.8	22.9	13.5	32.9	24.3
Any illicit drug, but no alcohol	4.6	7.3	8.7	3.7	4.3	8.0	6.2	4.5	7.8	6.4	6.3	4.7	8.0	6.4

Note: The first 18 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. Binge drinking is defined as having had five or more alcoholic drinks in a row in the past two weeks. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "--" indicates that data are not available.

		2006			2008			2010			2012			2014			2016	
	Middle School	0	Total	Middle School	.0.	Total	Middle School		Total	Middle School	High School	Total	Middle School	High School	Total	Middle School	High School	Total
Alcohol	46.0	69.3	58.8		65.4	65.4	39.6	67.0	54.5	39.6	69.6	56.2	34.8	65.2	51.6	24.2	51.5	40.1
Blacking Out														26.6	26.6		21.8	21.8
Cigarettes	27.9	49.6	40.0		47.0	47.0	21.8	46.5	35.1	22.7	49.2	37.3	16.9	35.1	27.0	9.8	25.7	19.0
Vaporizer/E-Cigarette																13.0	39.0	28.0
Marijuana or Hashish	13.4	34.0	24.6		29.4	29.4	12.6	37.2	26.0	10.5	37.2	25.4	10.3	31.8	22.3	5.6	25.5	17.0
Synthetic Marijuana											23.2	23.2		16.4	16.4		8.4	8.4
Inhalants	14.7	16.7	15.6		15.5	15.5	13.8	12.4	13.1	8.2	9.2	8.8	7.3	8.9	8.2	7.8	3.7	5.5
Flakka																	0.5	0.5
Club Drugs							0.0	3.3	1.8	1.2	5.9	3.8	0.8	3.5	2.3	0.0	2.6	1.5
LSD, PCP or Mushrooms							3.8	3.1	3.4	1.6	7.0	4.6	1.2	4.9	3.3	0.9	3.5	2.4
Methamphetamine	1.6	2.7	2.2		2.6	2.6	0.5	1.1	0.8	0.0	2.2	1.2	1.6	4.0	3.0	0.0	0.8	0.4
Cocaine or Crack Cocaine							1.3	3.1	2.3	1.2	5.5	3.6	1.9	2.9	2.4	0.9	4.6	3.0
Heroin	0.5	1.9	1.2		1.2	1.2	1.0	1.1	1.1	0.7	1.3	1.0	1.4	1.3	1.3	0.4	0.8	0.6
Depressants	3.0	11.8	7.7		9.7	9.7	3.1	6.2	4.8	2.8	7.3	5.3	0.7	6.7	4.0	0.9	7.1	4.4
Prescription Pain Relievers	10.2	16.2	13.3		14.5	14.5	6.9	10.2	8.7	6.7	15.0	11.4	2.7	6.7	5.0	4.5	8.6	6.9
Prescription Amphetamines	2.1	5.3	3.8		5.6	5.6	2.2	4.8	3.6	2.2	7.2	5.0	1.5	3.0	2.3	2.5	3.4	3.0
Steroids (without a doctor's order)	0.9	1.2	1.0		3.5	3.5	2.3	0.9	1.5	1.2	2.2	1.8	0.9	1.8	1.4	0.5	0.0	0.2
Over-the-Counter Drugs							5.3	10.8	8.3	2.6	7.4	5.2	3.6	6.1	5.0	2.1	3.1	2.7
Needle to Inject Drugs																	1.6	1.6
Any illicit drug	28.0	42.0	35.4		37.8	37.8	25.2	41.8	34.2	21.7	46.0	35.1	15.6	40.0	29.1	14.6	32.5	24.8
Any illicit drug other than marijuana	21.0	28.8	24.8		26.6	26.6	21.7	24.1	23.0	16.8	27.8	22.9	12.3	22.3	17.9	11.8	17.2	14.9
Alcohol only	25.7	30.7	28.7		30.4	30.4	20.2	27.7	24.3	21.6	26.9	24.6	22.4	30.4	26.8	15.0	26.2	21.5
Alcohol or any illicit drug	53.6	71.8	63.6		68.2	68.2	44.9	68.9	57.9	43.1	72.9	59.6	37.8	69.9	55.6	29.0	57.7	45.6
Any illicit drug, but no alcohol	8.2	3.5	5.6		2.8	2.8	5.6	1.9	3.6	3.5	3.3	3.4	2.8	5.3	4.2	5.0	6.4	5.8

Table 4. Lifetime trend in alcohol, tobacco and other drug use for Gilchrist, 2006 to 2016

Note: The first 19 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. Respondents were asked on how many occasions in their lifetime they woke up after a night of drinking and did not remember the things they did or the places they went. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "---" indicates that data are not available.

		2006			2008			2010			2012			2014			2016	
	Middle School		Total	Middle School	High School	Total	Middle School		Total	Middle School		Total	Middle School	0	Total	Middle School		Total
Alcohol	22.3	43.8	33.3		47.2	47.2	19.8	41.5	31.6	15.5	41.6	30.0	14.8	33.5	25.2	7.4	31.8	21.5
Binge Drinking	10.3	24.3	18.0		30.6	30.6	5.6	21.8	14.4	6.0	23.1	15.5	7.7	18.8	13.9	2.4	15.1	9.8
Cigarettes	11.2	22.6	17.1		20.7	20.7	9.0	15.3	12.4	6.4	20.4	14.1	3.5	16.3	10.6	1.5	8.2	5.4
Vaporizer/E-Cigarette																6.7	18.7	13.6
Marijuana or Hashish	6.6	16.5	11.8		15.6	15.6	6.7	17.6	12.6	6.2	19.9	13.9	4.1	13.7	9.5	1.2	11.3	7.1
Synthetic Marijuana											7.7	7.7		2.2	2.2		3.4	3.4
Inhalants	7.6	4.5	5.9		6.7	6.7	5.6	2.0	3.7	0.7	1.5	1.2	2.3	1.3	1.7	2.9	1.1	1.8
Flakka																	0.5	0.5
Club Drugs							0.0	0.5	0.3	0.4	1.1	0.8	0.0	1.2	0.7	0.0	0.5	0.3
LSD, PCP or Mushrooms							1.0	0.4	0.7	0.7	1.7	1.3	0.4	2.5	1.6	0.0	0.9	0.5
Methamphetamine	0.8	0.4	0.6		1.8	1.8	0.5	1.1	0.8	0.0	1.1	0.6	1.0	2.6	1.9	0.0	0.2	0.1
Cocaine or Crack Cocaine							0.0	0.8	0.4	0.7	1.7	1.3	0.4	1.2	0.9	0.0	2.0	1.1
Heroin	0.0	1.2	0.6		0.0	0.0	1.5	0.0	0.7	0.7	0.5	0.6	1.7	0.7	1.2	0.0	0.0	0.0
Depressants	0.8	6.9	4.0		4.8	4.8	0.5	1.6	1.1	0.7	2.8	1.9	0.4	2.5	1.6	0.0	3.8	2.2
Prescription Pain Relievers	4.7	5.1	4.9		6.8	6.8	1.8	3.3	2.7	2.0	4.6	3.4	0.9	2.5	1.8	1.7	3.5	2.8
Prescription Amphetamines	0.8	0.0	0.3		1.7	1.7	0.4	0.9	0.7	0.7	1.7	1.3	1.3	1.9	1.6	0.0	2.3	1.4
Steroids (without a doctor's order)	0.0	0.8	0.4		2.6	2.6	0.5	0.3	0.4	0.4	0.5	0.4	0.6	0.5	0.5	0.5	0.0	0.2
Over-the-Counter Drugs							2.8	4.3	3.6	1.8	3.9	3.0	1.5	3.6	2.6	0.4	1.4	1.0
Any illicit drug	16.2	20.6	18.3		19.8	19.8	15.1	20.3	17.9	9.7	25.4	18.4	7.3	17.3	12.8	6.0	17.0	12.4
Any illicit drug other than marijuana	12.5	11.7	11.9		14.1	14.1	10.1	8.9	9.5	5.5	10.6	8.4	6.6	8.4	7.6	5.4	9.6	7.8
Alcohol only	14.2	25.3	19.8		32.3	32.3	11.6	24.2	18.5	9.0	21.4	15.9	12.5	21.2	17.3	5.9	21.8	15.2
Alcohol or any illicit drug	30.0	45.4	37.7		50.8	50.8	26.8	43.8	36.0	18.5	46.9	34.2	19.6	38.0	29.8	11.5	38.5	27.0
Any illicit drug, but no alcohol	8.1	2.3	4.9		3.3	3.3	7.4	2.6	4.7	3.4	5.5	4.6	5.1	4.9	5.0	4.6	7.3	6.2

Table 5. Past-30-day trend in alcohol, tobacco and other drug use for Gilchrist, 2006 to 2016

Note: The first 18 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. Binge drinking is defined as having had five or more alcoholic drinks in a row in the past two weeks. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "--" indicates that data are not available.

Table 6. Percentages of Gilchrist and Florida Statewide youth who reported early ATOD use, perceived risk of harm, personal disapproval, parental disapproval and peer disapproval, 2016

			(Gilchrist	;					Flori	da State	ewide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Early ATOD use														
More than a sip of alcohol		24.8	23.5	24.8		24.9	24.8		19.4	18.3	20.2		19.3	19.4
Drinking at least once a month		7.5	6.4	7.8		7.5	7.5		3.5	3.7	3.4		3.5	3.5
Smoke cigarettes		16.8	16.1	16.2		16.5	16.8		9.3	8.6	9.8		9.4	9.3
Smoke marijuana		10.8	7.8	12.9		11.6	10.8		10.6	9.0	12.0		10.8	10.6
Great risk of harm														
1 or 2 drinks nearly every day	40.7	34.0	42.1	31.7	41.3	33.4	36.8	45.7	40.7	46.4	39.3	45.9	40.0	42.8
5+ drinks once or twice a week	53.2	49.2	62.1	39.9	53.5	50.8	50.9	58.1	52.3	58.4	51.2	58.4	51.8	54.7
1+ packs of cigarettes per day	67.2	58.8	64.7	60.2	65.5	58.5	62.3	67.9	68.8	68.4	68.5	68.7	68.2	68.4
Marijuana once or twice a week	55.8	32.9	45.1	39.0	53.1	34.6	42.4	51.7	25.2	37.5	35.2	50.1	24.4	36.3
Try marijuana once or twice	43.2	22.0	31.1	29.4	40.5	21.6	30.9	36.1	16.8	24.8	25.0	34.7	16.4	24.9
Use prescription drugs	70.5	66.3	71.2	65.1	72.5	64.3	68.1	69.1	68.2	70.0	67.3	69.4	67.9	68.5
Think it would be wrong to														
Smoke cigarettes	94.7	77.7	86.4	83.5	92.9	80.4	84.8	95.6	87.7	91.1	91.0	95.3	88.9	91.0
Drink alcohol regularly	84.6	57.1	74.7	61.9	79.5	59.6	68.6	88.8	64.5	74.1	75.2	87.6	64.8	74.7
Smoke marijuana	93.4	66.3	77.6	77.9	90.3	69.4	77.6	89.7	62.3	74.4	73.3	88.1	62.3	73.8
Smoke synthetic marijuana	96.1	86.4	88.3	92.6	94.7	86.5	90.4	93.1	88.6	90.1	90.9	92.7	88.6	90.5
Use other illicit drugs	98.7	93.8	96.7	94.9	97.5	95.7	95.9	97.5	93.6	95.8	94.7	97.3	93.6	95.2
Parents think it would be wrong to														
1 or 2 drinks nearly every day	98.2	90.9	94.6	93.1	98.4	91.3	94.0	97.3	93.8	95.4	95.1	97.2	94.1	95.2
Smoke cigarettes	99.1	92.6	95.3	95.2	98.3	93.1	95.3	98.6	96.8	97.6	97.5	98.6	97.3	97.6
Smoke marijuana	97.1	87.5	90.0	92.8	96.8	87.8	91.5	97.0	88.6	92.3	92.0	96.6	88.9	92.1
Use prescription drugs	97.7	95.9	95.2	98.1	97.7	95.4	96.6	98.2	97.3	97.4	97.9	98.1	97.4	97.7
Friends think it would be wrong to														
1 or 2 drinks nearly every day	92.7	72.7	87.0	75.5	92.2	73.8	81.0	91.3	79.0	85.6	82.9	90.7	79.1	84.2
Smoke tobacco	95.3	80.1	90.4	82.4	95.4	80.0	86.4	94.8	87.0	91.1	89.5	94.4	87.8	90.3
Smoke marijuana	92.5	68.5	80.3	76.7	90.4	69.0	78.5	88.6	59.9	72.6	71.3	86.6	59.8	72.0
Use prescription drugs	96.6	91.3	94.1	93.0	96.0	92.0	93.5	95.7	90.6	93.4	92.2	95.5	90.3	92.7

Note: Early ATOD use is defined as the percentage of high school students who started using at age 13 or younger. The symbol "--" indicates that data are not available.

		2006			2008			2010			2012			2014			2016	
	Middle School	0	Total	Middle School	0	Total	Middle School	High School	Total	Middle School	0	Total	Middle School	.0.	Total	Middle School	0	Total
Early ATOD use																		
Try alcohol		37.3	37.3		37.0	37.0		28.6	28.6		29.4	29.4		26.2	26.2		24.8	24.8
Drink monthly		6.5	6.5		9.7	9.7		11.5	11.5		7.3	7.3		9.5	9.5		7.5	7.5
Smoke cigarettes		33.8	33.8		31.2	31.2		24.9	24.9		28.0	28.0		18.9	18.9		16.8	16.8
Smoke marijuana		17.4	17.4		10.8	10.8		13.8	13.8		14.6	14.6		12.8	12.8		10.8	10.8
Great risk of harm																		
Daily alcohol use	34.9	35.9	35.3		34.1	34.1	40.5	36.1	38.6	42.2	30.3	35.6	38.8	29.8	33.8	40.7	34.0	36.8
5+ drinks weekly													48.0	42.4	44.9	53.2	49.2	50.9
Daily cigarette use	64.7	63.5	63.9		59.2	59.2	63.8	55.8	59.4	73.4	60.9	66.5	69.3	62.8	65.8	67.2	58.8	62.3
Marijuana weekly*													54.8	34.0	43.2	55.8	32.9	42.4
Try marijuana	48.3	25.9	36.1		27.7	27.7	36.5	23.6	29.4	40.7	20.2	29.4	39.0	26.3	32.0	43.2	22.0	30.9
Prescription drugs											71.1	71.1	74.9	75.9	75.4	70.5	66.3	68.1
Think it wrong																		
Smoke cigarettes	83.8	63.4	73.2		61.0	61.0	82.9	60.0	70.6	89.9	68.5	78.0	92.0	77.5	84.0	94.7	77.7	84.8
Drink regularly	71.8	51.3	60.1		46.2	46.2	73.6	55.9	64.0	75.2	49.2	60.8	82.6	59.4	69.8	84.6	57.1	68.6
Smoke marijuana	87.4	70.4	78.3		72.9	72.9	84.3	69.7	76.4	87.2	63.5	74.1	90.6	72.4	80.6	93.4	66.3	77.6
Smoke synthetic marijuana																96.1	86.4	90.4
Use other illicit drugs	97.4	93.6	95.4		91.7	91.7	92.7	93.2	93.0	96.9	93.1	94.8	96.7	96.3	96.5	98.7	93.8	95.9
Parents think it wrong																		
Daily alcohol use*													92.4	90.1	91.2	98.2	90.9	94.0
Smoke cigarettes	91.2	86.2	88.1		83.9	83.9	97.6	85.6	90.9	96.1	89.0	92.2	96.7	92.2	94.2	99.1	92.6	95.3
Smoke marijuana	95.8	91.3	93.3		86.6	86.6	96.2	90.5	93.0	95.8	90.3	92.8	97.6	89.3	93.0	97.1	87.5	91.5
Prescription drugs													97.6	97.5	97.6	97.7	95.9	96.6
Friends think it wrong																		
Daily alcohol use													85.0	69.1	76.2	92.7	72.7	81.0
Smoke tobacco													91.2	75.7	82.6	95.3	80.1	86.4
Smoke marijuana													92.7	68.3	79.2	92.5	68.5	78.5
Prescription drugs													97.0	92.9	94.7	96.6	91.3	93.5

Table 7. Trends in early ATOD use and attitudes toward substance use for Gilchrist, 2006 to 2016

Note: Early ATOD use is defined as the percentage of high school students who started using at age 13 or younger. The symbol "--" indicates that data are not available.

* In 2014, the description of marijuana use in the risk of harm question was changed from "regularly" to "once or twice a week." Also, in 2014, the description of alcohol use in the parent disapproval question was changed from "regularly" to "nearly every day." For these items, care should be exercised when comparing 2014 data to previous years.

Table 8. Among <u>high school</u> drinkers, usual source of alcohol within the past 30 days, Gilchrist and Florida Statewide youth,2016

				Gilchrist	t					Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Bought in a store		3.0	0.0	5.8		4.1	3.0		8.3	5.0	12.4		7.3	8.3
Bought in a restaurant, bar, or club		0.0	0.0	0.0		0.0	0.0		1.6	1.2	2.1		1.5	1.6
Bought at a public event		0.0	0.0	0.0		0.0	0.0		0.7	0.7	0.8		0.6	0.7
Someone bought it for me		21.0	20.8	21.3		24.9	21.0		14.7	13.3	16.3		14.5	14.7
Someone gave it to me		40.3	59.2	23.0		37.7	40.3		44.8	50.6	37.8		47.0	44.8
Took it from a store		0.0	0.0	0.0		0.0	0.0		0.3	0.2	0.4		0.4	0.3
Took it from a family member		7.9	6.5	9.2		6.3	7.9		11.7	14.2	8.7		12.4	11.7
Some other way		27.7	13.5	40.7		26.9	27.7		17.8	14.7	21.5		16.5	17.8

Note: The symbol "--" indicates that data are not available.

Table 9. Among <u>high school</u> drinkers, usual drinking location within the past 30 days, Gilchrist and Florida Statewide youth,2016

				Gilchrist	t					Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
My home		30.7	25.8	36.2		28.4	30.7		37.7	39.2	35.9		37.1	37.7
Another person's home		57.3	61.8	52.3		63.1	57.3		40.0	41.0	38.9		40.8	40.0
Car or other vehicle		0.8	0.0	1.6		0.0	0.8		1.7	1.6	1.9		1.7	1.7
Restaurant, bar, or club		0.8	1.4	0.0		1.0	0.8		2.7	2.9	2.5		2.4	2.7
Public place		1.6	3.0	0.0		2.2	1.6		3.9	3.9	3.7		4.3	3.9
Public event		1.7	0.0	3.7		0.0	1.7		1.6	1.8	1.5		1.7	1.6
School property		0.9	0.0	1.9		0.0	0.9		1.3	0.7	2.0		1.4	1.3
Some other place		6.3	8.0	4.3		5.4	6.3		11.1	9.0	13.7		10.6	11.1

Note: The symbol "--" indicates that data are not available.

Table 10. Among <u>high school</u> drinkers, number of drinks usually consumed per day, on the days students drank in the past 30 days, Gilchrist and Florida Statewide youth, 2016

			. (Gilchrist	t					Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
1		27.4	38.8	17.9		26.1	27.4		29.6	31.6	27.4		30.5	29.6
2		14.8	12.7	16.7		17.4	14.8		22.5	24.8	19.8		22.4	22.5
3		11.1	6.8	14.7		10.9	11.1		17.3	17.9	16.3		16.9	17.3
4		14.0	14.5	13.6		12.5	14.0		8.6	9.0	8.3		8.5	8.6
5 or more		32.6	27.2	37.1		33.0	32.6		22.0	16.7	28.2		21.8	22.0

Note: The symbol "--" indicates that data are not available.

another drug to get high <u>before c</u>	<u>)r duri</u>	ng scho	<u>ool</u> in th	ie past	12 mo	nths, 20	016							
			(Gilchrist	t					Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total

2.4

2.0

1.0

3.3

3.2

1.4

5.5

8.1

2.9

6.5

12.2

3.9

4.7

8.7

2.9

3.3

3.7

1.5

6.8

12.3

4.1

5.1

8.5

2.9

5.3

6.6

2.9

6.6

9.9

4.3

Table 11. Percentages of Gilchrist and Florida Statewide youth who reported drinking alcohol, smoking marijuana, or using another drug to get high <u>before or during school</u> in the past 12 months, 2016

2.6

2.1

1.1

7.1

9.8

4.3

4.1

4.9

2.5

6.6

8.5

3.4

Drinking alcohol

Smoking marijuana

Used another drug to get high

Table 12. Percentages of Gilchrist and Florida Statewide youth who reported engaging in delinquent behavior within the past12 months, 2016

				Gilchrist	t					Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Carrying a handgun	9.9	12.1	7.3	15.0	8.2	12.8	11.2	5.0	5.8	2.6	8.3	5.0	5.9	5.5
Selling drugs	1.1	5.1	2.6	4.4	1.7	4.4	3.4	1.6	6.0	3.0	5.4	1.8	6.1	4.2
Attempting to steal a vehicle	1.0	2.1	1.7	1.5	1.1	1.8	1.6	1.0	1.5	0.8	1.7	1.0	1.5	1.3
Being arrested	0.9	1.5	2.0	0.6	1.1	1.7	1.3	1.9	2.8	1.8	3.0	1.8	3.0	2.4
Taking a handgun to school	0.0	2.0	0.2	2.1	0.0	1.7	1.1	0.4	0.7	0.3	0.9	0.4	0.7	0.6
Getting suspended	10.6	12.9	8.4	15.8	9.4	14.6	11.9	10.6	9.1	7.0	12.5	10.1	9.8	9.8
Attacking someone with intent to harm	4.5	5.6	3.1	7.3	3.6	6.5	5.1	6.2	6.1	5.3	6.9	6.0	6.6	6.1

		2006			2008			2010			2012			2014			2016	
	Middle School	High School	Total															
Carrying a handgun	5.1	7.5	6.3		9.6	9.6	9.4	8.2	8.7	6.8	9.9	8.5	10.0	12.4	11.3	9.9	12.1	11.2
Selling drugs	0.8	4.9	3.1		8.7	8.7	1.9	9.1	5.9	2.9	10.9	7.3	3.1	6.2	4.9	1.1	5.1	3.4
Attempting to steal a vehicle	1.3	0.9	1.1		5.6	5.6	1.9	1.4	1.7	1.6	2.8	2.3	0.9	1.5	1.2	1.0	2.1	1.6
Being arrested	5.0	4.3	4.6		4.8	4.8	2.8	3.0	2.9	1.2	5.0	3.4	1.3	3.1	2.3	0.9	1.5	1.3
Taking a handgun to school	0.0	1.2	0.7		4.0	4.0	0.5	0.0	0.2	1.1	0.8	0.9	0.3	1.5	0.9	0.0	2.0	1.1
Getting suspended	17.2	13.7	15.0		15.6	15.6	15.5	14.4	14.9	11.5	13.3	12.5	11.4	6.7	8.8	10.6	12.9	11.9
Attacking someone with intent to harm	14.1	13.6	13.6		13.2	13.2	10.9	9.7	10.3	3.7	9.6	7.0	5.2	8.8	7.2	4.5	5.6	5.1

Table 13. Trends in delinquent behaviors for Gilchrist, 2006 to 2016

Table 14. Percentages of Gilchrist and Florida Statewide youth with family members who have been or are now in jail or friends who have been in trouble because of alcohol or drugs, 2016

			. (Gilchrist	t					Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Family members who have been or are now in jail or prison	50.0	47.1	52.8	43.8	47.5	46.0	48.3	44.4	44.6	47.8	41.2	44.2	45.4	44.5
Friends who have have been or are in trouble because of alcohol or drugs	23.2	36.4	31.5	30.4	23.6	34.2	30.9	23.9	46.5	38.4	35.8	26.0	47.1	37.1

 $\label{eq:2016-Florida-Youth-Substance-Abuse-Survey-Gilchrist Report} 2016\ Florida-Youth-Substance-Abuse-Survey-Gilchrist-Report}$

				Gilchrist						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Skipped school because of bullying	8.5	10.0	14.6	4.3	6.9	11.2	9.4	7.6	8.2	11.5	4.4	7.6	8.3	7.9
Was kicked or shoved in past 30 days	24.0	9.4	15.0	15.2	21.3	11.5	15.5	19.6	8.1	12.0	13.8	18.8	8.1	13.0
Was taunted or teased in past 30 days	34.6	25.8	35.5	23.8	35.1	25.8	29.5	35.3	22.5	30.1	25.6	34.4	22.8	27.9
Was victim of cyber bullying in past 30 days	10.9	11.9	16.4	6.8	12.0	11.4	11.5	7.8	7.6	10.2	5.2	7.8	7.9	7.7
Physically bullied others in past 30 days	10.0	5.8	4.7	9.6	9.2	5.2	7.6	7.6	4.9	5.0	7.0	7.2	5.2	6.1
Verbally bullied others in past 30 days	13.6	9.2	10.3	11.2	13.2	8.4	11.0	14.3	10.4	11.0	13.0	14.0	10.7	12.0
Cyber bullied others in past 30 days	5.0	4.3	5.0	3.8	4.5	3.5	4.6	3.6	4.1	4.0	3.7	3.7	4.1	3.9

Table 15. Percentages of Gilchrist and Florida Statewide youth who reported involvement in bullying behavior, 2016

				Gilchris	t					Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Have you ever belonged to a gang?	2.6	5.2	2.8	5.5	3.1	4.9	4.1	3.3	3.5	2.3	4.5	3.1	3.6	3.4
Did that gang have a name?	22.9	12.1	12.7	16.7	15.4	13.9	14.6	23.4	14.1	13.1	19.7	21.9	14.4	16.9
Are you a gang member now?		3.1	3.1	3.2		2.5	3.1		2.0	1.1	2.8		2.0	2.0

Table 16. Percentages of Gilchrist and Florida Statewide youth who reported gang membership, 2016

Note: The prevalence rates for "Did that gang have a name?" exclude students who reported that they have never belonged to a gang. The symbol "--" indicates that data are not available.

Table 17. Reasons for joining a gang, reported by <u>high school students who have belonged to a gang</u>, among Gilchrist and Florida Statewide youth, 2016

				Gilchrist	t					Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Fun and excitement		7.8	15.8	3.1		5.4	7.8		5.0	4.3	5.5		5.0	5.0
Protection		2.6	4.8	1.3		2.6	2.6		4.2	2.6	5.4		4.5	4.2
Friend or relative in a gang		4.5	0.0	7.4		6.5	4.5		4.1	2.9	4.9		4.1	4.1
Forced to join		2.7	0.0	4.5		4.0	2.7		1.1	1.1	1.2		1.2	1.1
To get respect		5.4	3.9	6.5		6.8	5.4		3.5	2.0	4.7		3.5	3.5
Money		2.2	0.0	3.6		3.2	2.2		3.8	2.5	4.8		4.1	3.8
To fit in better		3.0	2.0	3.8		3.3	3.0		1.2	1.1	1.4		1.2	1.2
Other reasons		7.5	0.0	12.4		10.9	7.5		5.9	4.5	7.0		5.6	5.9

Note: The symbol "--" indicates that data are not available.

 $\label{eq:2016-Florida-Youth-Substance-Abuse-Survey-Gilchrist Report} 2016\ Florida-Youth-Substance-Abuse-Survey-Gilchrist-Report}$

				Gilchrist						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
School Sports	54.6	48.5	46.6	56.0	53.9	50.4	51.1	36.0	40.4	34.5	42.5	36.4	41.4	38.5
Organized Sports Outside of School	32.5	15.4	21.8	24.0	32.5	15.1	22.6	41.6	22.4	28.2	32.9	40.1	22.8	30.5
School Band	25.9	13.2	21.0	16.3	26.0	13.6	18.6	16.7	8.1	12.3	11.1	16.2	8.1	11.7
School Club(s)	30.2	40.1	42.8	29.2	34.3	38.9	35.9	22.1	31.8	35.4	20.4	22.8	31.6	27.6
Community Club(s)	6.3	10.4	9.1	8.5	6.7	11.3	8.7	9.3	12.8	14.6	8.2	9.4	12.6	11.3

Table 18. Percentages of Gilchrist and Florida Statewide youth who reported participation in extracurricular activities, 2016

......

Table 19. Percentages of <u>high school</u> students who reported *riding in* a vehicle within the past 30 days driven by someone who had been drinking alcohol or using marijuana, or *driving* a vehicle within the past 30 days after drinking alcohol or using marijuana, among Gilchrist and Florida Statewide youth, 2016

				Gilchrist						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Riding in a vehicle														
Alcohol		24.4	28.7	19.2		23.0	24.4		16.4	17.5	15.3		16.3	16.4
Marijuana		16.2	16.8	15.6		17.4	16.2		22.7	23.7	21.7		22.9	22.7
Driving a vehicle														
Alcohol		9.0	8.8	9.0		7.2	9.0		5.4	5.0	5.8		5.2	5.4
Marijuana		8.3	6.1	10.0		7.8	8.3		10.3	9.1	11.4		9.9	10.3

Note: The symbol "--" indicates that data are not available.

Table 20. Trends for <u>high school</u> students *riding in* a vehicle within the past 30 days driven by someone who had been drinking alcohol or using marijuana, or *driving* a vehicle within the past 30 days after drinking alcohol or using marijuana, for Gilchrist, 2012 and 2016

		2006			2008			2010			2012			2014		2016	
	Middle School	High School	Total	Middle School	0	Total	High School	Total									
Riding in a vehicle																	
Alcohol											33.3	33.3		23.1	23.1	 24.4	24.4
Marijuana											24.0	24.0		17.2	17.2	 16.2	16.2
Driving a vehicle																	
Alcohol											14.3	14.3		11.1	11.1	 9.0	9.0
Marijuana											15.8	15.8		8.6	8.6	 8.3	8.3

Note: The symbol "--" indicates that data are not available.

Domain	Scale	Gilc	hrist	Florida S	Statewide	Nationa	l Norms
		Middle School	High School	Middle School	High School	Middle School	High School
Community	Community Rewards for Prosocial Involvement	58	80	46	69	56	63
Family	Family Opportunities for Prosocial Involvement Family Rewards for Prosocial Involvement	64 64	60 51	60 56	59 56	59 54	54 55
School	School Opportunities for Prosocial Involvement School Rewards for Prosocial Involvement	55 58	65 65	53 49	63 59	57 53	60 58
Peer and Individual	Religiosity	62	68	49	57	56	62
Average Prevalence R	ate	60	65	52	61	56	59

Table 21. Protective factor prevalence rates for Gilchrist, Florida Statewide youth and the national normative database, 2016

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.

Domain	Scale	Gilc	hrist	Florida S	Statewide	Nationa	l Norms
		Middle School	High School	Middle School	High School	Middle School	High School
Community	Community Disorganization	40	48	42	44	47	47
	Transitions and Mobility	56	49	59	61	47	46
	Laws and Norms Favorable to Drug Use	42	34	37	31	42	42
	Perceived Availability of Drugs	35	27	37	27	45	45
	Perceived Availability of Handguns	45	57	24	36	25	42
Family	Poor Family Management	32	44	40	38	44	45
	Family Conflict	31	35	38	33	42	37
School	Poor Academic Performance	41	34	42	44	45	48
	Lack of Commitment to School	58	53	53	54	47	46
Peer and	Favorable Attitudes toward Antisocial Behavior	35	39	39	35	40	46
Individual	Favorable Attitudes toward ATOD Use	36	42	32	36	39	45
	Early Initiation of Drug Use	26	27	23	22	41	46
Average Preva	lence Rate	40	41	39	38	42	45

Table 22. Risk factor prevalence rates for Gilchrist, Florida Statewide youth and the national normative database, 2016

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values. The symbol "---" indicates that data are not available.

Domain	Scale			Gilc	hrist		
		2006	2008	2010	2012	2014	2016
Community	Community Rewards for Prosocial Involvement	63		53	61	59	58
Family	Family Opportunities for Prosocial Involvement	54		53	61	58	64
	Family Rewards for Prosocial Involvement	50		43	52	53	64
School	School Opportunities for Prosocial Involvement	48		51	48	55	55
	School Rewards for Prosocial Involvement	44		44	59	56	58
Peer and Individual	Religiosity	56		54	59	59	62
Average Prevalence R	ate	53		50	57	57	60

Table 23. Protective factor prevalence rate trends among middle school students for Gilchrist, 2006 to 2016

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values. The symbol "---" indicates that data are not available.

Domain	Scale			Gilc	hrist		
		2006	2008	2010	2012	2014	2016
Community	Community Disorganization	42		45	48	53	40
	Transitions and Mobility	56		46	53	49	56
	Laws and Norms Favorable to Drug Use	55		52	45	46	42
	Perceived Availability of Drugs	51		52	48	41	35
	Perceived Availability of Handguns	45		40	44	45	45
Family	Poor Family Management	49		52	35	41	32
	Family Conflict	43		44	38	36	31
School	Poor Academic Performance	47		50	44	43	41
	Lack of Commitment to School	57		62	53	57	58
Peer and	Favorable Attitudes toward Antisocial Behavior	53		50	37	37	35
Individual	Favorable Attitudes toward ATOD Use	51		44	41	39	36
	Early Initiation of Drug Use	52		40	39	34	26
Average Preva	lence Rate	50		48	43	43	40

Table 24. Risk factor prevalence rate trends among middle school students for Gilchrist, 2006 to 2016

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.

Domain	Scale	Gilchrist						
		2006	2008	2010	2012	2014	2016	
Community	Community Rewards for Prosocial Involvement	72	65	69	72	76	80	
Family	Family Opportunities for Prosocial Involvement	58	54	61	54	56	60	
	Family Rewards for Prosocial Involvement	61	54	50	54	52	51	
School	School Opportunities for Prosocial Involvement	48	47	53	58	58	65	
	School Rewards for Prosocial Involvement	64	53	60	70	63	65	
Peer and Individual	Religiosity	68	61	65	63	67	68	
Average Prevalence Rate		62	56	60	62	62	65	

Table 25. Protective factor prevalence rate trends among high school students for Gilchrist, 2006 to 2016

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values. The symbol "---" indicates that data are not available.

Domain	Scale	Gilchrist						
		2006	2008	2010	2012	2014	2016	
Community	Community Disorganization	52	44	51	47	48	48	
	Transitions and Mobility	58	54	49	51	51	49	
	Laws and Norms Favorable to Drug Use	46	52	48	45	34	34	
	Perceived Availability of Drugs	43	43	46	37	31	27	
	Perceived Availability of Handguns	61	59	59	56	61	57	
Family	Poor Family Management	44	52	43	41	42	44	
	Family Conflict	34	33	35	35	34	35	
School	Poor Academic Performance	52	50	54	49	42	34	
	Lack of Commitment to School	54	62	61	53	60	53	
Peer and	Favorable Attitudes toward Antisocial Behavior	44	49	47	43	37	39	
Individual	Favorable Attitudes toward ATOD Use	46	45	45	47	36	42	
	Early Initiation of Drug Use	43	46	38	41	28	27	
Average Prevalence Rate		46	50	50	48	42	41	

Table 26. Risk factor prevalence rate trends among high school students for Gilchrist, 2006 to 2016

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.