2016 Florida Youth Substance Abuse Survey

Table 1. Major demographic characteristics of surveyed Hardee and Florida Statewide, 2016

	Hai	rdee	Florida S	tatewide
	N	%	N	%
Sex				
Female	392	47.3	31,515	47.9
Male	419	50.5	32,905	50.0
Race/Ethnic group				
African American	39	4.7	14,666	22.3
American Indian	20	2.4	568	0.9
Asian	11	1.3	743	1.1
Hispanic/Latino	446	53.8	13,174	20.0
Native Hawaiian/Pacific Islander	1	0.1	144	0.2
Other/Multiple	119	14.4	7,523	11.4
White, non-Hispanic	189	22.8	28,309	43.0
Age				
10	1	0.1	65	0.1
11	59	7.1	3,856	5.9
12	129	15.6	8,338	12.7
13	147	17.7	9,230	14.0
14	127	15.3	9,454	14.4
15	129	15.6	10,070	15.3
16	104	12.5	9,684	14.7
17	88	10.6	9,348	14.2
18	32	3.9	4,799	7.3
19 or older	8	1.0	618	0.9
Grade				
6th	157	18.9	9,301	14.1
7th	153	18.5	9,215	14.0
8th	152	18.3	9,326	14.2
9th	121	14.6	10,140	15.4
10th	118	14.2	9,834	15.0
11th	95	11.5	9,254	14.1
12th	33	4.0	8,705	13.2
Overall Middle School	462	55.7	27,678	42.1
Overall High School	367	44.3	37,765	57.4
Total	829	100.0	65,776	100.0

Note: Some categories do not sum to 100% of the total due to missing values (e.g., not all survey questions were answered). In addition, rounding can produce totals that do not equal 100%. "N" represents the number of valid cases. In this table, county data are unweighted while statewide data are weighted.

Table 2. Percentages of Hardee and Florida Statewide youth who reported having used various drugs in their lifetimes, 2016

				Hardee						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Alcohol	25.0	44.9	37.1	35.4	26.9	40.2	36.0	21.8	51.4	41.3	37.1	23.6	51.5	39.1
Blacking Out		12.8	12.3	13.1		12.6	12.8		15.9	16.5	15.4		16.1	15.9
Cigarettes	12.0	25.6	17.0	21.6	13.6	22.4	19.3	8.0	18.5	14.0	14.1	8.2	18.5	14.1
Vaporizer/E-Cigarette	24.2	43.3	29.9	39.4	26.0	38.7	34.5	14.6	33.9	24.4	27.1	15.7	34.5	25.8
Marijuana or Hashish	7.8	26.9	18.0	19.0	8.0	25.0	18.2	7.0	31.7	21.4	21.3	8.1	31.8	21.3
Synthetic Marijuana		12.1	11.7	12.9		11.7	12.1		4.9	4.8	5.0		4.8	4.9
Inhalants	6.3	6.6	5.3	7.8	7.5	4.2	6.5	6.4	4.6	6.1	4.7	6.4	4.7	5.4
Flakka		2.1	2.6	1.8		1.2	2.1		1.0	0.9	1.1		1.0	1.0
Club Drugs	1.0	3.7	2.4	2.7	1.9	2.9	2.5	0.7	3.0	1.9	2.2	0.8	2.9	2.1
LSD, PCP or Mushrooms	2.0	3.4	2.3	3.3	2.8	2.6	2.8	1.0	4.8	2.7	3.7	1.1	4.9	3.2
Methamphetamine	1.2	2.6	1.4	2.6	1.2	2.0	2.0	0.5	0.8	0.6	0.8	0.5	0.8	0.7
Cocaine or Crack Cocaine	1.8	3.6	2.2	3.4	2.7	2.1	2.8	0.8	2.5	1.6	1.9	0.8	2.3	1.8
Heroin	0.5	1.5	1.2	0.9	0.6	1.7	1.0	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Depressants	2.3	6.6	4.7	4.7	2.9	6.1	4.6	2.2	6.5	5.3	4.1	2.4	6.6	4.7
Prescription Pain Relievers	4.1	7.3	6.6	4.9	4.7	6.7	5.8	3.3	5.8	5.3	4.2	3.5	5.7	4.8
Prescription Amphetamines	1.5	3.4	2.6	2.6	1.9	2.1	2.6	1.3	4.7	3.3	3.2	1.4	4.6	3.2
Steroids (without a doctor's order)	1.2	0.3	1.1	0.3	1.0	0.5	0.7	0.6	0.5	0.4	0.6	0.5	0.5	0.5
Over-the-Counter Drugs	4.3	5.7	6.3	4.2	4.4	5.8	5.1	3.2	5.3	4.7	4.2	3.4	5.4	4.4
Needle to Inject Illegal Drugs		2.5	3.3	1.9		3.3	2.5		0.8	0.6	0.8		0.8	0.8
Any illicit drug	17.2	31.1	24.4	25.7	18.7	28.4	24.8	15.7	36.4	29.0	26.4	16.6	36.6	27.7
Any illicit drug other than marijuana	13.1	17.2	16.0	15.0	14.8	15.7	15.4	12.0	17.7	16.4	14.2	12.3	17.7	15.3
Alcohol only	13.5	20.2	18.2	16.4	14.2	19.2	17.2	12.5	21.0	18.1	17.0	13.4	21.0	17.5
Alcohol or any illicit drug	30.2	51.2	42.4	41.7	32.4	47.5	41.6	27.8	57.2	46.8	43.1	29.6	57.4	44.8
Any illicit drug, but no alcohol	5.7	6.5	5.6	6.9	6.3	7.1	6.1	6.5	6.1	6.0	6.5	6.5	6.1	6.3

Note: The first 19 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. Respondents were asked on how many occasions in their lifetime they woke up after a night of drinking and did not remember the things they did or the places they went. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "--" indicates that data are not available.

Table 3. Percentages of Hardee and Florida Statewide youth who reported having used various drugs in the past 30 days, 2016

				Hardee						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Alcohol	11.8	22.4	18.1	17.1	12.3	18.6	17.6	8.3	25.5	19.9	17.0	9.1	25.2	18.3
Binge Drinking	6.0	9.8	5.8	9.8	7.4	7.0	8.1	3.2	10.9	7.9	7.7	3.4	10.8	7.7
Cigarettes	1.8	7.4	4.1	5.0	2.6	4.8	4.8	1.4	4.8	3.3	3.5	1.5	4.6	3.4
Vaporizer/E-Cigarette	10.2	16.3	10.4	16.1	10.8	14.5	13.5	5.1	12.8	8.4	10.6	5.5	12.9	9.6
Marijuana or Hashish	2.9	11.8	7.0	8.7	2.9	11.2	7.7	3.2	17.0	10.9	11.5	3.9	16.9	11.2
Synthetic Marijuana		2.4	2.5	2.3		2.3	2.4		1.0	0.9	1.2		1.1	1.0
Inhalants	1.7	1.8	1.0	2.5	2.5	1.0	1.7	2.2	1.2	1.9	1.4	2.1	1.2	1.6
Flakka		0.6	1.1	0.0		0.3	0.6		0.5	0.4	0.7		0.5	0.5
Club Drugs	0.4	1.1	0.3	1.2	0.8	0.3	0.8	0.3	0.8	0.5	0.6	0.3	0.7	0.6
LSD, PCP or Mushrooms	0.7	1.3	0.3	1.8	0.8	1.0	1.1	0.3	1.4	0.7	1.1	0.3	1.4	0.9
Methamphetamine	0.5	0.8	0.5	0.9	0.5	0.9	0.7	0.3	0.4	0.3	0.4	0.3	0.4	0.4
Cocaine or Crack Cocaine	0.6	1.3	0.7	1.2	1.2	0.9	1.0	0.3	0.8	0.6	0.6	0.3	0.8	0.6
Heroin	0.4	0.8	0.5	0.7	0.2	1.2	0.6	0.1	0.2	0.1	0.2	0.1	0.2	0.2
Depressants	0.3	2.4	1.4	1.6	0.6	2.7	1.5	0.8	2.4	2.0	1.5	0.9	2.5	1.8
Prescription Pain Relievers	3.3	2.2	3.0	2.5	3.0	2.9	2.7	1.6	2.0	2.2	1.5	1.6	2.1	1.8
Prescription Amphetamines	0.3	0.7	0.7	0.4	0.3	0.8	0.5	0.5	1.6	1.1	1.2	0.6	1.7	1.2
Steroids (without a doctor's order)	0.7	0.0	0.5	0.2	0.6	0.0	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Over-the-Counter Drugs	2.1	2.7	1.9	3.0	2.3	2.0	2.4	1.8	2.1	2.0	1.9	1.8	2.1	2.0
Any illicit drug	9.4	16.6	12.4	14.6	10.2	15.9	13.3	7.5	20.0	15.1	14.2	8.1	20.0	14.7
Any illicit drug other than marijuana	7.5	8.5	7.4	9.0	8.5	7.5	8.1	5.6	7.7	7.3	6.2	5.7	7.7	6.8
Alcohol only	8.4	13.3	10.4	11.5	9.1	9.6	11.1	5.2	13.4	11.1	9.1	5.6	13.3	10.0
Alcohol or any illicit drug	17.7	29.5	22.9	25.6	19.2	25.5	24.2	12.5	32.9	25.8	22.9	13.5	32.9	24.3
Any illicit drug, but no alcohol	6.4	7.5	5.2	8.9	7.4	7.1	7.0	4.5	7.8	6.4	6.3	4.7	8.0	6.4

Note: The first 18 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. Binge drinking is defined as having had five or more alcoholic drinks in a row in the past two weeks. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "--" indicates that data are not available.

Table 4. Lifetime trend in alcohol, tobacco and other drug use for Hardee, 2006 to 2016

		2006			2008			2010			2012			2014			2016	
	Middle School	U	Total	Middle School	8	Total	Middle School		Total	Middle School	High School	Total	Middle School	O	Total	Middle School	U	Total
Alcohol	45.1	64.8	55.5	42.9	63.0	53.8	39.8	63.8	52.5	33.2	59.3	47.2	27.4	0.0	27.4	25.0	44.9	36.0
Blacking Out														0.0	0.0		12.8	12.8
Cigarettes	31.5	42.5	37.6	32.3	47.5	40.5	23.1	50.4	37.5	20.7	39.4	30.7	15.6	0.0	15.6	12.0	25.6	19.3
Vaporizer/E-Cigarette																24.2	43.3	34.5
Marijuana or Hashish	13.3	24.7	20.1	10.0	27.0	19.1	11.7	33.7	23.4	14.5	36.5	26.5	13.1	0.0	13.1	7.8	26.9	18.2
Synthetic Marijuana											10.4	10.4		0.0	0.0		12.1	12.1
Inhalants	15.2	8.4	11.4	12.2	8.2	10.0	13.6	8.4	10.8	9.3	2.7	5.7	8.0	0.0	8.0	6.3	6.6	6.5
Flakka																	2.1	2.1
Club Drugs				0.5		0.5	1.9	4.8	3.4	2.2	4.4	3.4	1.2	0.0	1.2	1.0	3.7	2.5
LSD, PCP or Mushrooms				0.5		0.5	2.1	3.8	3.0	2.7	6.3	4.7	1.1	0.0	1.1	2.0	3.4	2.8
Methamphetamine	6.5	6.0	6.5	0.9	5.9	3.6	2.9	3.9	3.4	2.2	5.4	3.9	1.6	0.0	1.6	1.2	2.6	2.0
Cocaine or Crack Cocaine				2.0		2.0	4.4	4.8	4.6	2.2	10.1	6.5	2.7	0.0	2.7	1.8	3.6	2.8
Heroin	1.2	0.8	1.0	0.5	1.0	0.7	1.7	1.5	1.6	1.1	0.0	0.5	1.0	0.0	1.0	0.5	1.5	1.0
Depressants	1.3	4.2	3.1	1.0	6.2	3.8	1.9	4.4	3.2	1.9	3.8	2.9	3.2	0.0	3.2	2.3	6.6	4.6
Prescription Pain Relievers	4.8	10.4	7.9	4.1	11.6	8.1	5.4	9.6	7.6	4.4	8.9	6.9	3.7	0.0	3.7	4.1	7.3	5.8
Prescription Amphetamines	0.6	3.3	2.3	0.7	1.9	1.4	0.8	2.5	1.7	1.8	3.1	2.5	1.1	0.0	1.1	1.5	3.4	2.6
Steroids (without a doctor's order)	1.7	1.4	1.5	0.9	0.3	0.6	0.7	0.7	0.7	0.2	0.0	0.1	0.9	0.0	0.9	1.2	0.3	0.7
Over-the-Counter Drugs				4.0		4.0	3.2	7.9	5.7	6.9	15.1	11.3	4.5	0.0	4.5	4.3	5.7	5.1
Needle to Inject Drugs																	2.5	2.5
Any illicit drug	25.7	32.4	29.8	21.3	33.9	28.1	23.9	39.8	32.2	24.7	43.7	35.0	21.1	0.0	21.1	17.2	31.1	24.8
Any illicit drug other than marijuana	21.1	20.1	20.8	16.2	22.4	19.5	18.7	20.9	19.9	18.3	29.1	24.2	13.7	0.0	13.7	13.1	17.2	15.4
Alcohol only	24.4	33.8	29.2	26.4	31.2	29.0	21.7	27.8	24.9	15.0	23.8	19.8	12.6	0.0	12.6	13.5	20.2	17.2
Alcohol or any illicit drug	49.9	66.1	58.7	47.5	64.9	56.9	45.3	67.6	57.0	39.7	66.8	54.3	33.9	0.0	33.9	30.2	51.2	41.6
Any illicit drug, but no alcohol	5.1	1.4	3.5	5.0	2.0	3.3	5.3	3.8	4.5	7.1	6.4	6.7	6.4	0.0	6.4	5.7	6.5	6.1

Note: The first 19 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. Respondents were asked on how many occasions in their lifetime they woke up after a night of drinking and did not remember the things they did or the places they went. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "--" indicates that data are not available.

2016 Florida Youth Substance Abuse Survey - Hardee Report

Table 5. Past-30-day trend in alcohol, tobacco and other drug use for Hardee, 2006 to 2016

		2006			2008			2010			2012			2014			2016	
	Middle School	High School	Total	Middle School	0	Total	Middle School	0	Total	Middle School		Total	Middle School	High School	Total	Middle School	0	Total
Alcohol	20.4	44.7	33.0	20.0	36.8	29.1	22.0	34.8	28.7	15.7	36.9	27.2	13.0	0.0	13.0	11.8	22.4	17.6
Binge Drinking	12.9	26.0	19.7	9.6	22.0	16.3	9.1	19.6	14.6	10.0	22.0	16.5	8.1	0.0	8.1	6.0	9.8	8.1
Cigarettes	9.5	13.3	11.6	8.5	18.1	13.7	6.9	17.6	12.5	5.4	24.2	15.5	5.0	0.0	5.0	1.8	7.4	4.8
Vaporizer/E-Cigarette																10.2	16.3	13.5
Marijuana or Hashish	6.1	11.6	9.9	4.9	11.8	8.6	5.4	16.3	11.2	7.2	16.2	12.2	7.4	0.0	7.4	2.9	11.8	7.7
Synthetic Marijuana											6.1	6.1		0.0	0.0		2.4	2.4
Inhalants	6.4	1.7	3.9	6.0	1.8	3.8	7.3	3.2	5.1	3.7	0.9	2.2	4.0	0.0	4.0	1.7	1.8	1.7
Flakka																	0.6	0.6
Club Drugs				0.0		0.0	1.5	1.7	1.6	0.3	3.8	2.2	0.4	0.0	0.4	0.4	1.1	0.8
LSD, PCP or Mushrooms				0.3		0.3	1.1	1.2	1.2	0.3	5.0	2.9	0.0	0.0	0.0	0.7	1.3	1.1
Methamphetamine	2.4	3.5	3.3	0.3	1.6	1.0	0.7	0.9	0.8	0.6	5.3	3.2	1.1	0.0	1.1	0.5	0.8	0.7
Cocaine or Crack Cocaine				0.8		0.8	1.5	0.8	1.1	0.6	5.0	3.0	0.9	0.0	0.9	0.6	1.3	1.0
Heroin	0.8	0.4	0.6	0.0	0.3	0.2	1.2	0.0	0.6	0.3	0.0	0.1	0.9	0.0	0.9	0.4	0.8	0.6
Depressants	0.9	2.4	2.0	0.3	2.7	1.6	1.0	1.1	1.0	0.5	0.2	0.4	0.8	0.0	0.8	0.3	2.4	1.5
Prescription Pain Relievers	1.6	4.5	3.4	1.6	4.4	3.1	3.0	2.9	2.9	1.7	0.2	0.9	2.7	0.0	2.7	3.3	2.2	2.7
Prescription Amphetamines	0.3	1.7	1.4	0.4	0.9	0.7	0.2	0.8	0.5	0.8	0.0	0.3	1.3	0.0	1.3	0.3	0.7	0.5
Steroids (without a doctor's order)	1.5	1.4	1.4	0.4	0.0	0.2	0.0	0.7	0.4	0.3	0.0	0.1	0.2	0.0	0.2	0.7	0.0	0.3
Over-the-Counter Drugs				2.0		2.0	1.3	3.5	2.5	3.0	12.6	8.3	2.5	0.0	2.5	2.1	2.7	2.4
Any illicit drug	14.0	14.6	15.2	11.7	14.9	13.4	14.0	21.3	17.8	12.4	22.1	17.7	13.4	0.0	13.4	9.4	16.6	13.3
Any illicit drug other than marijuana	11.1	7.9	9.9	8.2	8.8	8.5	11.4	10.1	10.7	8.2	18.6	13.9	8.0	0.0	8.0	7.5	8.5	8.1
Alcohol only	11.3	33.3	22.3	15.2	24.5	20.3	14.3	21.5	18.1	7.9	25.2	17.4	6.6	0.0	6.6	8.4	13.3	11.1
Alcohol or any illicit drug	24.8	46.7	36.7	26.4	39.5	33.5	27.8	42.5	35.5	20.4	47.2	34.9	19.7	0.0	19.7	17.7	29.5	24.2
Any illicit drug, but no alcohol	4.7	3.0	4.2	6.8	2.9	4.6	5.9	8.1	7.1	4.7	10.7	8.0	7.0	0.0	7.0	6.4	7.5	7.0

Note: The first 18 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. Binge drinking is defined as having had five or more alcoholic drinks in a row in the past two weeks. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "--" indicates that data are not available.

Table 6. Percentages of Hardee and Florida Statewide youth who reported early ATOD use, perceived risk of harm, personal disapproval, parental disapproval and peer disapproval, 2016

				Hardee						Flori	da State	ewide		
	Middle School		Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Early ATOD use														
More than a sip of alcohol		21.0	17.3	24.6		20.3	21.0		19.4	18.3	20.2		19.3	19.4
Drinking at least once a month		6.0	3.9	7.5		4.7	6.0		3.5	3.7	3.4		3.5	3.5
Smoke cigarettes		15.1	11.6	18.3		14.2	15.1		9.3	8.6	9.8		9.4	9.3
Smoke marijuana		14.6	10.0	19.5		13.8	14.6		10.6	9.0	12.0		10.8	10.6
Great risk of harm														
1 or 2 drinks nearly every day	42.8	34.7	43.5	32.7	41.8	35.9	38.4	45.7	40.7	46.4	39.3	45.9	40.0	42.8
5+ drinks once or twice a week	53.4	47.7	53.6	46.7	52.4	49.4	50.3	58.1	52.3	58.4	51.2	58.4	51.8	54.7
1+ packs of cigarettes per day	63.7	53.5	57.1	58.7	62.5	56.3	58.2	67.9	68.8	68.4	68.5	68.7	68.2	68.4
Marijuana once or twice a week	51.9	27.5	41.7	34.8	50.2	29.2	38.7	51.7	25.2	37.5	35.2	50.1	24.4	36.3
Try marijuana once or twice	40.5	24.2	32.8	29.3	38.1	26.9	31.7	36.1	16.8	24.8	25.0	34.7	16.4	24.9
Use prescription drugs	69.5	61.7	66.3	63.9	67.6	64.6	65.3	69.1	68.2	70.0	67.3	69.4	67.9	68.5
Think it would be wrong to														
Smoke cigarettes	94.2	81.4	87.0	87.4	91.7	84.8	87.2	95.6	87.7	91.1	91.0	95.3	88.9	91.0
Drink alcohol regularly	84.5	63.8	73.5	72.8	82.3	66.9	73.2	88.8	64.5	74.1	75.2	87.6	64.8	74.7
Smoke marijuana	89.9	68.6	81.0	75.3	87.4	70.4	78.3	89.7	62.3	74.4	73.3	88.1	62.3	73.8
Smoke synthetic marijuana	91.5	89.1	91.4	89.0	90.7	88.7	90.2	93.1	88.6	90.1	90.9	92.7	88.6	90.5
Use other illicit drugs	96.6	92.2	95.0	93.6	95.8	93.0	94.2	97.5	93.6	95.8	94.7	97.3	93.6	95.2
Parents think it would be wrong to														
1 or 2 drinks nearly every day	96.7	94.6	95.6	95.4	97.2	94.9	95.6	97.3	93.8	95.4	95.1	97.2	94.1	95.2
Smoke cigarettes	98.4	96.6	97.5	97.3	98.7	97.6	97.4	98.6	96.8	97.6	97.5	98.6	97.3	97.6
Smoke marijuana	96.4	93.4	95.6	93.7	96.4	94.1	94.8	97.0	88.6	92.3	92.0	96.6	88.9	92.1
Use prescription drugs	95.6	96.7	96.1	96.2	95.5	96.8	96.2	98.2	97.3	97.4	97.9	98.1	97.4	97.7
Friends think it would be wrong to														
1 or 2 drinks nearly every day	88.1	75.9	83.0	80.3	86.7	77.4	81.5	91.3	79.0	85.6	82.9	90.7	79.1	84.2
Smoke tobacco	92.5	85.8	92.3	85.1	91.7	87.3	88.8	94.8	87.0	91.1	89.5	94.4	87.8	90.3
Smoke marijuana	86.8	65.7	78.2	72.4	84.0	68.3	75.4	88.6	59.9	72.6	71.3	86.6	59.8	72.0
Use prescription drugs	94.9	90.1	94.5	90.3	93.7	90.7	92.3	95.7	90.6	93.4	92.2	95.5	90.3	92.7

Note: Early ATOD use is defined as the percentage of high school students who started using at age 13 or younger. The symbol "--" indicates that data are not available.

2016 Florida Youth Substance Abuse Survey - Hardee Report

Table 7. Trends in early ATOD use and attitudes toward substance use for Hardee, 2006 to 2016

		2006			2008			2010			2012			2014			2016	
	Middle School	High School	Total	Middle School		Total	Middle School		Total	Middle School		Total	Middle School	High School	Total	Middle School	High School	Total
Early ATOD use																		
Try alcohol		27.3	27.3		30.6	30.6		31.9	31.9		33.1	33.1		0.0	0.0		21.0	21.0
Drink monthly		8.8	8.8		7.6	7.6		8.4	8.4		14.4	14.4		0.0	0.0		6.0	6.0
Smoke cigarettes		28.8	28.8		28.7	28.7		32.0	32.0		34.2	34.2		0.0	0.0		15.1	15.1
Smoke marijuana		12.7	12.7		11.2	11.2		14.0	14.0		7.8	7.8		0.0	0.0		14.6	14.6
Great risk of harm																		
Daily alcohol use	34.3	36.9	36.1	37.2	33.0	34.9	38.6	32.0	35.1	46.1	38.3	41.9	44.0	0.0	44.0	42.8	34.7	38.4
5+ drinks weekly													52.9	0.0	52.9	53.4	47.7	50.3
Daily cigarette use	55.8	58.1	57.4	54.8	55.3	55.1	57.0	53.3	55.0	57.7	73.3	66.1	62.9	0.0	62.9	63.7	53.5	58.2
Marijuana weekly*													50.6	0.0	50.6	51.9	27.5	38.7
Try marijuana	44.6	34.7	39.4	41.1	28.1	34.1	39.7	25.8	32.4	40.2	29.9	34.6	40.5	0.0	40.5	40.5	24.2	31.7
Prescription drugs											82.2	82.2	70.0	0.0	70.0	69.5	61.7	65.3
Think it wrong																		
Smoke cigarettes	84.1	67.8	75.6	87.4	63.9	74.7	86.2	68.9	77.0	90.7	76.3	83.0	91.3	0.0	91.3	94.2	81.4	87.2
Drink regularly	76.2	55.4	65.3	79.5	56.6	67.1	77.8	57.2	66.9	81.7	63.8	72.1	82.9	0.0	82.9	84.5	63.8	73.2
Smoke marijuana	89.6	75.1	81.9	92.0	75.8	83.3	91.1	71.3	80.6	86.1	83.2	84.5	85.1	0.0	85.1	89.9	68.6	78.3
Smoke synthetic marijuana																91.5	89.1	90.2
Use other illicit drugs	94.8	93.3	93.8	97.2	93.4	95.2	96.0	93.7	94.8	96.0	97.6	96.8	94.9	0.0	94.9	96.6	92.2	94.2
Parents think it wrong																		
Daily alcohol use*													95.7	0.0	95.7	96.7	94.6	95.6
Smoke cigarettes	93.1	92.2	92.4	94.7	90.4	92.4	97.9	89.2	93.4	98.2	96.4	97.2	97.4	0.0	97.4	98.4	96.6	97.4
Smoke marijuana	96.5	93.2	94.5	97.0	95.3	96.1	97.9	95.0	96.4	96.9	95.1	96.0	96.3	0.0	96.3	96.4	93.4	94.8
Prescription drugs													97.8	0.0	97.8	95.6	96.7	96.2
Friends think it wrong																		
Daily alcohol use													82.2	0.0	82.2	88.1	75.9	81.5
Smoke tobacco													90.8	0.0	90.8	92.5	85.8	88.8
Smoke marijuana													82.8	0.0	82.8	86.8	65.7	75.4
Prescription drugs													93.0	0.0	93.0	94.9	90.1	92.3

Note: Early ATOD use is defined as the percentage of high school students who started using at age 13 or younger. The symbol "--" indicates that data are not available.

^{*} In 2014, the description of marijuana use in the risk of harm question was changed from "regularly" to "once or twice a week." Also, in 2014, the description of alcohol use in the parent disapproval question was changed from "regularly" to "nearly every day." For these items, care should be exercised when comparing 2014 data to previous years.

Table 8. Among $\underline{\text{high school}}$ drinkers, usual source of alcohol within the past 30 days, Hardee and Florida Statewide youth, 2016

				Hardee						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Bought in a store		3.3	0.0	7.7		2.1	3.3		8.3	5.0	12.4		7.3	8.3
Bought in a restaurant, bar, or club		1.3	0.0	3.0		2.2	1.3		1.6	1.2	2.1		1.5	1.6
Bought at a public event		0.0	0.0	0.0		0.0	0.0		0.7	0.7	0.8		0.6	0.7
Someone bought it for me		20.5	4.9	41.5		15.3	20.5		14.7	13.3	16.3		14.5	14.7
Someone gave it to me		47.2	65.5	20.5		37.9	47.2		44.8	50.6	37.8		47.0	44.8
Took it from a store		0.0	0.0	0.0		0.0	0.0		0.3	0.2	0.4		0.4	0.3
Took it from a family member		11.6	12.9	10.9		19.4	11.6		11.7	14.2	8.7		12.4	11.7
Some other way		16.1	16.7	16.5		23.1	16.1		17.8	14.7	21.5		16.5	17.8

Table 9. Among $\underline{\text{high school}}$ drinkers, usual drinking location within the past 30 days, Hardee and Florida Statewide youth, 2016

				Hardee						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
My home		35.9	35.8	38.6		24.4	35.9		37.7	39.2	35.9		37.1	37.7
Another person's home		25.4	26.2	22.9		31.9	25.4		40.0	41.0	38.9		40.8	40.0
Car or other vehicle		5.2	10.0	0.0		9.3	5.2		1.7	1.6	1.9		1.7	1.7
Restaurant, bar, or club		8.3	4.7	13.0		7.3	8.3		2.7	2.9	2.5		2.4	2.7
Public place		2.6	1.9	3.6		1.7	2.6		3.9	3.9	3.7		4.3	3.9
Public event		0.9	0.0	2.0		1.6	0.9		1.6	1.8	1.5		1.7	1.6
School property		1.7	0.0	3.9		3.1	1.7		1.3	0.7	2.0		1.4	1.3
Some other place		20.0	21.4	16.0		20.5	20.0		11.1	9.0	13.7		10.6	11.1

Table 10. Among <u>high school</u> drinkers, number of drinks usually consumed per day, on the days students drank in the past 30 days, Hardee and Florida Statewide youth, 2016

				Hardee						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
1		30.8	43.9	13.8		38.9	30.8		29.6	31.6	27.4		30.5	29.6
2		13.0	9.7	17.7		3.9	13.0		22.5	24.8	19.8		22.4	22.5
3		22.9	29.4	13.5		23.3	22.9		17.3	17.9	16.3		16.9	17.3
4		13.3	1.8	27.7		14.6	13.3		8.6	9.0	8.3		8.5	8.6
5 or more		20.0	15.1	27.3		19.3	20.0		22.0	16.7	28.2		21.8	22.0

Table 11. Percentages of Hardee and Florida Statewide youth who reported drinking alcohol, smoking marijuana, or using another drug to get high $\underline{\text{before or during school}}$ in the past 12 months, 2016

				Hardee						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Drinking alcohol	6.4	7.3	7.9	5.8	7.1	7.2	6.9	3.3	6.5	5.5	4.7	3.3	6.8	5.1
Smoking marijuana	5.1	11.4	8.0	9.3	4.8	11.2	8.5	3.2	12.2	8.1	8.7	3.7	12.3	8.5
Used another drug to get high	1.6	2.8	2.1	2.6	1.8	3.2	2.3	1.4	3.9	2.9	2.9	1.5	4.1	2.9

Table 12. Percentages of Hardee and Florida Statewide youth who reported engaging in delinquent behavior within the past 12 months, 2016

				Hardee						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Carrying a handgun	9.3	8.8	4.5	13.4	8.8	7.9	9.0	5.0	5.8	2.6	8.3	5.0	5.9	5.5
Selling drugs	2.1	8.0	3.6	7.1	2.8	6.7	5.3	1.6	6.0	3.0	5.4	1.8	6.1	4.2
Attempting to steal a vehicle	0.6	2.8	0.8	2.9	1.1	2.4	1.8	1.0	1.5	0.8	1.7	1.0	1.5	1.3
Being arrested	1.4	3.3	1.0	3.8	1.4	3.0	2.4	1.9	2.8	1.8	3.0	1.8	3.0	2.4
Taking a handgun to school	0.7	1.1	0.3	1.6	0.7	0.5	0.9	0.4	0.7	0.3	0.9	0.4	0.7	0.6
Getting suspended	11.9	5.4	5.7	10.2	12.0	4.4	8.4	10.6	9.1	7.0	12.5	10.1	9.8	9.8
Attacking someone with intent to harm	5.7	5.7	5.1	6.3	6.8	4.9	5.7	6.2	6.1	5.3	6.9	6.0	6.6	6.1

Table 13. Trends in delinquent behaviors for Hardee, 2006 to 2016

		2006			2008			2010			2012			2014			2016	
	Middle School	High School	Total															
Carrying a handgun	5.8	8.6	7.6	5.0	9.0	7.1	7.8	6.9	7.3	5.7	4.1	4.8	5.2	0.0	5.2	9.3	8.8	9.0
Selling drugs	4.9	5.3	5.4	2.9	6.3	4.7	4.2	7.4	5.9	3.8	9.5	6.8	2.0	0.0	2.0	2.1	8.0	5.3
Attempting to steal a vehicle	2.2	5.3	4.2	2.0	1.4	1.7	3.0	2.2	2.6	1.3	1.2	1.3	0.9	0.0	0.9	0.6	2.8	1.8
Being arrested	4.8	4.8	4.7	2.8	6.7	4.9	3.3	4.2	3.7	4.3	1.4	2.7	1.8	0.0	1.8	1.4	3.3	2.4
Taking a handgun to school	1.6	3.0	2.7	1.2	1.7	1.5	1.1	2.1	1.7	0.3	0.7	0.5	0.3	0.0	0.3	0.7	1.1	0.9
Getting suspended	16.5	11.1	13.3	17.3	9.7	13.3	12.8	9.8	11.3	11.2	4.2	7.4	10.3	0.0	10.3	11.9	5.4	8.4
Attacking someone with intent to harm	15.9	12.9	14.5	11.8	11.3	11.5	9.4	11.9	10.7	6.2	7.6	6.9	6.7	0.0	6.7	5.7	5.7	5.7

Table 14. Percentages of Hardee and Florida Statewide youth with family members who have been or are now in jail or friends who have been in trouble because of alcohol or drugs, 2016

				Hardee						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Family members who have been or are now in jail or prison	53.4	52.4	56.8	48.5	54.1	53.9	52.9	44.4	44.6	47.8	41.2	44.2	45.4	44.5
Friends who have have been or are in trouble because of alcohol or drugs	32.0	48.8	43.5	38.8	34.1	47.4	41.1	23.9	46.5	38.4	35.8	26.0	47.1	37.1

Table 15. Percentages of Hardee and Florida Statewide youth who reported involvement in bullying behavior, 2016

				Hardee						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Skipped school because of bullying	6.1	9.3	11.1	5.0	7.7	9.0	7.8	7.6	8.2	11.5	4.4	7.6	8.3	7.9
Was kicked or shoved in past 30 days	21.4	10.2	13.9	16.7	21.1	10.1	15.3	19.6	8.1	12.0	13.8	18.8	8.1	13.0
Was taunted or teased in past 30 days	34.9	24.5	30.9	27.6	35.0	25.2	29.3	35.3	22.5	30.1	25.6	34.4	22.8	27.9
Was victim of cyber bullying in past 30 days	5.2	7.3	8.5	4.6	4.8	8.1	6.4	7.8	7.6	10.2	5.2	7.8	7.9	7.7
Physically bullied others in past 30 days	11.2	6.3	6.0	11.1	11.7	5.6	8.5	7.6	4.9	5.0	7.0	7.2	5.2	6.1
Verbally bullied others in past 30 days	18.6	11.6	14.2	15.7	17.7	12.4	14.8	14.3	10.4	11.0	13.0	14.0	10.7	12.0
Cyber bullied others in past 30 days	3.5	3.2	4.0	2.8	3.5	3.5	3.3	3.6	4.1	4.0	3.7	3.7	4.1	3.9

Table 16. Percentages of Hardee and Florida Statewide youth who reported gang membership, 2016

				Hardee						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Have you ever belonged to a gang?	5.7	5.7	5.6	6.0	4.2	7.7	5.7	3.3	3.5	2.3	4.5	3.1	3.6	3.4
Did that gang have a name?	33.4	12.1	14.0	22.0	28.3	14.3	18.2	23.4	14.1	13.1	19.7	21.9	14.4	16.9
Are you a gang member now?		4.2	5.1	3.4		4.4	4.2		2.0	1.1	2.8		2.0	2.0

Note: The prevalence rates for "Did that gang have a name?" exclude students who reported that they have never belonged to a gang. The symbol "--" indicates that data are not available.

Table 17. Reasons for joining a gang, reported by <u>high school students who have belonged to a gang</u>, among Hardee and Florida Statewide youth, 2016

				Hardee						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Fun and excitement		8.0	11.5	6.3		9.9	8.0		5.0	4.3	5.5		5.0	5.0
Protection		12.2	0.0	17.9		12.0	12.2		4.2	2.6	5.4		4.5	4.2
Friend or relative in a gang		6.2	4.8	6.9		5.8	6.2		4.1	2.9	4.9		4.1	4.1
Forced to join		0.9	0.0	1.3		1.1	0.9		1.1	1.1	1.2		1.2	1.1
To get respect		9.5	0.0	14.1		7.0	9.5		3.5	2.0	4.7		3.5	3.5
Money		14.1	4.0	18.8		13.3	14.1		3.8	2.5	4.8		4.1	3.8
To fit in better		0.0	0.0	0.0		0.0	0.0		1.2	1.1	1.4		1.2	1.2
Other reasons		12.2	6.9	14.8		13.8	12.2		5.9	4.5	7.0		5.6	5.9

Table 18. Percentages of Hardee and Florida Statewide youth who reported participation in extracurricular activities, 2016

				Hardee						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
School Sports	34.3	41.7	34.7	42.3	33.3	43.6	38.4	36.0	40.4	34.5	42.5	36.4	41.4	38.5
Organized Sports Outside of School	39.0	23.0	28.8	31.6	37.0	23.7	30.3	41.6	22.4	28.2	32.9	40.1	22.8	30.5
School Band	10.9	7.3	10.9	6.7	12.6	6.1	8.9	16.7	8.1	12.3	11.1	16.2	8.1	11.7
School Club(s)	18.0	23.5	29.3	13.5	18.5	23.8	21.0	22.1	31.8	35.4	20.4	22.8	31.6	27.6
Community Club(s)	11.6	11.9	16.0	7.7	12.3	12.5	11.7	9.3	12.8	14.6	8.2	9.4	12.6	11.3

Table 19. Percentages of <u>high school</u> students who reported *riding in* a vehicle within the past 30 days driven by someone who had been drinking alcohol or using marijuana, or *driving* a vehicle within the past 30 days after drinking alcohol or using marijuana, among Hardee and Florida Statewide youth, 2016

				Hardee						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Riding in a vehicle														
Alcohol		17.3	18.4	16.9		16.0	17.3		16.4	17.5	15.3		16.3	16.4
Marijuana		17.3	15.7	19.4		16.9	17.3		22.7	23.7	21.7		22.9	22.7
Driving a vehicle														
Alcohol		7.6	7.0	7.9		6.3	7.6		5.4	5.0	5.8		5.2	5.4
Marijuana		8.9	6.9	11.2		8.3	8.9		10.3	9.1	11.4		9.9	10.3

Table 20. Trends for <u>high school</u> students *riding in* a vehicle within the past 30 days driven by someone who had been drinking alcohol or using marijuana, or *driving* a vehicle within the past 30 days after drinking alcohol or using marijuana, for Hardee, 2012 and 2016

		2006			2008			2010			2012			2014			2016	
	Middle School	High School	Total	Middle School	O	Total	Middle School	High School	Total	Middle School	0		Middle School	High School	Total	Middle School	High School	Total
Riding in a vehicle																		
Alcohol											23.6	23.6		0.0	0.0		17.3	17.3
Marijuana											17.9	17.9		0.0	0.0		17.3	17.3
Driving a vehicle																		
Alcohol											10.2	10.2		0.0	0.0		7.6	7.6
Marijuana											9.7	9.7		0.0	0.0		8.9	8.9

Table 21. Protective factor prevalence rates for Hardee, Florida Statewide youth and the national normative database, 2016

Domain	Scale	Hai	dee	Florida S	Statewide	Nationa	l Norms
		Middle School	High School	Middle School	High School	Middle School	High School
Community	Community Rewards for Prosocial Involvement	45	66	46	69	56	63
Family	Family Opportunities for Prosocial Involvement	55	51	60	59	59	54
School	Family Rewards for Prosocial Involvement School Opportunities for Prosocial Involvement	50 35	46 57	56 53	56 63	54	55 60
	School Rewards for Prosocial Involvement	42	54	49	59	53	58
Peer and Individual	Religiosity	57	70	49	57	56	62
Average Prevalence R	ate	47	57	52	61	56	59

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.

Table 22. Risk factor prevalence rates for Hardee, Florida Statewide youth and the national normative database, 2016

Domain	Scale	Hai	rdee	Florida S	Statewide	Nationa	l Norms
		Middle School	High School	Middle School	High School	Middle School	High School
Community	Community Disorganization	48	49	42	44	47	47
	Transitions and Mobility	47	49	59	61	47	46
	Laws and Norms Favorable to Drug Use	50	30	37	31	42	42
	Perceived Availability of Drugs	39	25	37	27	45	45
	Perceived Availability of Handguns	30	41	24	36	25	42
Family	Poor Family Management	48	42	40	38	44	45
	Family Conflict	34	28	38	33	42	37
School	Poor Academic Performance	48	51	42	44	45	48
	Lack of Commitment to School	60	59	53	54	47	46
Peer and	Favorable Attitudes toward Antisocial Behavior	41	31	39	35	40	46
Individual	Favorable Attitudes toward ATOD Use	40	36	32	36	39	45
	Early Initiation of Drug Use	29	26	23	22	41	46
Average Preva	lence Rate	43	39	39	38	42	45

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values. The symbol "--" indicates that data are not available.

Table 23. Protective factor prevalence rate trends among middle school students for Hardee, 2006 to 2016

Domain	Scale			Hai	dee		
		2006	2008	2010	2012	2014	2016
Community	Community Rewards for Prosocial Involvement	50	52	50	50	43	45
Family	Family Opportunities for Prosocial Involvement	51	53	47	54	57	55
	Family Rewards for Prosocial Involvement	48	42	45	49	45	50
School	School Opportunities for Prosocial Involvement	44	36	38	40	41	35
	School Rewards for Prosocial Involvement	42	41	44	53	43	42
Peer and Individual	Religiosity	55	56	55	56	61	57
Average Prevalence R	ate	48	47	47	50	48	47

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values. The symbol "--" indicates that data are not available.

Table 24. Risk factor prevalence rate trends among middle school students for Hardee, 2006 to 2016

Domain	Scale			Hai	rdee		
		2006	2008	2010	2012	2014	2016
Community	Community Disorganization	48	62	56	53	51	48
	Transitions and Mobility	54	50	52	55	50	47
	Laws and Norms Favorable to Drug Use	46	52	44	40	46	50
	Perceived Availability of Drugs	50	55	46	42	38	39
	Perceived Availability of Handguns	36	38	32	28	31	30
Family	Poor Family Management	53	59	55	49	49	48
	Family Conflict	42	41	42	35	38	34
School	Poor Academic Performance	43	52	45	48	46	48
	Lack of Commitment to School	49	54	53	47	55	60
Peer and	Favorable Attitudes toward Antisocial Behavior	52	53	50	42	45	41
Individual	Favorable Attitudes toward ATOD Use	52	49	47	40	39	40
	Early Initiation of Drug Use	51	47	42	36	29	29
Average Preval	lence Rate	49	50	48	43	43	43

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.

Table 25. Protective factor prevalence rate trends among high school students for Hardee, 2006 to 2016

Domain	Scale	Hardee						
		2006	2008	2010	2012	2014	2016	
Community	Community Rewards for Prosocial Involvement	63	62	63	64	0	66	
Family	Family Opportunities for Prosocial Involvement	49	53	50	58	0	51	
	Family Rewards for Prosocial Involvement	50	51	48	47	0	46	
School	School Opportunities for Prosocial Involvement	58	58	53	39	0	57	
	School Rewards for Prosocial Involvement	44	56	58	61	0	54	
Peer and Individual	Religiosity	76	67	67	66	0	70	
Average Prevalence Rate		57	58	57	56	0	57	

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values. The symbol "--" indicates that data are not available.

Table 26. Risk factor prevalence rate trends among high school students for Hardee, 2006 to 2016

Domain	Scale	Hardee						
		2006	2008	2010	2012	2014	2016	
Community	Community Disorganization	47	52	52	61	0	49	
	Transitions and Mobility	62	56	52	61	0	49	
	Laws and Norms Favorable to Drug Use	41	38	38	39	0	30	
	Perceived Availability of Drugs	43	47	41	29	0	25	
	Perceived Availability of Handguns	62	53	49	40	0	41	
Family	Poor Family Management	54	50	52	33	0	42	
	Family Conflict	33	34	27	23	0	28	
School	Poor Academic Performance	45	47	47	55	0	51	
	Lack of Commitment to School	42	41	48	49	0	59	
Peer and	Favorable Attitudes toward Antisocial Behavior	46	47	45	36	0	31	
Individual	Favorable Attitudes toward ATOD Use	43	42	43	34	0	36	
	Early Initiation of Drug Use	41	41	43	45	0	26	
Average Prevalence Rate		44	43	47	41	0	39	

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.