2016 Florida Youth Substance Abuse Survey

Table 1. Major demographic characteristics of surveyed Okaloosa and Florida Statewide, 2016

	Oka	loosa	Florida S	tatewide
	N	%	N	%
Sex				
Female	1,135	49.0	31,515	47.9
Male	1,126	48.6	32,905	50.0
Race/Ethnic group				
African American	229	9.9	14,666	22.3
American Indian	67	2.9	568	0.9
Asian	43	1.9	743	1.1
Hispanic/Latino	149	6.4	13,174	20.0
Native Hawaiian/Pacific Islander	15	0.6	144	0.2
Other/Multiple	484	20.9	7,523	11.4
White, non-Hispanic	1,298	56.0	28,309	43.0
Age				
10	4	0.2	65	0.1
11	170	7.3	3,856	5.9
12	374	16.1	8,338	12.7
13	370	16.0	9,230	14.0
14	366	15.8	9,454	14.4
15	364	15.7	10,070	15.3
16	270	11.7	9,684	14.7
17	238	10.3	9,348	14.2
18	137	5.9	4,799	7.3
19 or older	10	0.4	618	0.9
Grade				
6th	412	17.8	9,301	14.1
7th	371	16.0	9,215	14.0
8th	361	15.6	9,326	14.2
9th	378	16.3	10,140	15.4
10th	322	13.9	9,834	15.0
11th	230	9.9	9,254	14.1
12th	242	10.4	8,705	13.2
Overall Middle School	1,144	49.4	27,678	42.1
Overall High School	1,172	50.6	37,765	57.4
Total	2,316	100.0	65,776	100.0

Note: Some categories do not sum to 100% of the total due to missing values (e.g., not all survey questions were answered). In addition, rounding can produce totals that do not equal 100%. "N" represents the number of valid cases. In this table, county data are unweighted while statewide data are weighted.

Table 2. Percentages of Okaloosa and Florida Statewide youth who reported having used various drugs in their lifetimes, 2016

			(Okaloosa	ì					Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Alcohol	22.2	52.6	41.4	38.4	24.0	51.9	39.5	21.8	51.4	41.3	37.1	23.6	51.5	39.1
Blacking Out		18.7	18.8	18.8		19.8	18.7		15.9	16.5	15.4		16.1	15.9
Cigarettes	10.2	24.9	18.2	19.0	9.8	24.3	18.5	8.0	18.5	14.0	14.1	8.2	18.5	14.1
Vaporizer/E-Cigarette	13.2	37.9	27.8	27.0	13.9	37.9	27.1	14.6	33.9	24.4	27.1	15.7	34.5	25.8
Marijuana or Hashish	6.9	33.5	22.0	22.1	8.0	32.1	21.8	7.0	31.7	21.4	21.3	8.1	31.8	21.3
Synthetic Marijuana		6.2	4.9	7.7		6.1	6.2		4.9	4.8	5.0		4.8	4.9
Inhalants	8.3	4.8	6.5	6.3	8.2	4.4	6.3	6.4	4.6	6.1	4.7	6.4	4.7	5.4
Flakka		0.6	0.1	1.0		0.6	0.6		1.0	0.9	1.1		1.0	1.0
Club Drugs	1.1	4.3	2.9	2.9	0.9	4.7	2.9	0.7	3.0	1.9	2.2	0.8	2.9	2.1
LSD, PCP or Mushrooms	1.3	6.6	3.9	4.6	1.2	6.5	4.3	1.0	4.8	2.7	3.7	1.1	4.9	3.2
Methamphetamine	0.5	0.6	0.3	0.9	0.4	0.9	0.6	0.5	0.8	0.6	0.8	0.5	0.8	0.7
Cocaine or Crack Cocaine	1.0	3.6	2.3	2.7	0.8	4.0	2.5	0.8	2.5	1.6	1.9	0.8	2.3	1.8
Heroin	0.4	0.2	0.2	0.4	0.4	0.2	0.3	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Depressants	3.3	13.4	9.9	8.2	3.5	14.7	9.0	2.2	6.5	5.3	4.1	2.4	6.6	4.7
Prescription Pain Relievers	3.4	8.5	6.6	6.2	3.6	8.5	6.3	3.3	5.8	5.3	4.2	3.5	5.7	4.8
Prescription Amphetamines	1.6	7.7	5.1	5.2	1.7	7.9	5.0	1.3	4.7	3.3	3.2	1.4	4.6	3.2
Steroids (without a doctor's order)	0.7	0.6	0.4	0.9	0.7	0.3	0.7	0.6	0.5	0.4	0.6	0.5	0.5	0.5
Over-the-Counter Drugs	2.9	6.4	5.0	4.4	3.2	6.3	4.8	3.2	5.3	4.7	4.2	3.4	5.4	4.4
Needle to Inject Illegal Drugs		0.6	0.3	0.9		0.7	0.6		0.8	0.6	0.8		0.8	0.8
Any illicit drug	16.4	38.8	30.0	28.4	16.9	38.1	29.0	15.7	36.4	29.0	26.4	16.6	36.6	27.7
Any illicit drug other than marijuana	13.3	24.1	19.7	19.0	13.1	24.2	19.4	12.0	17.7	16.4	14.2	12.3	17.7	15.3
Alcohol only	12.2	18.7	16.6	15.5	12.9	19.5	15.9	12.5	21.0	18.1	17.0	13.4	21.0	17.5
Alcohol or any illicit drug	28.2	57.6	46.5	43.8	29.4	57.7	44.8	27.8	57.2	46.8	43.1	29.6	57.4	44.8
Any illicit drug, but no alcohol	6.5	5.1	5.5	5.8	6.0	5.8	5.7	6.5	6.1	6.0	6.5	6.5	6.1	6.3

Note: The first 19 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. Respondents were asked on how many occasions in their lifetime they woke up after a night of drinking and did not remember the things they did or the places they went. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "--" indicates that data are not available.

 $Table \ 3. \ Percentages \ of \ Okaloosa \ and \ Florida \ Statewide \ youth \ who \ reported \ having \ used \ various \ drugs \ in \ the \ past \ 30 \ days, \\ 2016$

			(Okaloosa	ì					Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Alcohol	9.1	25.9	20.5	17.2	9.8	25.1	18.6	8.3	25.5	19.9	17.0	9.1	25.2	18.3
Binge Drinking	2.3	11.4	6.3	8.8	2.7	11.3	7.5	3.2	10.9	7.9	7.7	3.4	10.8	7.7
Cigarettes	1.8	6.3	3.9	4.8	1.6	6.1	4.3	1.4	4.8	3.3	3.5	1.5	4.6	3.4
Vaporizer/E-Cigarette	4.4	13.6	8.7	10.6	5.4	11.8	9.6	5.1	12.8	8.4	10.6	5.5	12.9	9.6
Marijuana or Hashish	3.7	15.5	9.9	11.0	3.8	15.7	10.4	3.2	17.0	10.9	11.5	3.9	16.9	11.2
Synthetic Marijuana		0.6	0.7	0.6		0.9	0.6		1.0	0.9	1.2		1.1	1.0
Inhalants	2.9	0.4	1.6	1.5	2.4	0.7	1.5	2.2	1.2	1.9	1.4	2.1	1.2	1.6
Flakka		0.3	0.2	0.4		0.3	0.3		0.5	0.4	0.7		0.5	0.5
Club Drugs	0.4	0.4	0.2	0.6	0.3	0.7	0.4	0.3	0.8	0.5	0.6	0.3	0.7	0.6
LSD, PCP or Mushrooms	0.4	1.0	0.7	0.8	0.2	1.3	0.7	0.3	1.4	0.7	1.1	0.3	1.4	0.9
Methamphetamine	0.2	0.2	0.1	0.4	0.1	0.4	0.2	0.3	0.4	0.3	0.4	0.3	0.4	0.4
Cocaine or Crack Cocaine	0.4	1.2	1.0	0.8	0.3	1.7	0.9	0.3	0.8	0.6	0.6	0.3	0.8	0.6
Heroin	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.2	0.1	0.2	0.1	0.2	0.2
Depressants	1.3	4.5	3.1	3.0	1.4	5.0	3.1	0.8	2.4	2.0	1.5	0.9	2.5	1.8
Prescription Pain Relievers	1.6	2.2	2.3	1.7	1.7	2.1	1.9	1.6	2.0	2.2	1.5	1.6	2.1	1.8
Prescription Amphetamines	0.3	2.3	1.1	1.8	0.4	2.4	1.4	0.5	1.6	1.1	1.2	0.6	1.7	1.2
Steroids (without a doctor's order)	0.0	0.2	0.0	0.3	0.0	0.1	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Over-the-Counter Drugs	1.0	2.1	1.8	1.3	1.2	1.8	1.6	1.8	2.1	2.0	1.9	1.8	2.1	2.0
Any illicit drug	7.3	19.4	13.8	14.5	7.1	19.6	14.1	7.5	20.0	15.1	14.2	8.1	20.0	14.7
Any illicit drug other than marijuana	5.0	9.0	6.9	7.6	4.8	9.6	7.3	5.6	7.7	7.3	6.2	5.7	7.7	6.8
Alcohol only	5.6	13.3	11.6	8.6	6.1	12.7	10.0	5.2	13.4	11.1	9.1	5.6	13.3	10.0
Alcohol or any illicit drug	12.6	32.5	25.4	22.7	12.9	32.0	23.8	12.5	32.9	25.8	22.9	13.5	32.9	24.3
Any illicit drug, but no alcohol	3.8	6.7	5.1	5.8	3.3	6.9	5.5	4.5	7.8	6.4	6.3	4.7	8.0	6.4

Note: The first 18 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. Binge drinking is defined as having had five or more alcoholic drinks in a row in the past two weeks. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "--" indicates that data are not available.

Table 4. Lifetime trend in alcohol, tobacco and other drug use for Okaloosa, 2006 to 2016

		2006			2008			2010			2012			2014			2016	
	Middle School	0	Total	Middle School		Total	Middle School		Total	Middle School	High School	Total	Middle School		Total	Middle School	U	Total
Alcohol				37.2	71.1	57.0	33.4	61.5	49.7	26.7	58.6	45.1	22.5	56.1	41.9	22.2	52.6	39.5
Blacking Out														19.5	19.5		18.7	18.7
Cigarettes				18.5	43.5	33.1	18.9	36.8	29.3	14.3	32.8	25.0	6.9	25.5	17.6	10.2	24.9	18.5
Vaporizer/E-Cigarette																13.2	37.9	27.1
Marijuana or Hashish				8.7	35.8	24.6	9.7	31.3	22.3	6.3	29.9	20.0	5.2	29.0	19.0	6.9	33.5	21.8
Synthetic Marijuana											14.4	14.4		7.7	7.7		6.2	6.2
Inhalants				15.2	11.7	13.2	14.1	8.5	10.8	12.0	6.7	8.9	9.5	6.7	7.9	8.3	4.8	6.3
Flakka																	0.6	0.6
Club Drugs				1.7		1.7	2.5	4.9	3.9	1.1	5.0	3.4	0.6	6.4	3.9	1.1	4.3	2.9
LSD, PCP or Mushrooms				2.8		2.8	3.3	6.0	4.9	0.9	5.2	3.4	0.5	6.2	3.8	1.3	6.6	4.3
Methamphetamine				1.7	2.0	1.9	1.8	0.8	1.2	1.0	1.2	1.1	0.0	1.0	0.6	0.5	0.6	0.6
Cocaine or Crack Cocaine				2.4		2.4	2.5	3.9	3.3	0.6	4.5	2.8	0.6	3.1	2.1	1.0	3.6	2.5
Heroin				1.3	1.2	1.2	1.3	1.1	1.2	0.3	0.4	0.4	0.0	1.7	1.0	0.4	0.2	0.3
Depressants				3.1	13.4	9.1	4.3	10.0	7.6	3.0	7.9	5.8	2.0	7.7	5.3	3.3	13.4	9.0
Prescription Pain Relievers				6.3	16.1	12.0	6.7	12.1	9.8	4.0	11.3	8.2	3.0	9.1	6.5	3.4	8.5	6.3
Prescription Amphetamines				2.8	11.3	7.7	2.1	9.4	6.4	1.7	9.2	6.0	1.0	8.7	5.5	1.6	7.7	5.0
Steroids (without a doctor's order)				2.2	1.4	1.8	0.6	1.6	1.2	0.5	1.4	1.0	0.2	1.2	0.8	0.7	0.6	0.7
Over-the-Counter Drugs				5.3		5.3	6.3	9.7	8.3	3.4	7.9	5.9	3.1	6.1	4.9	2.9	6.4	4.8
Needle to Inject Drugs																	0.6	0.6
Any illicit drug				23.3	44.2	35.5	24.5	37.5	32.0	19.1	38.4	30.2	17.4	37.7	29.1	16.4	38.8	29.0
Any illicit drug other than marijuana				20.0	30.9	26.4	21.1	24.4	23.0	16.8	24.2	21.0	14.6	22.3	19.1	13.3	24.1	19.4
Alcohol only				20.4	29.4	25.7	17.6	27.5	23.4	14.9	24.9	20.7	14.8	21.6	18.7	12.2	18.7	15.9
Alcohol or any illicit drug				43.7	73.5	61.1	42.0	64.7	55.2	33.9	63.0	50.7	32.1	59.3	47.8	28.2	57.6	44.8
Any illicit drug, but no alcohol				6.8	2.8	4.5	8.7	3.3	5.6	7.3	4.3	5.5	9.7	3.4	6.0	6.5	5.1	5.7

Note: The first 19 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. Respondents were asked on how many occasions in their lifetime they woke up after a night of drinking and did not remember the things they did or the places they went. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "--" indicates that data are not available.

2016 Florida Youth Substance Abuse Survey - Okaloosa Report

Table 5. Past-30-day trend in alcohol, tobacco and other drug use for Okaloosa, 2006 to 2016

		2006			2008			2010			2012			2014			2016	
	Middle School	U	Total	Middle School	0	Total	Middle School	High School	Total	Middle School		Total	Middle School		Total	Middle School	High School	Total
Alcohol				17.4	40.2	30.7	15.4	34.8	26.7	9.8	30.1	21.4	9.3	27.9	20.1	9.1	25.9	18.6
Binge Drinking				5.7	21.7	15.0	6.7	19.6	14.2	3.6	15.0	10.2	2.6	12.6	8.3	2.3	11.4	7.5
Cigarettes				4.8	16.9	11.9	5.6	15.8	11.5	3.4	11.8	8.3	1.1	4.0	2.8	1.8	6.3	4.3
Vaporizer/E-Cigarette																4.4	13.6	9.6
Marijuana or Hashish				4.5	18.0	12.4	5.3	16.5	11.8	2.0	14.7	9.4	2.9	13.9	9.2	3.7	15.5	10.4
Synthetic Marijuana											4.3	4.3		1.6	1.6		0.6	0.6
Inhalants				6.6	2.5	4.2	6.2	1.7	3.6	3.4	1.4	2.3	2.3	1.6	1.9	2.9	0.4	1.5
Flakka																	0.3	0.3
Club Drugs				0.7		0.7	0.5	1.1	0.8	0.3	1.3	0.9	0.3	1.9	1.3	0.4	0.4	0.4
LSD, PCP or Mushrooms				1.0		1.0	1.3	1.6	1.5	0.0	1.0	0.6	0.3	1.4	0.9	0.4	1.0	0.7
Methamphetamine				0.8	0.1	0.4	0.6	0.4	0.5	0.4	0.3	0.3	0.0	0.6	0.3	0.2	0.2	0.2
Cocaine or Crack Cocaine				1.3		1.3	1.1	1.0	1.1	0.3	1.1	0.8	0.5	1.8	1.3	0.4	1.2	0.9
Heroin				0.7	0.2	0.4	0.8	0.5	0.6	0.0	0.3	0.2	0.0	0.4	0.2	0.0	0.0	0.0
Depressants				1.3	4.8	3.4	1.6	2.9	2.4	0.9	2.8	2.0	0.5	1.6	1.2	1.3	4.5	3.1
Prescription Pain Relievers				3.0	5.5	4.5	3.1	4.6	4.0	1.9	3.5	2.8	2.1	2.2	2.2	1.6	2.2	1.9
Prescription Amphetamines				1.6	3.4	2.7	1.4	3.1	2.4	1.0	3.4	2.4	1.0	1.8	1.5	0.3	2.3	1.4
Steroids (without a doctor's order)				1.4	0.7	1.0	0.1	1.0	0.6	0.3	1.0	0.7	0.0	0.0	0.0	0.0	0.2	0.1
Over-the-Counter Drugs				2.0		2.0	3.7	3.1	3.4	1.2	3.0	2.2	1.5	3.7	2.7	1.0	2.1	1.6
Any illicit drug				12.3	23.0	18.6	13.3	20.0	17.2	6.7	20.4	14.6	6.2	19.5	13.9	7.3	19.4	14.1
Any illicit drug other than marijuana				10.4	11.4	11.0	11.1	9.5	10.1	6.0	11.1	8.9	4.3	8.4	6.7	5.0	9.0	7.3
Alcohol only				10.3	22.4	17.4	8.7	19.9	15.2	6.6	16.2	12.1	6.3	15.2	11.4	5.6	13.3	10.0
Alcohol or any illicit drug				22.4	44.9	35.5	21.9	39.5	32.1	13.0	36.2	26.4	12.3	34.9	25.3	12.6	32.5	23.8
Any illicit drug, but no alcohol				5.3	5.1	5.2	6.6	4.8	5.5	3.4	5.7	4.8	3.1	6.9	5.3	3.8	6.7	5.5

Note: The first 18 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. Binge drinking is defined as having had five or more alcoholic drinks in a row in the past two weeks. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "--" indicates that data are not available.

Table 6. Percentages of Okaloosa and Florida Statewide youth who reported early ATOD use, perceived risk of harm, personal disapproval, parental disapproval and peer disapproval, 2016

			(Okaloosa	1					Flori	da State	ewide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Early ATOD use														
More than a sip of alcohol		18.6	18.1	19.2		18.0	18.6		19.4	18.3	20.2		19.3	19.4
Drinking at least once a month		2.6	2.4	2.6		2.5	2.6		3.5	3.7	3.4		3.5	3.5
Smoke cigarettes		12.1	11.2	12.8		11.4	12.1		9.3	8.6	9.8		9.4	9.3
Smoke marijuana		9.5	8.8	10.4		10.0	9.5		10.6	9.0	12.0		10.8	10.6
Great risk of harm														
1 or 2 drinks nearly every day	44.0	35.8	42.3	36.5	43.6	36.1	39.4	45.7	40.7	46.4	39.3	45.9	40.0	42.8
5+ drinks once or twice a week	59.6	49.6	57.5	50.0	58.5	50.4	53.9	58.1	52.3	58.4	51.2	58.4	51.8	54.7
1+ packs of cigarettes per day	72.6	64.6	67.2	68.8	72.7	64.9	68.1	67.9	68.8	68.4	68.5	68.7	68.2	68.4
Marijuana once or twice a week	58.5	26.1	39.9	39.7	56.0	26.0	40.2	51.7	25.2	37.5	35.2	50.1	24.4	36.3
Try marijuana once or twice	38.0	15.0	24.4	25.4	36.2	14.9	25.1	36.1	16.8	24.8	25.0	34.7	16.4	24.9
Use prescription drugs	71.9	65.8	69.0	68.2	71.1	65.5	68.5	69.1	68.2	70.0	67.3	69.4	67.9	68.5
Think it would be wrong to														
Smoke cigarettes	95.0	81.7	87.8	86.7	94.5	83.3	87.5	95.6	87.7	91.1	91.0	95.3	88.9	91.0
Drink alcohol regularly	89.6	64.0	75.0	74.9	88.1	63.4	75.1	88.8	64.5	74.1	75.2	87.6	64.8	74.7
Smoke marijuana	90.5	63.8	76.3	73.9	89.3	64.4	75.4	89.7	62.3	74.4	73.3	88.1	62.3	73.8
Smoke synthetic marijuana	93.7	86.8	90.0	89.4	93.2	86.2	89.8	93.1	88.6	90.1	90.9	92.7	88.6	90.5
Use other illicit drugs	98.4	93.1	96.7	94.0	98.4	93.0	95.4	97.5	93.6	95.8	94.7	97.3	93.6	95.2
Parents think it would be wrong to														
1 or 2 drinks nearly every day	98.1	92.4	94.5	95.3	98.0	93.0	94.9	97.3	93.8	95.4	95.1	97.2	94.1	95.2
Smoke cigarettes	98.8	95.2	97.0	96.5	98.8	96.4	96.8	98.6	96.8	97.6	97.5	98.6	97.3	97.6
Smoke marijuana	97.1	88.8	93.5	91.4	97.1	89.4	92.4	97.0	88.6	92.3	92.0	96.6	88.9	92.1
Use prescription drugs	99.0	96.4	97.4	97.8	98.5	96.9	97.6	98.2	97.3	97.4	97.9	98.1	97.4	97.7
Friends think it would be wrong to														
1 or 2 drinks nearly every day	91.6	75.4	84.7	80.2	91.3	75.3	82.5	91.3	79.0	85.6	82.9	90.7	79.1	84.2
Smoke tobacco	92.7	83.1	89.6	84.9	92.8	84.8	87.3	94.8	87.0	91.1	89.5	94.4	87.8	90.3
Smoke marijuana	88.6	61.5	74.2	71.7	88.0	60.8	73.3	88.6	59.9	72.6	71.3	86.6	59.8	72.0
Use prescription drugs	96.0	87.4	92.2	89.9	95.4	87.2	91.2	95.7	90.6	93.4	92.2	95.5	90.3	92.7

Note: Early ATOD use is defined as the percentage of high school students who started using at age 13 or younger. The symbol "--" indicates that data are not available.

2016 Florida Youth Substance Abuse Survey - Okaloosa Report

Table 7. Trends in early ATOD use and attitudes toward substance use for Okaloosa, 2006 to 2016

		2006			2008			2010			2012			2014			2016	
	Middle School	High School	Total	Middle School	High School	Total	Middle School	High School	Total	Middle School		Total	Middle School	High School	Total	Middle School	High School	Total
Early ATOD use																		
Try alcohol					26.9	26.9		24.2	24.2		20.1	20.1		18.0	18.0		18.6	18.6
Drink monthly					4.7	4.7		3.7	3.7		3.4	3.4		1.6	1.6		2.6	2.6
Smoke cigarettes					23.3	23.3		17.8	17.8		16.0	16.0		11.0	11.0		12.1	12.1
Smoke marijuana					11.3	11.3		9.1	9.1		9.0	9.0		6.3	6.3		9.5	9.5
Great risk of harm																		
Daily alcohol use				44.5	41.7	42.9	44.0	43.9	44.0	43.2	39.3	41.0	43.1	37.3	39.8	44.0	35.8	39.4
5+ drinks weekly													58.6	59.7	59.2	59.6	49.6	53.9
Daily cigarette use				74.9	69.2	71.6	75.9	68.1	71.4	71.4	71.8	71.7	73.5	75.6	74.7	72.6	64.6	68.1
Marijuana weekly*													62.1	38.9	48.8	58.5	26.1	40.2
Try marijuana				44.0	23.3	31.8	42.2	23.3	31.2	43.9	20.1	30.1	41.6	18.0	28.0	38.0	15.0	25.1
Prescription drugs											67.3	67.3	79.9	75.2	77.2	71.9	65.8	68.5
Think it wrong																		
Smoke cigarettes				89.6	65.6	75.5	88.9	69.8	77.8	93.6	78.8	85.0	96.0	87.2	90.9	95.0	81.7	87.5
Drink regularly				81.4	50.4	63.2	81.4	58.9	68.4	87.3	60.7	71.9	91.9	66.7	77.4	89.6	64.0	75.1
Smoke marijuana				91.2	70.2	78.9	89.2	71.5	78.9	92.4	71.8	80.4	94.2	74.0	82.6	90.5	63.8	75.4
Smoke synthetic marijuana																93.7	86.8	89.8
Use other illicit drugs				97.5	91.2	93.8	95.9	94.3	95.0	97.3	94.4	95.7	99.0	96.0	97.3	98.4	93.1	95.4
Parents think it wrong																		
Daily alcohol use*													99.3	96.4	97.6	98.1	92.4	94.9
Smoke cigarettes				95.1	89.6	91.9	96.0	90.6	92.9	98.1	92.2	94.7	99.7	96.5	97.8	98.8	95.2	96.8
Smoke marijuana				96.8	93.2	94.7	95.9	93.9	94.8	97.9	92.2	94.6	98.9	95.0	96.6	97.1	88.8	92.4
Prescription drugs													99.4	96.7	97.8	99.0	96.4	97.6
Friends think it wrong																		
Daily alcohol use													92.3	78.8	84.5	91.6	75.4	82.5
Smoke tobacco													94.6	83.5	88.3	92.7	83.1	87.3
Smoke marijuana													91.7	65.4	76.5	88.6	61.5	73.3
Prescription drugs													97.5	89.9	93.1	96.0	87.4	91.2

Note: Early ATOD use is defined as the percentage of high school students who started using at age 13 or younger. The symbol "--" indicates that data are not available.

^{*} In 2014, the description of marijuana use in the risk of harm question was changed from "regularly" to "once or twice a week." Also, in 2014, the description of alcohol use in the parent disapproval question was changed from "regularly" to "nearly every day." For these items, care should be exercised when comparing 2014 data to previous years.

Table 8. Among $\underline{\text{high school}}$ drinkers, usual source of alcohol within the past 30 days, Okaloosa and Florida Statewide youth, 2016

			(Okaloosa	ı					Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Bought in a store		6.9	0.6	14.4		6.1	6.9		8.3	5.0	12.4		7.3	8.3
Bought in a restaurant, bar, or club		0.6	0.0	1.2		0.0	0.6		1.6	1.2	2.1		1.5	1.6
Bought at a public event		0.5	0.6	0.4		0.3	0.5		0.7	0.7	0.8		0.6	0.7
Someone bought it for me		19.2	12.9	26.9		20.2	19.2		14.7	13.3	16.3		14.5	14.7
Someone gave it to me		40.1	51.6	26.2		41.7	40.1		44.8	50.6	37.8		47.0	44.8
Took it from a store		1.8	2.3	1.2		1.9	1.8		0.3	0.2	0.4		0.4	0.3
Took it from a family member		12.6	14.1	11.1		12.4	12.6		11.7	14.2	8.7		12.4	11.7
Some other way		18.3	17.9	18.6		17.4	18.3		17.8	14.7	21.5		16.5	17.8

Table 9. Among $\underline{\text{high school}}$ drinkers, usual drinking location within the past 30 days, Okaloosa and Florida Statewide youth, 2016

			(Okaloosa	a					Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
My home		32.6	32.4	33.6		29.3	32.6		37.7	39.2	35.9		37.1	37.7
Another person's home		50.0	54.0	44.7		49.5	50.0		40.0	41.0	38.9		40.8	40.0
Car or other vehicle		1.7	1.5	1.9		2.3	1.7		1.7	1.6	1.9		1.7	1.7
Restaurant, bar, or club		2.5	3.2	1.6		2.7	2.5		2.7	2.9	2.5		2.4	2.7
Public place		2.4	2.9	1.8		3.3	2.4		3.9	3.9	3.7		4.3	3.9
Public event		0.2	0.0	0.5		0.3	0.2		1.6	1.8	1.5		1.7	1.6
School property		2.0	1.1	3.1		2.7	2.0		1.3	0.7	2.0		1.4	1.3
Some other place		8.6	5.0	12.8		9.8	8.6		11.1	9.0	13.7		10.6	11.1

Table 10. Among <u>high school</u> drinkers, number of drinks usually consumed per day, on the days students drank in the past 30 days, Okaloosa and Florida Statewide youth, 2016

			(Okaloosa	ì					Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
1		26.0	30.7	20.6		26.7	26.0		29.6	31.6	27.4		30.5	29.6
2		20.4	23.8	15.8		18.4	20.4		22.5	24.8	19.8		22.4	22.5
3		19.2	19.3	19.5		19.3	19.2		17.3	17.9	16.3		16.9	17.3
4		9.1	7.5	11.2		9.9	9.1		8.6	9.0	8.3		8.5	8.6
5 or more		25.3	18.8	32.8		25.7	25.3		22.0	16.7	28.2		21.8	22.0

Table 11. Percentages of Okaloosa and Florida Statewide youth who reported drinking alcohol, smoking marijuana, or using another drug to get high $\underline{\text{before or during school}}$ in the past 12 months, 2016

			(Okaloosa	1					Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Drinking alcohol	2.0	5.9	4.2	4.0	2.0	6.2	4.2	3.3	6.5	5.5	4.7	3.3	6.8	5.1
Smoking marijuana	2.8	10.6	6.8	7.8	2.8	11.3	7.2	3.2	12.2	8.1	8.7	3.7	12.3	8.5
Used another drug to get high	0.8	5.2	2.8	3.6	1.1	5.5	3.3	1.4	3.9	2.9	2.9	1.5	4.1	2.9

Table 12. Percentages of Okaloosa and Florida Statewide youth who reported engaging in delinquent behavior within the past 12 months, 2016

			(Okaloosa	1					Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Carrying a handgun	8.1	5.9	3.2	10.7	7.3	6.5	6.9	5.0	5.8	2.6	8.3	5.0	5.9	5.5
Selling drugs	1.4	7.0	3.5	5.8	1.6	7.4	4.5	1.6	6.0	3.0	5.4	1.8	6.1	4.2
Attempting to steal a vehicle	0.4	1.0	1.0	0.5	0.6	0.9	0.7	1.0	1.5	0.8	1.7	1.0	1.5	1.3
Being arrested	1.9	2.4	1.7	2.7	1.5	2.4	2.1	1.9	2.8	1.8	3.0	1.8	3.0	2.4
Taking a handgun to school	0.1	0.3	0.1	0.3	0.1	0.4	0.2	0.4	0.7	0.3	0.9	0.4	0.7	0.6
Getting suspended	5.5	6.1	3.3	8.2	5.0	6.5	5.8	10.6	9.1	7.0	12.5	10.1	9.8	9.8
Attacking someone with intent to harm	5.3	5.3	4.4	6.2	5.1	5.3	5.3	6.2	6.1	5.3	6.9	6.0	6.6	6.1

Table 13. Trends in delinquent behaviors for Okaloosa, 2006 to 2016

		2006			2008			2010			2012			2014			2016	
	Middle School	High School	Total															
Carrying a handgun				2.7	5.2	4.2	5.1	3.2	4.0	5.5	4.1	4.7	3.6	3.8	3.7	8.1	5.9	6.9
Selling drugs				1.5	9.0	5.9	3.7	8.3	6.4	1.7	7.1	4.8	0.7	3.4	2.3	1.4	7.0	4.5
Attempting to steal a vehicle				1.3	3.2	2.4	2.5	1.7	2.1	0.8	2.2	1.6	0.5	0.3	0.4	0.4	1.0	0.7
Being arrested				3.0	7.2	5.5	3.0	4.3	3.8	2.4	3.2	2.9	0.7	1.8	1.3	1.9	2.4	2.1
Taking a handgun to school				0.3	0.8	0.6	1.2	0.6	0.9	0.3	0.4	0.4	0.0	0.2	0.1	0.1	0.3	0.2
Getting suspended				11.1	16.1	14.0	11.9	10.8	11.2	11.8	10.1	10.8	4.1	5.5	4.9	5.5	6.1	5.8
Attacking someone with intent to harm				10.3	13.5	12.2	11.3	8.5	9.7	5.4	4.9	5.1	2.9	7.0	5.3	5.3	5.3	5.3

Table 14. Percentages of Okaloosa and Florida Statewide youth with family members who have been or are now in jail or friends who have been in trouble because of alcohol or drugs, 2016

			(Okaloosa	l					Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Family members who have been or are now in jail or prison	42.2	41.9	44.5	39.5	41.3	41.7	42.0	44.4	44.6	47.8	41.2	44.2	45.4	44.5
Friends who have have been or are in trouble because of alcohol or drugs	26.1	54.5	44.9	39.0	29.3	52.7	42.0	23.9	46.5	38.4	35.8	26.0	47.1	37.1

Table 15. Percentages of Okaloosa and Florida Statewide youth who reported involvement in bullying behavior, 2016

			(Okaloosa	ì					Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Skipped school because of bullying	8.1	9.1	13.1	4.5	8.6	8.7	8.6	7.6	8.2	11.5	4.4	7.6	8.3	7.9
Was kicked or shoved in past 30 days	18.0	8.8	11.9	13.4	17.7	8.8	12.8	19.6	8.1	12.0	13.8	18.8	8.1	13.0
Was taunted or teased in past 30 days	36.6	24.2	32.6	26.3	35.0	24.9	29.6	35.3	22.5	30.1	25.6	34.4	22.8	27.9
Was victim of cyber bullying in past 30 days	7.4	9.8	12.3	5.3	7.8	10.1	8.8	7.8	7.6	10.2	5.2	7.8	7.9	7.7
Physically bullied others in past 30 days	5.9	3.2	3.8	4.7	5.6	3.1	4.4	7.6	4.9	5.0	7.0	7.2	5.2	6.1
Verbally bullied others in past 30 days	12.0	9.2	10.0	10.9	11.4	9.5	10.4	14.3	10.4	11.0	13.0	14.0	10.7	12.0
Cyber bullied others in past 30 days	2.2	3.6	3.9	2.0	2.5	3.5	3.0	3.6	4.1	4.0	3.7	3.7	4.1	3.9

Table 16. Percentages of Okaloosa and Florida Statewide youth who reported gang membership, 2016

			. (Okaloosa	ı					Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Have you ever belonged to a gang?	3.5	2.8	2.1	4.0	2.8	3.6	3.1	3.3	3.5	2.3	4.5	3.1	3.6	3.4
Did that gang have a name?	32.9	14.2	13.3	25.0	23.7	19.9	19.9	23.4	14.1	13.1	19.7	21.9	14.4	16.9
Are you a gang member now?		1.3	0.8	1.8		1.3	1.3		2.0	1.1	2.8		2.0	2.0

Note: The prevalence rates for "Did that gang have a name?" exclude students who reported that they have never belonged to a gang. The symbol "--" indicates that data are not available.

Table 17. Reasons for joining a gang, reported by <u>high school students who have belonged to a gang</u>, among Okaloosa and Florida Statewide youth, 2016

			(Okaloosa	ì					Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Fun and excitement		3.8	2.7	4.6		3.8	3.8		5.0	4.3	5.5		5.0	5.0
Protection		4.5	3.0	5.6		5.5	4.5		4.2	2.6	5.4		4.5	4.2
Friend or relative in a gang		3.8	1.4	5.9		3.6	3.8		4.1	2.9	4.9		4.1	4.1
Forced to join		1.2	2.3	0.3		1.4	1.2		1.1	1.1	1.2		1.2	1.1
To get respect		2.6	0.8	3.7		2.8	2.6		3.5	2.0	4.7		3.5	3.5
Money		2.9	2.3	3.5		1.9	2.9		3.8	2.5	4.8		4.1	3.8
To fit in better		1.8	0.8	2.7		2.2	1.8		1.2	1.1	1.4		1.2	1.2
Other reasons		4.9	1.2	7.8		5.0	4.9		5.9	4.5	7.0		5.6	5.9

Table 18. Percentages of Okaloosa and Florida Statewide youth who reported participation in extracurricular activities, 2016

			(Okaloosa	a					Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
School Sports	48.1	38.1	38.4	45.8	46.9	39.6	42.5	36.0	40.4	34.5	42.5	36.4	41.4	38.5
Organized Sports Outside of School	42.4	20.1	28.8	30.5	40.2	20.3	29.8	41.6	22.4	28.2	32.9	40.1	22.8	30.5
School Band	22.2	16.0	20.3	16.6	23.2	15.3	18.7	16.7	8.1	12.3	11.1	16.2	8.1	11.7
School Club(s)	28.2	34.7	40.6	23.4	29.7	33.8	31.8	22.1	31.8	35.4	20.4	22.8	31.6	27.6
Community Club(s)	8.5	12.9	13.3	8.3	9.3	12.8	11.0	9.3	12.8	14.6	8.2	9.4	12.6	11.3

Table 19. Percentages of <u>high school</u> students who reported *riding in* a vehicle within the past 30 days driven by someone who had been drinking alcohol or using marijuana, or *driving* a vehicle within the past 30 days after drinking alcohol or using marijuana, among Okaloosa and Florida Statewide youth, 2016

				Okaloosa	ı					Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Riding in a vehicle														
Alcohol		11.7	12.8	10.8		10.9	11.7		16.4	17.5	15.3		16.3	16.4
Marijuana		19.9	20.2	19.8		19.0	19.9		22.7	23.7	21.7		22.9	22.7
Driving a vehicle														
Alcohol		5.8	6.3	5.0		4.7	5.8		5.4	5.0	5.8		5.2	5.4
Marijuana		8.7	6.9	10.6		8.4	8.7		10.3	9.1	11.4		9.9	10.3

Table 20. Trends for <u>high school</u> students *riding in* a vehicle within the past 30 days driven by someone who had been drinking alcohol or using marijuana, or *driving* a vehicle within the past 30 days after drinking alcohol or using marijuana, for

Okaloosa, 2012 and 2016

		2006		2008		2010			2012			2014			2016	
	Middle School	High School	Middle School	High School	Middle School	O	Total	Middle School	O	Total	Middle School	O	Total	Middle School	High School	Total
Riding in a vehicle																
Alcohol			 		 				18.8	18.8		13.8	13.8		11.7	11.7
Marijuana			 		 				18.1	18.1		15.3	15.3		19.9	19.9
Driving a vehicle																
Alcohol			 		 				7.1	7.1		6.0	6.0		5.8	5.8
Marijuana			 		 				10.2	10.2		8.9	8.9		8.7	8.7

Table 21. Protective factor prevalence rates for Okaloosa, Florida Statewide youth and the national normative database, 2016

Domain	Scale	Oka	loosa	Florida S	Statewide	Nationa	l Norms
		Middle School	High School	Middle School	High School	Middle School	High School
Community	Community Rewards for Prosocial Involvement	52	75	46	69	56	63
Family	Family Opportunities for Prosocial Involvement	63	61	60	59	59	54
	Family Rewards for Prosocial Involvement	58	57	56	56	54	55
School	School Opportunities for Prosocial Involvement	62	70	53	63	57	60
	School Rewards for Prosocial Involvement	52	60	49	59	53	58
Peer and Individual	Religiosity	56	59	49	57	56	62
Average Prevalence R	ate	57	64	52	61	56	59

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.

Table 22. Risk factor prevalence rates for Okaloosa, Florida Statewide youth and the national normative database, 2016

Domain	Scale	Oka	loosa	Florida S	Statewide	Nationa	l Norms
		Middle School	High School	Middle School	High School	Middle School	High School
Community	Community Disorganization	37	39	42	44	47	47
	Transitions and Mobility	60	67	59	61	47	46
	Laws and Norms Favorable to Drug Use	34	30	37	31	42	42
	Perceived Availability of Drugs	42	28	37	27	45	45
	Perceived Availability of Handguns	34	39	24	36	25	42
Family	Poor Family Management	33	32	40	38	44	45
	Family Conflict	39	34	38	33	42	37
School	Poor Academic Performance	41	45	42	44	45	48
	Lack of Commitment to School	50	54	53	54	47	46
Peer and	Favorable Attitudes toward Antisocial Behavior	39	36	39	35	40	46
Individual	Favorable Attitudes toward ATOD Use	30	38	32	36	39	45
	Early Initiation of Drug Use	23	22	23	22	41	46
Average Preva	lence Rate	38	39	39	38	42	45

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values. The symbol "--" indicates that data are not available.

Table 23. Protective factor prevalence rate trends among middle school students for Okaloosa, 2006 to 2016

Domain	Scale			Oka	loosa		
		2006	2008	2010	2012	2014	2016
Community	Community Rewards for Prosocial Involvement		59	57	58	58	52
Family	Family Opportunities for Prosocial Involvement		54	56	60	67	63
	Family Rewards for Prosocial Involvement		52	55	54	61	58
School	School Opportunities for Prosocial Involvement		51	52	59	61	62
	School Rewards for Prosocial Involvement		47	50	52	59	52
Peer and Individual	Religiosity		62	59	57	52	56
Average Prevalence R	ate		54	55	57	60	57

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values. The symbol "--" indicates that data are not available.

Table 24. Risk factor prevalence rate trends among middle school students for Okaloosa, 2006 to 2016

Domain	Scale			Oka	loosa		
		2006	2008	2010	2012	2014	2016
Community	Community Disorganization		43	45	40	36	37
	Transitions and Mobility		61	64	62	56	60
	Laws and Norms Favorable to Drug Use		40	39	34	32	34
	Perceived Availability of Drugs		52	48	39	35	42
	Perceived Availability of Handguns		31	31	29	28	34
Family	Poor Family Management		46	39	37	26	33
	Family Conflict		45	42	41	31	39
School	Poor Academic Performance		43	41	49	40	41
	Lack of Commitment to School		55	50	48	44	50
Peer and	Favorable Attitudes toward Antisocial Behavior		45	44	37	32	39
Individual	Favorable Attitudes toward ATOD Use		36	39	31	23	30
	Early Initiation of Drug Use		36	35	26	21	23
Average Preval	lence Rate		41	42	37	34	38

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.

Table 25. Protective factor prevalence rate trends among high school students for Okaloosa, 2006 to 2016

Domain	Scale	Okaloosa						
		2006	2008	2010	2012	2014	2016	
Community	Community Rewards for Prosocial Involvement		65	67	65	71	75	
Family	Family Opportunities for Prosocial Involvement		54	58	57	62	61	
	Family Rewards for Prosocial Involvement		54	58	56	61	57	
School	School Opportunities for Prosocial Involvement		61	64	71	68	70	
	School Rewards for Prosocial Involvement		60	64	63	70	60	
Peer and Individual	Religiosity		67	66	67	68	59	
Average Prevalence Rate			60	63	63	67	64	

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values. The symbol "--" indicates that data are not available.

Table 26. Risk factor prevalence rate trends among high school students for Okaloosa, 2006 to 2016

Domain	Scale	Okaloosa						
		2006	2008	2010	2012	2014	2016	
Community	Community Disorganization		36	37	37	35	39	
	Transitions and Mobility		66	66	63	62	67	
	Laws and Norms Favorable to Drug Use		32	30	26	25	30	
	Perceived Availability of Drugs		47	35	30	27	28	
	Perceived Availability of Handguns		47	39	39	38	39	
Family	Poor Family Management		44	38	36	27	32	
	Family Conflict		39	34	34	31	34	
School	Poor Academic Performance		46	43	46	35	45	
	Lack of Commitment to School		49	51	43	50	54	
Peer and	Favorable Attitudes toward Antisocial Behavior		50	39	35	31	36	
Individual	Favorable Attitudes toward ATOD Use		45	41	35	29	38	
	Early Initiation of Drug Use		36	30	26	21	22	
Average Prevalence Rate			46	43	39	34	39	

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.