2016 Florida Youth Substance Abuse Survey

Table 1. Major demographic characteristics of surveyed Pasco and Florida Statewide, 2016

	Pas	sco	Florida St	tatewide
	N	%	N	%
Sex				
Female	735	46.8	31,515	47.9
Male	790	50.4	32,905	50.0
Race/Ethnic group				
African American	86	5.5	14,666	22.3
American Indian	26	1.7	568	0.9
Asian	32	2.0	743	1.1
Hispanic/Latino	204	13.0	13,174	20.0
Native Hawaiian/Pacific Islander	7	0.4	144	0.2
Other/Multiple	315	20.1	7,523	11.4
White, non-Hispanic	875	55.8	28,309	43.0
Age				
10	1	0.1	65	0.1
11	143	9.1	3,856	5.9
12	270	17.2	8,338	12.7
13	304	19.4	9,230	14.0
14	243	15.5	9,454	14.4
15	210	13.4	10,070	15.3
16	151	9.6	9,684	14.7
17	154	9.8	9,348	14.2
18	78	5.0	4,799	7.3
19 or older	8	0.5	618	0.9
Grade				
6th	326	20.8	9,301	14.1
7th	261	16.6	9,215	14.0
8th	320	20.4	9,326	14.2
9th	179	11.4	10,140	15.4
10th	197	12.6	9,834	15.0
11th	121	7.7	9,254	14.1
12th	165	10.5	8,705	13.2
Overall Middle School	907	57.8	27,678	42.1
Overall High School	662	42.2	37,765	57.4
Total	1,569	100.0	65,776	100.0

Note: Some categories do not sum to 100% of the total due to missing values (e.g., not all survey questions were answered). In addition, rounding can produce totals that do not equal 100%. "N" represents the number of valid cases. In this table, county data are unweighted while statewide data are weighted.

Table 2. Percentages of Pasco and Florida Statewide youth who reported having used various drugs in their lifetimes, 2016

				Pasco						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Alcohol	21.0	52.5	42.4	35.7	21.6	54.3	39.0	21.8	51.4	41.3	37.1	23.6	51.5	39.1
Blacking Out		17.0	17.3	16.5		19.2	17.0		15.9	16.5	15.4		16.1	15.9
Cigarettes	6.4	23.4	17.6	14.3	6.4	25.4	16.0	8.0	18.5	14.0	14.1	8.2	18.5	14.1
Vaporizer/E-Cigarette	12.8	37.6	25.7	28.1	13.7	39.5	26.8	14.6	33.9	24.4	27.1	15.7	34.5	25.8
Marijuana or Hashish	6.7	34.4	22.7	22.3	7.0	36.9	22.4	7.0	31.7	21.4	21.3	8.1	31.8	21.3
Synthetic Marijuana		7.1	7.4	6.5		7.2	7.1		4.9	4.8	5.0		4.8	4.9
Inhalants	5.1	2.4	3.1	4.1	4.6	2.8	3.6	6.4	4.6	6.1	4.7	6.4	4.7	5.4
Flakka		0.1	0.2	0.0		0.1	0.1		1.0	0.9	1.1		1.0	1.0
Club Drugs	0.6	2.5	2.0	1.5	1.1	2.6	1.7	0.7	3.0	1.9	2.2	0.8	2.9	2.1
LSD, PCP or Mushrooms	0.9	5.6	3.4	3.9	0.8	6.5	3.5	1.0	4.8	2.7	3.7	1.1	4.9	3.2
Methamphetamine	0.2	1.0	0.9	0.5	0.6	0.9	0.7	0.5	0.8	0.6	0.8	0.5	0.8	0.7
Cocaine or Crack Cocaine	0.9	2.6	2.2	1.7	0.7	3.0	1.9	0.8	2.5	1.6	1.9	0.8	2.3	1.8
Heroin	0.4	0.6	0.8	0.2	0.5	0.6	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Depressants	2.5	8.1	7.0	4.2	2.6	9.2	5.7	2.2	6.5	5.3	4.1	2.4	6.6	4.7
Prescription Pain Relievers	4.4	7.8	7.2	5.2	4.2	8.4	6.3	3.3	5.8	5.3	4.2	3.5	5.7	4.8
Prescription Amphetamines	0.9	5.9	3.9	3.3	0.8	6.8	3.7	1.3	4.7	3.3	3.2	1.4	4.6	3.2
Steroids (without a doctor's order)	0.5	0.5	0.3	0.8	0.4	0.7	0.5	0.6	0.5	0.4	0.6	0.5	0.5	0.5
Over-the-Counter Drugs	3.4	7.5	5.8	5.6	3.6	8.1	5.7	3.2	5.3	4.7	4.2	3.4	5.4	4.4
Needle to Inject Illegal Drugs		1.3	2.2	0.5		1.2	1.3		0.8	0.6	0.8		0.8	0.8
Any illicit drug	15.2	38.1	29.4	27.1	15.1	40.0	28.2	15.7	36.4	29.0	26.4	16.6	36.6	27.7
Any illicit drug other than marijuana	11.6	19.8	16.3	16.0	11.1	21.3	16.2	12.0	17.7	16.4	14.2	12.3	17.7	15.3
Alcohol only	12.5	21.1	19.0	15.7	13.2	21.4	17.4	12.5	21.0	18.1	17.0	13.4	21.0	17.5
Alcohol or any illicit drug	27.3	58.9	47.9	42.5	27.9	61.2	45.2	27.8	57.2	46.8	43.1	29.6	57.4	44.8
Any illicit drug, but no alcohol	6.9	6.4	6.0	7.2	6.7	7.0	6.6	6.5	6.1	6.0	6.5	6.5	6.1	6.3

Note: The first 19 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. Respondents were asked on how many occasions in their lifetime they woke up after a night of drinking and did not remember the things they did or the places they went. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "--" indicates that data are not available.

Table 3. Percentages of Pasco and Florida Statewide youth who reported having used various drugs in the past 30 days, 2016

				Pasco						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Alcohol	9.2	28.5	21.5	19.0	9.1	31.2	20.2	8.3	25.5	19.9	17.0	9.1	25.2	18.3
Binge Drinking	2.3	11.0	6.7	7.6	2.2	11.9	7.3	3.2	10.9	7.9	7.7	3.4	10.8	7.7
Cigarettes	1.5	7.4	5.6	3.6	1.3	7.9	4.8	1.4	4.8	3.3	3.5	1.5	4.6	3.4
Vaporizer/E-Cigarette	5.3	16.9	10.5	12.7	5.7	17.4	11.8	5.1	12.8	8.4	10.6	5.5	12.9	9.6
Marijuana or Hashish	3.2	19.7	12.3	12.7	3.5	20.8	12.5	3.2	17.0	10.9	11.5	3.9	16.9	11.2
Synthetic Marijuana		0.6	0.5	0.7		0.7	0.6		1.0	0.9	1.2		1.1	1.0
Inhalants	1.4	0.8	1.0	1.1	1.1	0.9	1.0	2.2	1.2	1.9	1.4	2.1	1.2	1.6
Flakka		0.0	0.0	0.0		0.0	0.0		0.5	0.4	0.7		0.5	0.5
Club Drugs	0.4	1.0	1.1	0.4	0.7	0.9	0.7	0.3	0.8	0.5	0.6	0.3	0.7	0.6
LSD, PCP or Mushrooms	0.3	1.1	0.7	0.8	0.2	1.3	0.7	0.3	1.4	0.7	1.1	0.3	1.4	0.9
Methamphetamine	0.0	0.6	0.7	0.0	0.4	0.3	0.3	0.3	0.4	0.3	0.4	0.3	0.4	0.4
Cocaine or Crack Cocaine	0.1	0.4	0.2	0.3	0.1	0.5	0.3	0.3	0.8	0.6	0.6	0.3	0.8	0.6
Heroin	0.1	0.1	0.1	0.1	0.2	0.0	0.1	0.1	0.2	0.1	0.2	0.1	0.2	0.2
Depressants	1.1	2.3	2.6	0.7	1.4	2.5	1.8	0.8	2.4	2.0	1.5	0.9	2.5	1.8
Prescription Pain Relievers	2.5	2.0	3.2	1.0	2.6	1.7	2.2	1.6	2.0	2.2	1.5	1.6	2.1	1.8
Prescription Amphetamines	0.2	2.1	1.1	1.3	0.3	2.3	1.3	0.5	1.6	1.1	1.2	0.6	1.7	1.2
Steroids (without a doctor's order)	0.5	0.6	0.4	0.6	0.5	0.7	0.5	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Over-the-Counter Drugs	2.0	3.7	2.9	2.9	2.4	3.6	3.0	1.8	2.1	2.0	1.9	1.8	2.1	2.0
Any illicit drug	7.1	23.5	16.9	15.5	7.7	23.7	16.3	7.5	20.0	15.1	14.2	8.1	20.0	14.7
Any illicit drug other than marijuana	5.5	9.2	8.8	6.1	5.4	9.8	7.6	5.6	7.7	7.3	6.2	5.7	7.7	6.8
Alcohol only	6.2	14.1	10.9	10.6	6.0	16.1	10.6	5.2	13.4	11.1	9.1	5.6	13.3	10.0
Alcohol or any illicit drug	13.0	37.1	27.8	25.5	13.4	39.4	26.6	12.5	32.9	25.8	22.9	13.5	32.9	24.3
Any illicit drug, but no alcohol	4.1	8.7	6.6	6.7	4.4	8.4	6.7	4.5	7.8	6.4	6.3	4.7	8.0	6.4

Note: The first 18 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. Binge drinking is defined as having had five or more alcoholic drinks in a row in the past two weeks. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "--" indicates that data are not available.

Table 4. Lifetime trend in alcohol, tobacco and other drug use for Pasco, 2006 to 2016

		2006			2008			2010			2012			2014			2016	
	Middle School	0	Total	Middle School		Total	Middle School		Total	Middle School		Total	Middle School	U	Total	Middle School		Total
Alcohol	37.7	72.8	57.6	35.2	69.1	53.9	34.3	67.9	53.1	24.5	62.5	45.6	27.3	55.5	43.2	21.0	52.5	39.0
Blacking Out														21.9	21.9		17.0	17.0
Cigarettes	19.5	41.7	32.1	19.9	38.9	30.4	18.5	37.6	29.2	11.8	27.3	20.4	10.4	24.3	18.2	6.4	23.4	16.0
Vaporizer/E-Cigarette																12.8	37.6	26.8
Marijuana or Hashish	9.1	35.1	23.9	9.3	34.8	23.5	11.2	40.2	27.4	9.3	32.8	22.4	8.8	34.5	23.3	6.7	34.4	22.4
Synthetic Marijuana											17.7	17.7		10.0	10.0		7.1	7.1
Inhalants	13.7	12.7	13.2	11.3	9.8	10.4	12.3	8.5	10.1	9.6	6.7	8.0	8.5	6.2	7.2	5.1	2.4	3.6
Flakka																	0.1	0.1
Club Drugs				2.3		2.3	1.9	4.5	3.3	1.2	4.9	3.3	0.5	4.0	2.5	0.6	2.5	1.7
LSD, PCP or Mushrooms				1.5		1.5	2.0	4.8	3.6	1.7	5.6	3.9	1.2	6.1	4.0	0.9	5.6	3.5
Methamphetamine	1.5	3.0	2.4	1.0	2.0	1.6	1.2	0.8	1.0	0.5	1.3	0.9	1.6	1.0	1.3	0.2	1.0	0.7
Cocaine or Crack Cocaine				2.1		2.1	2.0	2.6	2.3	1.3	3.3	2.4	1.1	1.5	1.3	0.9	2.6	1.9
Heroin	1.1	1.9	1.6	0.9	1.5	1.3	1.3	0.8	1.0	0.7	1.1	0.9	0.8	1.4	1.1	0.4	0.6	0.5
Depressants	3.9	12.9	9.0	3.8	12.0	8.4	3.7	7.2	5.6	1.3	7.2	4.6	1.6	6.2	4.2	2.5	8.1	5.7
Prescription Pain Relievers	6.4	16.2	12.0	6.5	14.2	10.8	4.7	12.4	9.0	4.9	8.5	6.9	2.5	9.3	6.4	4.4	7.8	6.3
Prescription Amphetamines	2.4	7.4	5.3	1.6	3.7	2.7	1.7	3.4	2.6	1.4	3.8	2.8	1.6	4.2	3.1	0.9	5.9	3.7
Steroids (without a doctor's order)	1.7	1.2	1.4	0.6	1.0	0.8	0.9	1.1	1.0	0.6	0.8	0.7	0.8	0.4	0.5	0.5	0.5	0.5
Over-the-Counter Drugs				3.8		3.8	4.8	8.9	7.1	4.0	6.5	5.4	3.0	7.0	5.2	3.4	7.5	5.7
Needle to Inject Drugs																	1.3	1.3
Any illicit drug	22.4	44.3	34.9	19.8	40.6	31.3	22.2	45.5	35.1	19.1	39.7	30.5	17.6	39.9	30.2	15.2	38.1	28.2
Any illicit drug other than marijuana	19.1	28.6	24.5	16.1	25.7	21.4	18.4	25.0	22.1	15.4	20.8	18.4	13.3	20.1	17.1	11.6	19.8	16.2
Alcohol only	20.3	32.9	27.4	20.6	30.5	26.1	17.5	25.3	21.8	11.9	26.9	20.2	14.6	20.0	17.7	12.5	21.1	17.4
Alcohol or any illicit drug	42.6	77.0	62.1	40.2	71.2	57.3	39.7	70.8	57.0	30.7	66.6	50.6	32.1	59.9	47.7	27.3	58.9	45.2
Any illicit drug, but no alcohol	5.2	4.3	4.7	5.2	2.2	3.6	5.7	3.1	4.3	6.2	4.3	5.1	5.0	4.5	4.7	6.9	6.4	6.6

Note: The first 19 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. Respondents were asked on how many occasions in their lifetime they woke up after a night of drinking and did not remember the things they did or the places they went. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "--" indicates that data are not available.

2016 Florida Youth Substance Abuse Survey - Pasco Report

Table 5. Past-30-day trend in alcohol, tobacco and other drug use for Pasco, 2006 to 2016

		2006			2008			2010			2012			2014			2016	
	Middle School	O	Total	Middle School		Total	Middle School		Total	Middle School		Total	Middle School	. 0	Total	Middle School		Total
Alcohol	16.6	46.4	33.5	15.1	42.0	30.0	16.2	39.3	29.2	13.8	32.6	24.3	11.8	28.8	21.3	9.2	28.5	20.2
Binge Drinking	8.2	29.4	20.3	6.0	23.7	15.8	7.2	18.5	13.5	5.6	13.5	10.0	4.5	12.2	8.8	2.3	11.0	7.3
Cigarettes	6.8	17.5	12.8	5.9	17.4	12.3	5.8	16.6	11.8	4.2	9.7	7.2	2.5	7.3	5.2	1.5	7.4	4.8
Vaporizer/E-Cigarette																5.3	16.9	11.8
Marijuana or Hashish	5.5	19.5	13.6	4.7	19.5	12.9	6.7	21.1	14.8	5.0	16.9	11.6	5.1	20.1	13.6	3.2	19.7	12.5
Synthetic Marijuana											5.4	5.4		2.1	2.1		0.6	0.6
Inhalants	5.1	5.2	5.2	3.8	1.1	2.3	4.9	1.7	3.1	2.4	1.5	1.9	2.8	1.8	2.2	1.4	0.8	1.0
Flakka																	0.0	0.0
Club Drugs				0.7		0.7	0.6	1.1	0.9	0.4	2.0	1.3	0.1	0.7	0.5	0.4	1.0	0.7
LSD, PCP or Mushrooms				1.0		1.0	0.6	1.0	0.8	1.0	1.9	1.5	0.1	1.3	0.8	0.3	1.1	0.7
Methamphetamine	0.4	0.6	0.5	0.1	0.5	0.4	0.3	0.5	0.4	0.3	1.0	0.7	1.0	0.3	0.6	0.0	0.6	0.3
Cocaine or Crack Cocaine				0.4		0.4	0.7	1.0	0.9	0.3	1.5	1.0	0.2	0.6	0.4	0.1	0.4	0.3
Heroin	0.4	1.0	0.7	0.4	0.6	0.5	0.6	0.5	0.5	0.1	0.6	0.4	0.4	0.3	0.3	0.1	0.1	0.1
Depressants	0.8	6.5	4.1	1.5	3.5	2.7	0.7	2.3	1.6	0.6	2.4	1.6	0.6	2.3	1.6	1.1	2.3	1.8
Prescription Pain Relievers	2.4	6.9	5.0	3.2	3.8	3.5	1.5	4.1	3.0	1.9	2.8	2.4	1.1	3.1	2.2	2.5	2.0	2.2
Prescription Amphetamines	0.8	2.2	1.6	0.6	0.8	0.7	0.8	0.7	0.8	0.6	1.2	0.9	0.5	0.5	0.5	0.2	2.1	1.3
Steroids (without a doctor's order)	0.7	0.1	0.4	0.4	0.4	0.4	0.3	0.4	0.4	0.2	0.4	0.3	0.3	0.0	0.1	0.5	0.6	0.5
Over-the-Counter Drugs				1.7		1.7	2.1	3.7	3.0	1.9	2.5	2.2	1.6	2.9	2.4	2.0	3.7	3.0
Any illicit drug	11.2	24.8	19.1	8.6	22.3	16.2	12.2	25.1	19.4	8.9	21.3	15.8	9.1	24.4	17.8	7.1	23.5	16.3
Any illicit drug other than marijuana	8.4	14.4	11.9	6.5	9.9	8.4	8.2	9.7	9.1	5.9	9.0	7.6	5.6	9.1	7.6	5.5	9.2	7.6
Alcohol only	9.8	25.5	18.7	9.6	24.9	18.1	8.9	20.8	15.5	8.6	17.7	13.6	7.8	14.1	11.4	6.2	14.1	10.6
Alcohol or any illicit drug	20.4	50.2	37.3	18.0	47.0	34.0	20.6	45.6	34.6	17.3	38.8	29.2	16.7	38.1	28.7	13.0	37.1	26.6
Any illicit drug, but no alcohol	4.0	4.6	4.4	3.0	5.2	4.3	4.6	6.4	5.6	3.7	6.5	5.2	4.9	9.4	7.5	4.1	8.7	6.7

Note: The first 18 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. Binge drinking is defined as having had five or more alcoholic drinks in a row in the past two weeks. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "--" indicates that data are not available.

Table 6. Percentages of Pasco and Florida Statewide youth who reported early ATOD use, perceived risk of harm, personal disapproval, parental disapproval and peer disapproval, 2016

				Pasco						Flori	da State	ewide		
	Middle School		Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Early ATOD use														
More than a sip of alcohol		20.8	21.7	19.1		21.4	20.8		19.4	18.3	20.2		19.3	19.4
Drinking at least once a month		3.2	4.5	2.0		3.9	3.2		3.5	3.7	3.4		3.5	3.5
Smoke cigarettes		10.7	11.4	9.9		11.2	10.7		9.3	8.6	9.8		9.4	9.3
Smoke marijuana		12.3	11.6	13.0		12.9	12.3		10.6	9.0	12.0		10.8	10.6
Great risk of harm														
1 or 2 drinks nearly every day	46.6	34.5	42.7	36.7	45.6	35.8	39.7	45.7	40.7	46.4	39.3	45.9	40.0	42.8
5+ drinks once or twice a week	59.1	50.8	57.6	51.2	58.4	51.9	54.4	58.1	52.3	58.4	51.2	58.4	51.8	54.7
1+ packs of cigarettes per day	69.2	67.3	67.9	68.5	68.7	67.3	68.1	67.9	68.8	68.4	68.5	68.7	68.2	68.4
Marijuana once or twice a week	54.9	21.9	35.7	36.2	52.1	21.2	36.2	51.7	25.2	37.5	35.2	50.1	24.4	36.3
Try marijuana once or twice	35.5	14.2	22.4	24.1	33.6	14.0	23.4	36.1	16.8	24.8	25.0	34.7	16.4	24.9
Use prescription drugs	73.6	67.2	71.0	68.6	72.6	66.8	70.0	69.1	68.2	70.0	67.3	69.4	67.9	68.5
Think it would be wrong to														
Smoke cigarettes	96.8	85.3	91.0	89.9	96.1	86.8	90.2	95.6	87.7	91.1	91.0	95.3	88.9	91.0
Drink alcohol regularly	90.5	61.3	73.1	74.7	88.4	61.8	73.9	88.8	64.5	74.1	75.2	87.6	64.8	74.7
Smoke marijuana	91.4	55.7	71.0	70.7	88.6	55.1	70.9	89.7	62.3	74.4	73.3	88.1	62.3	73.8
Smoke synthetic marijuana	94.8	89.9	91.8	92.4	94.4	90.1	92.0	93.1	88.6	90.1	90.9	92.7	88.6	90.5
Use other illicit drugs	98.4	92.7	96.1	94.3	97.6	92.4	95.2	97.5	93.6	95.8	94.7	97.3	93.6	95.2
Parents think it would be wrong to														
1 or 2 drinks nearly every day	97.9	93.7	96.1	95.1	97.6	94.1	95.5	97.3	93.8	95.4	95.1	97.2	94.1	95.2
Smoke cigarettes	99.3	95.6	96.8	97.7	99.0	95.8	97.2	98.6	96.8	97.6	97.5	98.6	97.3	97.6
Smoke marijuana	96.8	81.6	87.6	88.6	96.2	81.0	88.2	97.0	88.6	92.3	92.0	96.6	88.9	92.1
Use prescription drugs	98.1	97.3	97.1	98.1	98.0	97.5	97.7	98.2	97.3	97.4	97.9	98.1	97.4	97.7
Friends think it would be wrong to														
1 or 2 drinks nearly every day	92.8	74.9	85.1	80.4	92.3	74.5	82.6	91.3	79.0	85.6	82.9	90.7	79.1	84.2
Smoke tobacco	96.9	85.3	91.0	89.9	96.6	86.0	90.3	94.8	87.0	91.1	89.5	94.4	87.8	90.3
Smoke marijuana	90.5	56.3	73.0	68.6	87.9	56.1	71.1	88.6	59.9	72.6	71.3	86.6	59.8	72.0
Use prescription drugs	96.9	91.4	93.5	94.2	96.8	90.4	93.8	95.7	90.6	93.4	92.2	95.5	90.3	92.7

Note: Early ATOD use is defined as the percentage of high school students who started using at age 13 or younger. The symbol "--" indicates that data are not available.

2016 Florida Youth Substance Abuse Survey - Pasco Report

Table 7. Trends in early ATOD use and attitudes toward substance use for Pasco, 2006 to 2016

		2006			2008			2010			2012			2014			2016	
	Middle School		Total															
Early ATOD use																		
Try alcohol		36.3	36.3		33.8	33.8		27.0	27.0		22.7	22.7		20.1	20.1		20.8	20.8
Drink monthly		8.1	8.1		6.2	6.2		5.6	5.6		3.2	3.2		2.1	2.1		3.2	3.2
Smoke cigarettes		26.6	26.6		24.7	24.7		21.0	21.0		14.9	14.9		12.5	12.5		10.7	10.7
Smoke marijuana		13.6	13.6		11.8	11.8		13.8	13.8		11.1	11.1		12.1	12.1		12.3	12.3
Great risk of harm																		
Daily alcohol use	38.7	35.0	36.6	41.4	37.3	39.2	44.6	37.1	40.4	44.6	37.7	40.8	39.2	37.5	38.2	46.6	34.5	39.7
5+ drinks weekly													55.2	49.3	52.0	59.1	50.8	54.4
Daily cigarette use	63.6	64.9	64.4	71.1	67.1	68.9	70.7	67.3	68.8	71.0	66.9	68.7	66.9	68.8	68.0	69.2	67.3	68.1
Marijuana weekly*													48.7	23.4	34.6	54.9	21.9	36.2
Try marijuana	45.5	25.0	33.9	41.8	24.2	32.0	38.9	14.6	25.3	37.1	17.2	26.0	32.5	14.5	22.5	35.5	14.2	23.4
Prescription drugs											76.0	76.0	72.8	73.8	73.3	73.6	67.2	70.0
Think it wrong																		
Smoke cigarettes	88.7	67.7	76.9	88.8	70.6	78.7	89.1	67.8	77.2	93.4	78.9	85.3	93.3	84.3	88.3	96.8	85.3	90.2
Drink regularly	80.5	48.1	62.2	82.2	51.6	65.2	80.0	50.8	63.7	87.2	60.8	72.5	87.0	64.6	74.4	90.5	61.3	73.9
Smoke marijuana	91.9	69.8	79.4	91.4	70.1	79.5	87.1	64.8	74.6	91.2	67.7	78.1	88.2	62.6	73.9	91.4	55.7	70.9
Smoke synthetic marijuana																94.8	89.9	92.0
Use other illicit drugs	96.6	91.4	93.7	96.9	93.4	95.0	96.4	92.9	94.4	98.1	94.7	96.2	96.6	93.9	95.1	98.4	92.7	95.2
Parents think it wrong																		
Daily alcohol use*													96.0	92.6	94.1	97.9	93.7	95.5
Smoke cigarettes	95.3	86.1	90.0	96.4	89.9	92.8	97.5	89.4	93.0	97.7	92.1	94.5	97.7	94.9	96.2	99.3	95.6	97.2
Smoke marijuana	97.6	91.4	94.0	97.7	91.6	94.3	96.1	89.2	92.2	97.4	90.5	93.5	96.5	89.0	92.3	96.8	81.6	88.2
Prescription drugs													98.6	98.5	98.5	98.1	97.3	97.7
Friends think it wrong																		
Daily alcohol use													89.7	75.5	81.7	92.8	74.9	82.6
Smoke tobacco													93.8	82.4	87.4	96.9	85.3	90.3
Smoke marijuana													87.4	59.1	71.5	90.5	56.3	71.1
Prescription drugs													96.4	92.3	94.1	96.9	91.4	93.8

Note: Early ATOD use is defined as the percentage of high school students who started using at age 13 or younger. The symbol "--" indicates that data are not available.

^{*} In 2014, the description of marijuana use in the risk of harm question was changed from "regularly" to "once or twice a week." Also, in 2014, the description of alcohol use in the parent disapproval question was changed from "regularly" to "nearly every day." For these items, care should be exercised when comparing 2014 data to previous years.

Table 8. Among high school drinkers, usual source of alcohol within the past 30 days, Pasco and Florida Statewide youth, 2016

				Pasco						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Bought in a store		8.2	5.5	11.9		8.6	8.2		8.3	5.0	12.4		7.3	8.3
Bought in a restaurant, bar, or club		1.4	1.0	0.0		1.0	1.4		1.6	1.2	2.1		1.5	1.6
Bought at a public event		0.0	0.0	0.0		0.0	0.0		0.7	0.7	0.8		0.6	0.7
Someone bought it for me		16.0	13.4	20.0		15.9	16.0		14.7	13.3	16.3		14.5	14.7
Someone gave it to me		49.4	55.7	41.3		49.1	49.4		44.8	50.6	37.8		47.0	44.8
Took it from a store		0.0	0.0	0.0		0.0	0.0		0.3	0.2	0.4		0.4	0.3
Took it from a family member		10.6	11.3	10.1		11.7	10.6		11.7	14.2	8.7		12.4	11.7
Some other way		14.4	13.2	16.6		13.7	14.4		17.8	14.7	21.5		16.5	17.8

Table 9. Among $\underline{\text{high school}}$ drinkers, usual drinking location within the past 30 days, Pasco and Florida Statewide youth, 2016

				Pasco						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
My home		39.4	41.0	39.5		37.9	39.4		37.7	39.2	35.9		37.1	37.7
Another person's home		38.4	39.8	35.9		40.1	38.4		40.0	41.0	38.9		40.8	40.0
Car or other vehicle		2.9	3.1	2.7		3.0	2.9		1.7	1.6	1.9		1.7	1.7
Restaurant, bar, or club		1.6	0.0	1.5		1.8	1.6		2.7	2.9	2.5		2.4	2.7
Public place		4.9	4.5	5.7		5.5	4.9		3.9	3.9	3.7		4.3	3.9
Public event		0.8	1.5	0.0		0.9	0.8		1.6	1.8	1.5		1.7	1.6
School property		1.2	0.0	2.7		1.4	1.2		1.3	0.7	2.0		1.4	1.3
Some other place		10.8	10.2	12.0		9.4	10.8		11.1	9.0	13.7		10.6	11.1

Table 10. Among <u>high school</u> drinkers, number of drinks usually consumed per day, on the days students drank in the past 30 days, Pasco and Florida Statewide youth, 2016

				Pasco						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
1		26.7	22.9	32.0		27.6	26.7		29.6	31.6	27.4		30.5	29.6
2		20.8	24.2	17.9		18.6	20.8		22.5	24.8	19.8		22.4	22.5
3		19.6	22.7	17.0		20.1	19.6		17.3	17.9	16.3		16.9	17.3
4		7.2	9.2	5.3		8.5	7.2		8.6	9.0	8.3		8.5	8.6
5 or more		25.7	21.0	27.8		25.3	25.7		22.0	16.7	28.2		21.8	22.0

Table 11. Percentages of Pasco and Florida Statewide youth who reported drinking alcohol, smoking marijuana, or using another drug to get high $\underline{\text{before or during school}}$ in the past 12 months, 2016

				Pasco						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Drinking alcohol	3.3	7.8	5.7	5.4	3.3	8.8	5.9	3.3	6.5	5.5	4.7	3.3	6.8	5.1
Smoking marijuana	3.3	13.7	9.0	9.5	4.1	14.5	9.2	3.2	12.2	8.1	8.7	3.7	12.3	8.5
Used another drug to get high	1.4	4.0	2.9	2.3	1.6	4.5	2.9	1.4	3.9	2.9	2.9	1.5	4.1	2.9

Table 12. Percentages of Pasco and Florida Statewide youth who reported engaging in delinquent behavior within the past 12 months, 2016

				Pasco						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Carrying a handgun	4.6	4.6	1.7	7.4	4.7	4.3	4.6	5.0	5.8	2.6	8.3	5.0	5.9	5.5
Selling drugs	1.6	6.3	3.2	5.5	1.9	6.3	4.3	1.6	6.0	3.0	5.4	1.8	6.1	4.2
Attempting to steal a vehicle	0.4	2.0	0.7	2.0	0.7	2.1	1.3	1.0	1.5	0.8	1.7	1.0	1.5	1.3
Being arrested	1.6	3.6	1.4	4.1	1.8	3.7	2.7	1.9	2.8	1.8	3.0	1.8	3.0	2.4
Taking a handgun to school	0.0	1.2	0.4	1.1	0.4	1.1	0.7	0.4	0.7	0.3	0.9	0.4	0.7	0.6
Getting suspended	7.0	9.5	4.5	12.2	7.9	9.6	8.4	10.6	9.1	7.0	12.5	10.1	9.8	9.8
Attacking someone with intent to harm	4.7	7.0	4.3	8.0	4.8	6.9	6.0	6.2	6.1	5.3	6.9	6.0	6.6	6.1

 $Table\ 13.\ Trends\ in\ delinquent\ behaviors\ for\ Pasco,\ 2006\ to\ 2016$

		2006			2008			2010			2012			2014			2016	
	Middle School	High School	Total	I	High School	Total												
Carrying a handgun	3.5	6.5	5.3	3.0	6.1	4.7	6.4	3.8	4.9	3.4	3.4	3.4	4.9	6.3	5.7	4.6	4.6	4.6
Selling drugs	1.8	9.8	6.5	3.4	7.8	5.8	3.5	10.0	7.2	2.0	6.2	4.4	2.8	7.8	5.6	1.6	6.3	4.3
Attempting to steal a vehicle	2.4	4.6	3.7	2.3	3.2	2.7	1.8	3.2	2.6	1.0	2.2	1.7	0.7	1.4	1.1	0.4	2.0	1.3
Being arrested	4.6	8.2	6.7	4.5	6.4	5.5	4.1	4.9	4.5	1.9	4.0	3.1	1.4	2.7	2.2	1.6	3.6	2.7
Taking a handgun to school	0.8	1.8	1.4	0.6	0.6	0.6	0.6	0.5	0.6	0.3	0.7	0.5	0.9	0.7	0.8	0.0	1.2	0.7
Getting suspended	14.2	15.9	15.1	14.7	15.0	14.9	12.3	13.6	13.1	10.5	8.3	9.3	9.5	9.2	9.4	7.0	9.5	8.4
Attacking someone with intent to harm	13.4	18.1	16.1	10.7	12.6	11.8	11.0	10.1	10.5	8.9	5.5	7.0	6.5	6.0	6.2	4.7	7.0	6.0

Table 14. Percentages of Pasco and Florida Statewide youth with family members who have been or are now in jail or friends who have been in trouble because of alcohol or drugs, 2016

				Pasco						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Family members who have been or are now in jail or prison	38.6	48.0	48.6	39.1	39.1	50.2	44.0	44.4	44.6	47.8	41.2	44.2	45.4	44.5
Friends who have have been or are in trouble because of alcohol or drugs	21.3	51.0	40.9	36.0	23.0	52.3	38.2	23.9	46.5	38.4	35.8	26.0	47.1	37.1

Table 15. Percentages of Pasco and Florida Statewide youth who reported involvement in bullying behavior, 2016

				Pasco						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Skipped school because of bullying	7.6	11.9	13.9	5.9	8.2	12.3	10.0	7.6	8.2	11.5	4.4	7.6	8.3	7.9
Was kicked or shoved in past 30 days	19.7	7.7	12.0	13.4	18.7	8.1	12.9	19.6	8.1	12.0	13.8	18.8	8.1	13.0
Was taunted or teased in past 30 days	39.2	25.6	34.0	28.2	38.4	26.0	31.5	35.3	22.5	30.1	25.6	34.4	22.8	27.9
Was victim of cyber bullying in past 30 days	9.0	8.5	12.6	5.1	9.4	8.9	8.7	7.8	7.6	10.2	5.2	7.8	7.9	7.7
Physically bullied others in past 30 days	5.9	2.8	3.0	5.1	5.2	3.5	4.1	7.6	4.9	5.0	7.0	7.2	5.2	6.1
Verbally bullied others in past 30 days	10.6	9.2	9.5	10.2	10.1	10.3	9.8	14.3	10.4	11.0	13.0	14.0	10.7	12.0
Cyber bullied others in past 30 days	2.1	4.5	3.5	3.6	2.1	5.1	3.5	3.6	4.1	4.0	3.7	3.7	4.1	3.9

Table 16. Percentages of Pasco and Florida Statewide youth who reported gang membership, 2016

				Pasco						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Have you ever belonged to a gang?	2.6	4.3	1.7	5.0	3.0	3.6	3.5	3.3	3.5	2.3	4.5	3.1	3.6	3.4
Did that gang have a name?	24.9	18.6	13.3	23.6	27.8	14.3	20.3	23.4	14.1	13.1	19.7	21.9	14.4	16.9
Are you a gang member now?		2.5	1.7	2.9		2.2	2.5		2.0	1.1	2.8		2.0	2.0

Note: The prevalence rates for "Did that gang have a name?" exclude students who reported that they have never belonged to a gang. The symbol "--" indicates that data are not available.

Table 17. Reasons for joining a gang, reported by <u>high school students who have belonged to a gang</u>, among Pasco and Florida Statewide youth, 2016

				Pasco						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Fun and excitement		5.4	6.6	3.1		3.3	5.4		5.0	4.3	5.5		5.0	5.0
Protection		3.5	3.9	3.4		4.1	3.5		4.2	2.6	5.4		4.5	4.2
Friend or relative in a gang		2.7	1.3	3.7		2.5	2.7		4.1	2.9	4.9		4.1	4.1
Forced to join		0.8	0.0	1.4		0.7	0.8		1.1	1.1	1.2		1.2	1.1
To get respect		2.1	1.4	2.6		2.2	2.1		3.5	2.0	4.7		3.5	3.5
Money		4.1	6.8	2.5		3.7	4.1		3.8	2.5	4.8		4.1	3.8
To fit in better		0.5	0.0	0.9		0.7	0.5		1.2	1.1	1.4		1.2	1.2
Other reasons		7.1	4.1	9.5		6.4	7.1		5.9	4.5	7.0		5.6	5.9

Table 18. Percentages of Pasco and Florida Statewide youth who reported participation in extracurricular activities, 2016

				Pasco						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
School Sports	30.6	38.2	31.1	39.3	30.0	40.9	34.9	36.0	40.4	34.5	42.5	36.4	41.4	38.5
Organized Sports Outside of School	44.5	24.8	32.4	34.3	42.6	26.1	33.3	41.6	22.4	28.2	32.9	40.1	22.8	30.5
School Band	11.0	5.9	10.0	5.9	11.1	5.2	8.1	16.7	8.1	12.3	11.1	16.2	8.1	11.7
School Club(s)	25.0	27.0	32.4	20.5	24.0	28.2	26.1	22.1	31.8	35.4	20.4	22.8	31.6	27.6
Community Club(s)	10.7	12.2	16.3	7.5	10.1	12.9	11.5	9.3	12.8	14.6	8.2	9.4	12.6	11.3

Table 19. Percentages of <u>high school</u> students who reported *riding in* a vehicle within the past 30 days driven by someone who had been drinking alcohol or using marijuana, or *driving* a vehicle within the past 30 days after drinking alcohol or using marijuana, among Pasco and Florida Statewide youth, 2016

				Pasco						Flori	da State	wide		
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
Riding in a vehicle														
Alcohol		14.8	14.0	15.8		15.2	14.8		16.4	17.5	15.3		16.3	16.4
Marijuana		21.9	22.2	22.5		24.0	21.9		22.7	23.7	21.7		22.9	22.7
Driving a vehicle														
Alcohol		3.6	3.0	4.3		4.0	3.6		5.4	5.0	5.8		5.2	5.4
Marijuana		9.6	8.2	11.4		10.4	9.6		10.3	9.1	11.4		9.9	10.3

Table 20. Trends for <u>high school</u> students *riding in* a vehicle within the past 30 days driven by someone who had been drinking alcohol or using marijuana, or *driving* a vehicle within the past 30 days after drinking alcohol or using marijuana, for Pasco, 2012 and 2016

		2006			2008			2010			2012			2014			2016	
	Middle School	High School	Total	Middle School	High School	Total	Middle School	High School	Total	Middle School	U		Middle School	High School	Total	Middle School	High School	Total
Riding in a vehicle																		
Alcohol											17.1	17.1		15.5	15.5		14.8	14.8
Marijuana											22.9	22.9		23.4	23.4		21.9	21.9
Driving a vehicle																		
Alcohol											7.6	7.6		4.8	4.8		3.6	3.6
Marijuana											10.0	10.0		13.1	13.1		9.6	9.6

Table 21. Protective factor prevalence rates for Pasco, Florida Statewide youth and the national normative database, 2016

Domain	Scale	Pa	sco	Florida S	Statewide	Nationa	l Norms
		Middle School	High School	Middle School	High School	Middle School	High School
Community	Community Rewards for Prosocial Involvement	46	64	46	69	56	63
Family	Family Opportunities for Prosocial Involvement Family Rewards for Prosocial Involvement	59 53	56 52	60 56	59 56	59 54	54 55
School	School Opportunities for Prosocial Involvement School Rewards for Prosocial Involvement	57 48	55 53	53 49	63 59	57 53	60 58
Peer and Individual	Religiosity	44	41	49	57	56	62
Average Prevalence R	ate	51	54	52	61	56	59

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.

Table 22. Risk factor prevalence rates for Pasco, Florida Statewide youth and the national normative database, 2016

Domain	Scale	Pa	sco	Florida S	Statewide	Nationa	l Norms
		Middle School	High School	Middle School	High School	Middle School	High School
Community	Community Disorganization	45	54	42	44	47	47
	Transitions and Mobility	62	63	59	61	47	46
	Laws and Norms Favorable to Drug Use	34	44	37	31	42	42
	Perceived Availability of Drugs	38	33	37	27	45	45
	Perceived Availability of Handguns	26	38	24	36	25	42
Family	Poor Family Management	36	40	40	38	44	45
	Family Conflict	40	39	38	33	42	37
School	Poor Academic Performance	42	49	42	44	45	48
	Lack of Commitment to School	55	62	53	54	47	46
Peer and	Favorable Attitudes toward Antisocial Behavior	37	42	39	35	40	46
Individual	Favorable Attitudes toward ATOD Use	32	42	32	36	39	45
	Early Initiation of Drug Use	20	26	23	22	41	46
Average Preval	lence Rate	39	44	39	38	42	45

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values. The symbol "--" indicates that data are not available.

Table 23. Protective factor prevalence rate trends among middle school students for Pasco, 2006 to 2016

Domain	Scale			Pas	sco		
		2006	2008	2010	2012	2014	2016
Community	Community Rewards for Prosocial Involvement	53	56	53	57	48	46
Family	Family Opportunities for Prosocial Involvement	53	59	55	57	55	59
	Family Rewards for Prosocial Involvement	51	49	48	58	52	53
School	School Opportunities for Prosocial Involvement	53	53	53	56	53	57
	School Rewards for Prosocial Involvement	49	55	51	59	50	48
Peer and Individual	Religiosity	45	41	46	44	41	44
Average Prevalence R	ate	51	52	51	55	50	51

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values. The symbol "--" indicates that data are not available.

Table 24. Risk factor prevalence rate trends among middle school students for Pasco, 2006 to 2016

Domain	Scale			Pa	sco		
		2006	2008	2010	2012	2014	2016
Community	Community Disorganization	49	50	54	51	50	45
	Transitions and Mobility	62	66	62	57	57	62
	Laws and Norms Favorable to Drug Use	43	46	41	36	41	34
	Perceived Availability of Drugs	50	49	54	40	43	38
	Perceived Availability of Handguns	24	24	27	24	28	26
Family	Poor Family Management	50	45	47	39	42	36
	Family Conflict	42	44	40	39	37	40
School	Poor Academic Performance	46	44	44	38	43	42
	Lack of Commitment to School	51	47	52	46	55	55
Peer and	Favorable Attitudes toward Antisocial Behavior	50	46	48	37	40	37
Individual	Favorable Attitudes toward ATOD Use	42	37	42	30	34	32
	Early Initiation of Drug Use	42	36	37	25	24	20
Average Preval	lence Rate	44	41	45	37	41	39

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.

Table 25. Protective factor prevalence rate trends among high school students for Pasco, 2006 to 2016

Domain	Scale	Pasco						
		2006	2008	2010	2012	2014	2016	
Community	Community Rewards for Prosocial Involvement	62	64	60	64	62	64	
Family	Family Opportunities for Prosocial Involvement	54	54	53	58	57	56	
	Family Rewards for Prosocial Involvement	54	51	52	54	54	52	
School	School Opportunities for Prosocial Involvement	57	61	61	62	59	55	
	School Rewards for Prosocial Involvement	56	59	61	67	59	53	
Peer and Individual	Religiosity	53	50	51	53	49	41	
Average Prevalence Rate		56	57	56	60	57	54	

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values. The symbol "--" indicates that data are not available.

Table 26. Risk factor prevalence rate trends among high school students for Pasco, 2006 to 2016

Domain	Scale	Pasco						
		2006	2008	2010	2012	2014	2016	
Community	Community Disorganization	47	55	60	53	56	54	
	Transitions and Mobility	66	65	68	64	61	63	
	Laws and Norms Favorable to Drug Use	40	43	43	35	34	44	
	Perceived Availability of Drugs	47	47	45	33	33	33	
	Perceived Availability of Handguns	42	40	38	36	46	38	
Family	Poor Family Management	50	48	43	36	33	40	
	Family Conflict	38	39	41	38	40	39	
School	Poor Academic Performance	48	44	46	40	43	49	
	Lack of Commitment to School	54	50	52	44	57	62	
Peer and	Favorable Attitudes toward Antisocial Behavior	52	50	46	38	35	42	
Individual	Favorable Attitudes toward ATOD Use	46	42	46	37	39	42	
	Early Initiation of Drug Use	43	37	37	29	26	26	
Average Prevalence Rate		47	45	47	42	42	44	

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.