



# 2008 Florida Youth Substance Abuse Survey



## Columbia County Report



**Executive Office  
of the Governor**

# **2008 Florida Youth Substance Abuse Survey**

**Columbia County Report**



# EXECUTIVE SUMMARY

---

**T**he Florida Legislature's 1999 Drug Control Summit recommended the establishment of an annual, multi-agency-directed, statewide school-based survey effort, combining several survey instruments, with specific variations in odd and even years. The *Florida Youth Substance Abuse Survey (FYSAS)*, one of these instruments and the focus of this report, is administered to a county-level sample of students in even years, and a smaller statewide sample in odd years.

The *FYSAS* is based on the *Communities That Care Youth Survey*, developed from the nationally recognized work of Dr. J. David Hawkins and Dr. Richard F. Catalano. It not only measures the prevalence of alcohol, tobacco and other drug use and delinquent behavior, but also measures the risk and protective factors related to these behaviors.

The 2008 *FYSAS* was administered to 818 Columbia County students in grades 6 through 12 in the spring of 2008. The results supply a valuable source of information to help reduce and prevent the use of alcohol, tobacco and other drugs by school-aged youth.

## Key Survey Results

### Strengths to Build on

- Marijuana use has declined in Columbia County. Overall lifetime marijuana use has gone from 24.0% in 2001 to 19.3% in 2008. Overall past-30-day marijuana use has gone from 13.4% in 2001 to 10.3% in 2008.
- Surveyed students reported a substantial reduction in past-30-day cigarette use. The rate dropped from 18.0% in 2001 to 11.6% in 2008.
- No respondents in middle school reported past-30-day usage of heroin or steroids. No respondents in high school reported lifetime usage of ketamine. No high school past-30-day usage was reported for Rohypnol, ketamine or heroin.
- Among Columbia County middle school students, past-30-day prevalence rates for club drugs (1.6%), LSD, PCP or mushrooms (2.0%), cocaine or crack cocaine (1.1%), methamphetamine (1.1%), depressants (1.6%) and prescription amphetamines (1.2%) are all 2.0% or less. Among Columbia County high school students, past-30-day prevalence rates for Ecstasy (1.7%), hallucinogenic mushrooms (1.7%) and cocaine (1.3%) are all less than 2.0%.
- Among Columbia County high school students, past-30-day prevalence rates for GHB (0.8%), LSD or PCP (0.7%), crack cocaine (0.2%), methamphetamine (0.4%), prescription amphetamines (0.2%) and steroids (0.6%) are all less than 1.0%.
- Relatively few students reported that they would be seen as “cool” by their peers if they drink alcohol regularly (13.6%), smoke cigarettes (5.7%) or smoke marijuana (10.4%).
- A substantial proportion of students indicated that it would be “wrong” or “very wrong” for someone their age to smoke cigarettes (78.3%), smoke marijuana (80.8%) or use other illicit drugs (95.3%).
- A majority of respondents reported that each of the following behaviors poses a “great risk” of harm: smoking a pack or more of cigarettes every day (59.8%) and regular use of marijuana (62.4%).
- High school students reported particularly low rates of risk for two risk factor scales that are directly associated with alcohol, tobacco and other drug use: *Laws and Norms Favorable to Drug Use* (34%) and *Perceived Availability of Drugs* (38%).

- Prevalence rates for *Selling Drugs* (3.7%), *Attempting to Steal a Vehicle* (2.3%) and *Taking a Handgun to School* (0.9%) are all less than 5.0%.

## Opportunities for Improvement

- With overall prevalence rates of 48.9% for lifetime use and 24.9% for past-30-day use, alcohol is the most commonly used drug among Columbia County students.
- After alcohol, students reported cigarettes (30.8% lifetime and 11.6% past-30-day) and marijuana (19.3% lifetime and 10.3% past-30-day) as the most commonly used drugs. Prevalence rates for other drugs are substantially lower.
- Among Columbia County middle school students, 40.0% reported being verbally bullied within the past 30 days.
- Columbia County students reported some of their lowest rates of protection for two reward-based protective factor scales: *School Rewards for Prosocial Involvement* (40%) and *Family Rewards for Prosocial Involvement* (52%). This means that schools and families need to provide additional positive feedback to students, to help them form prosocial bonds in their schools and families.
- Of surveyed Columbia County students, 14.7% reported *Getting Suspended* and 12.6% reported *Being Drunk or High at School*.

These key findings illustrate the complexity of drug use and antisocial behavior among Columbia County's youth and the possible factors that may contribute to these activities. While some of the findings compare favorably to the national findings, Columbia County youth are still reporting drug use and delinquent behavior that will negatively affect their lives and our society.

The *FYSAS* data will enable Columbia County's planners to learn which risk and protective factors to target for their prevention, intervention and treatment programs.

# Table of Contents

<b>METHODOLOGY .....</b>	<b>1</b>
QUESTIONNAIRES .....	1
VALIDITY OF SURVEY DATA .....	2
WEIGHTING .....	2
CONFIDENCE INTERVALS .....	2
DEMOGRAPHICS .....	2
<b>ALCOHOL, TOBACCO AND OTHER DRUG USE .....</b>	<b>3</b>
ALCOHOL .....	4
TOBACCO .....	5
MARIJUANA OR HASHISH .....	6
INHALANTS .....	6
CLUB DRUGS .....	7
<i>Club Drugs in Middle School</i> .....	7
<i>Club Drugs in High School</i> .....	7
OVER-THE-COUNTER DRUGS IN MIDDLE SCHOOL .....	7
PRESCRIPTION DRUGS .....	7
OTHER ILLICIT DRUGS .....	8
<i>Other Illicit Drugs in Middle School</i> .....	8
<i>Other Illicit Drugs in High School</i> .....	8
DRUG COMBINATION RATES .....	9
<i>Any Illicit Drug</i> .....	9
<i>Any Illicit Drug Other than Marijuana</i> .....	9
<i>Alcohol Only</i> .....	9
<i>Alcohol or Any Illicit Drug</i> .....	9
<i>Any Illicit Drug, but No Alcohol</i> .....	10
<b>OTHER ANTISOCIAL BEHAVIORS .....</b>	<b>10</b>
<b>RISK AND PROTECTIVE FACTORS .....</b>	<b>12</b>
THE SOCIAL DEVELOPMENT STRATEGY .....	12
MEASUREMENT .....	14
<i>Calculation of Risk and Protective Factor Thresholds</i> .....	15
<i>Comparing Risk and Protective Factor Prevalence Rates</i> .....	16
<i>Normative Data</i> .....	16
<i>Trend Analysis</i> .....	16
<i>The Middle School Questionnaire</i> .....	16
USING YOUR RISK AND PROTECTIVE FACTOR DATA .....	16
<i>Risk and Protective Factor Prioritization</i> .....	17
<i>Choosing Effective Prevention Strategies</i> .....	20
<b>SPECIAL TOPICS .....</b>	<b>21</b>
EARLY INITIATION OF ATOD USE .....	21
PERCEIVED RISK OF HARM .....	21
PERSONAL DISAPPROVAL .....	22
PEER APPROVAL .....	22
EXTRACURRICULAR ACTIVITIES .....	23



BULLYING BEHAVIOR.....	24
<b>APPENDIX A: DETAILED TABLES .....</b>	<b>27</b>
<b>APPENDIX B: REFERENCES.....</b>	<b>49</b>
<b>APPENDIX C: THE SOCIAL DEVELOPMENT STRATEGY.....</b>	<b>51</b>
<b>APPENDIX D: OTHER RESOURCES.....</b>	<b>53</b>



# 2008 Florida Youth Substance Abuse Survey

## Columbia County Report

**T**he 2008 Florida Youth Substance Abuse Survey (FYSAS) provides scientifically sound information to communities on the prevalence of alcohol, tobacco and other drug (ATOD) use, and risk and protective factors among 6<sup>th</sup> through 12<sup>th</sup> grade students. This information is essential to support effective substance abuse needs-assessment and services planning, and to measure performance outcomes at local and state levels.

This report is one in a series of reports that describes the findings from the FYSAS. As part of the 2008 Florida Youth Survey effort, the FYSAS was administered to Florida youth jointly with the *Florida Youth Tobacco Survey* in May of 2008. The Florida Youth Survey effort was a collaboration among the Florida Departments of Health, Education, Children and Families, Juvenile Justice, and the Florida Office of Drug Control. This report was prepared by Rothenbach Research and Consulting, LLC.

The FYSAS was previously administered at the county level to Columbia County students in the spring of 2001, 2002, 2004 and 2006. While the survey form has been updated over this period, the majority of the instrument has remained unchanged. As a result, the present report includes both an analysis of current survey results and comparisons with the 2001, 2002, 2004 and 2006 survey findings.

This report contains only a brief discussion of methodology. More extensive information on survey administration, methodology and statewide findings can be found in the statewide report, available online at:

[www.dcf.state.fl.us/mentalhealth/publications/fysas/](http://www.dcf.state.fl.us/mentalhealth/publications/fysas/).

### *Methodology*

The sampling strategy was designed to produce survey results that are representative at both the state and county levels, with a minimal margin of error. In Columbia County, this method resulted in a final sample of 424 middle school students and 394 high school students.

### *Questionnaires*

In 2008, for the first time, two versions of the questionnaire were administered to Florida students. High school students received a questionnaire identical to the one used in recent FYSAS efforts. Middle school students received a shortened version of the questionnaire.

While the survey has an excellent track record of yielding high-quality data, concerns have been raised by the FYSAS Workgroup about the ability of some middle school students to complete the questionnaire in a standard classroom period. Analysis of historical data revealed that for some of the items positioned toward the end of the questionnaire, more than 25% of 6<sup>th</sup> grade students fail to provide valid responses.

To address this issue, a shorter version of the standard FYSAS questionnaire—with 176 items compared to 211 on the standard questionnaire—was developed for middle school students. To reach this reduced length, items were removed for eight risk factor scales and four protective factor scales deemed less critical for middle school prevention planning. Also, several ATOD items with very low prevalence among young respondents were either removed or aggregated. Finally, two items that measure the use of over-the-counter drugs in order to get high and eight items that assess bullying behavior were added.

A field test of the new middle school questionnaire, conducted as part of the 2007 FYSAS, yielded



missing value rates that were about 15 percentage points lower than standard questionnaire among 6<sup>th</sup> graders, and about 10 percentage points lower among 7<sup>th</sup> and 8<sup>th</sup> graders.

### ***Validity of Survey Data***

Five strategies were used to assess the validity of survey responses. Data were eliminated from the analysis for students who (1) reported unrealistically high levels of substance use, (2) reported unrealistically high levels of other antisocial behaviors, (3) reported use of a fictitious drug, (4) reported logically inconsistent patterns of substance use, or (5) answered less than 25% of the questions on the survey. These five strategies have been shown to consistently identify most surveys that were completed in a random fashion, those that were not taken seriously, and/or those that were not valid for other reasons.

### ***Weighting***

Before analysis, a set of statistical weights was applied to the 2008 FYSAS dataset. The application of the weights served three purposes. First, weighting compensates for certain elements of the sample design so that the sample selection probability for each student was equal. Second, weighting adjusts for nonresponse at both the school and classroom levels.

Third, weighting adjusts the distribution of the sample across grade levels and gender groups to match the distribution across the full population of Columbia County public school students. Through this process, responses from the grade levels and gender groups that were underrepresented relative to the population are given more weight in the data analysis, while responses from the grade levels and gender groups that were overrepresented are given less weight. The step, called post-stratification, is important because variations in participation across grade levels are common with statewide, school-based survey projects like the FYSAS. Post-stratification makes the sample more representative of the population, and improves the comparability of samples over time.

The 2001, 2002 and 2004 Columbia County datasets were weighted across grade levels but not gender groups.

### ***Confidence Intervals***

For the full sample of Columbia County respondents, the maximum 95% confidence interval estimate (“the margin of error”) is  $\pm 5.5$  percentage points for prevalence rates approximating 50% (such as alcohol

or tobacco). The maximum 95% confidence interval estimate is  $\pm 3.3$  percentage points for prevalence rates of 10% or lower (such as Ecstasy or cocaine). The level of certainty, in this case 95%, means that 95 out of 100 times the “true” population value will fall within the range of the confidence interval. For example, if 40% of the sample indicate using alcohol and the confidence interval is  $\pm 2.0\%$ , then the population value should fall within a range of 38% to 42%.

For subgroup analyses, confidence intervals are larger. Estimates for Columbia County middle school students have confidence intervals ranging from  $\pm 7.5$  percentage points (50% prevalence rates) to  $\pm 4.5$  percentage points (10% prevalence rates). Estimates for high school students have confidence intervals ranging from  $\pm 7.9$  percentage points (50% prevalence rates) to  $\pm 4.8$  percentage points (10% prevalence rates).

Also note that the variance estimates used for these confidence interval calculations include a design effect of 3.0 to adjust for the complex design of the 2008 FYSAS sample.

### ***Demographics***

The survey measures a variety of demographic characteristics. The first two data columns of Table 1 (see Appendix A for data tables) describe the demographic profile of the Columbia County sample before weights were applied. Please note that some categories do not sum to 100% due to missing values.

Despite covering only three out of seven surveyed grades, middle school students constituted slightly more than one half of the sample (51.8% middle school versus 48.2% high school). A higher percentage of the respondents were female (53.1% female versus 44.1% male). White, non-Hispanic students represent 64.9% of the sample. The largest minority population is African American students (17.4%), followed by Hispanic/Latino students (4.4%). The rest of the ethnic breakdown ranges from 0.7% for Native Hawaiian/Pacific Islander students to 9.3% for students who indicated Other/Multiple ethnic backgrounds.

The second set of data columns in Table 1 presents the demographic profile information for the statewide sample.

## Alcohol, Tobacco and Other Drug Use

Alcohol, tobacco and other drug (ATOD) use is measured by a set of 39 items on the 2008 *FYSAS*. While most of the survey items are identical to those used in previous waves of the survey, several key changes have been made.

Starting in 2001, the survey included items measuring: (a) the use of so-called “club drugs” such as Ecstasy, GHB, ketamine and Rohypnol, (b) the use of hallucinogenic mushrooms, and (c) the use of amphetamines, including Ritalin® and Adderall®, without a doctor’s orders. In addition, the use of marijuana and the use of hashish were combined into a single item, and the use of “LSD and other psychedelics” was reworded to read “LSD or PCP.” Also starting in 2001, a parenthetical mentioning the street names “ice” and “crystal meth” was added to the methamphetamine item.

Three changes were made to the ATOD section in 2002: (a) a new item measuring the use of OxyContin® without a doctor’s orders, (b) the prescription drug Xanax® was added to the list of examples given in the “depressants and downers” question, and (c) the “other narcotics” item was replaced by a new question measuring the use of “prescription pain relievers” without a doctor’s

orders.


On the 2006 questionnaire, OxyContin® was removed as an individual item and added to the list of examples included in the prescription pain reliever item. Also, the question for GHB was changed to include a more up-to-date set of slang or street names for the drug.

In 2008, the questionnaire administered to high school students remained unchanged, but the ATOD section of the new middle school questionnaire reduced the number of items by asking broader categories of ATOD use rather than only asking about individual drugs. The new middle school questionnaire also introduces an important new category of ATOD use to the *FYSAS*. A description of these changes is below:

- Items for smokeless tobacco have been removed.
- Items for the club drugs Ecstasy, GHB, ketamine and Rohypnol have been replaced by single items that ask about the use of “club drugs such as Ecstasy, Rohypnol, GHB or ketamine.”
- Items for LSD/PCP and hallucinogenic mushroom use have been combined into a pair of single items that ask about all three drugs.
- Items for cocaine and crack cocaine use have

**Graph 1**

Lifetime and past-30-day use of alcohol, tobacco and other drugs among Columbia County MIDDLE SCHOOL students, 2008



been combined into a pair of single items that ask about both drugs.

- Items that measure the use of over-the-counter drugs in order to get high have been added.

Tables 2 through 5 and Graphs 1 and 2 show the percentage of surveyed Columbia County students who reported using ATODs. These results are presented for both lifetime and past-30-day prevalence of use periods. Lifetime prevalence of use (whether the student has ever used the drug) is a good measure of student experimentation. Past-30-day prevalence of use (whether the student has used the drug within the last month) is a good measure of current use. In addition to the standard lifetime and past-30-day prevalence rates for alcohol use, binge drinking behavior (defined as a report of five or more drinks in a row within the past two weeks) is also measured.

Comparisons to the statewide results of the 2008 survey are presented in Tables 2 through 5 and Graphs 3 through 6. Trend comparisons to Columbia County results from the 2001, 2002, 2004 and 2006 surveys are presented in Tables 6 through 9 and Graphs 3 through 6.

### Alcohol

In most communities, alcohol is the drug used by the largest number of adolescents. As Graphs 1 and 2

show, this is indeed the case in Columbia County.


*Prevalence of Use.* Of the students surveyed in Columbia County in 2008, 48.9% have used alcohol on at least one occasion in their lifetimes. This corresponds to a rate of 40.4% among middle school students and 56.5% among high school students. Current use is substantially lower. Overall, 24.9% of surveyed Columbia County students reported the use of alcohol in the past 30 days, with grade-cohort averages of 21.0% for middle school students and 28.3% for high school students.

*Statewide Comparison.* As Graph 3 shows, the prevalence of past-30-day alcohol use for 2008 is lower in Columbia County compared to the state of Florida as a whole. Overall, 24.9% of surveyed Columbia County students reported the use of alcohol in the past 30 days compared to 29.8% of surveyed students statewide. Grade-cohort analysis shows that this overall lower rate of use is due to the lower rate of usage in high school (28.3% for Columbia County versus 39.5% statewide). Usage in middle school is actually higher in Columbia County than the state as a whole (21.0% for Columbia County versus 17.3% statewide).

*2001-2008 Trend.* In Columbia County, between 2001 and 2008, overall past-30-day alcohol use decreased 8.1 percentage points. Among middle school students, use increased 1.5 percentage points,


**Graph  
2**

Lifetime and past-30-day use of alcohol, tobacco and other drugs among Columbia County HIGH SCHOOL students, 2008



**Graph 3**

Past-30-day alcohol use, Columbia County 2001-2008 and Florida 2008



and among high school students, use decreased 17.9 percentage points. Between 2006 and 2008, the two most recent waves of the Columbia County survey, overall past-30-day alcohol use decreased 4.0 percentage points. Among middle school students, use increased 4.0 percentage points, and among high school students, use decreased 9.8 percentage points.

**Binge Drinking.** Findings on binge drinking (defined as consuming five or more drinks in a row within the past two weeks) are likely to be among the most important findings related to alcohol use (Johnston, O'Malley, Bachman & Schulenberg, 2008). In Columbia County, 16.2% of surveyed students reported binge drinking, with corresponding rates of 11.4% among middle school students and 20.5% among high school students. While this represents a similar rate of high school binge drinking compared to the state as a whole (21.5%), Columbia County middle school students reported a higher rate compared to results from across Florida (6.2%).

### Tobacco

This section of the report discusses the prevalence of tobacco use as measured by the 2008 FYSAS. Another survey, the 2008 Florida Youth Tobacco Survey (Florida Department of Health), was administered simultaneously with the 2008 FYSAS, and was specifically tobacco related. That survey is Florida's official source for youth tobacco use information. The information presented in this report is consistent with findings reported in the 2008 Florida Youth Tobacco Survey.

**Prevalence of Use.** Of the students surveyed in Columbia County in 2008,

30.8% have used cigarettes on at least one occasion in their lifetimes. This corresponds to a rate of 26.2% among middle school students and 35.0% among high school students. Current use is substantially lower. Overall, 11.6% of surveyed Columbia County students reported the use of cigarettes in the past 30 days, with grade-cohort averages of 8.8% for middle school students and 14.0% for high school students.

**Statewide Comparison.** As Graph 4 shows, the prevalence of past-30-day cigarette use for 2008 is higher in Columbia County compared to the state of Florida as a whole. Overall, 11.6% of


surveyed Columbia County students reported the use of cigarettes in the past 30 days compared to 9.1% of surveyed students statewide. This increased rate of use applies both to middle school (8.8% for Columbia County versus 4.7% statewide) and high school (14.0% for Columbia County versus 12.6% statewide) grade-cohorts.

**2001-2008 Trend.** In Columbia County, between 2001 and 2008, overall past-30-day cigarette use decreased 6.4 percentage points. Among middle school students, use decreased 1.2 percentage points, and among high school students, use decreased 11.7 percentage points. Between 2006 and 2008, the two most recent waves of the Columbia County survey, overall past-30-day cigarette use increased 1.0 percentage points. Among middle school students, use increased 0.3 percentage points, and among high school students, use increased 3.3 percentage points.

**Smokeless Tobacco.** The prevalence of current use of smokeless tobacco among high school students is


**Graph 4**

Past-30-day cigarette use, Columbia County 2001-2008 and Florida 2008



**Graph 5**

Past-30-day marijuana or hashish use, Columbia County 2001-2008 and Florida 2008



similar to the rate of cigarette use in Columbia County. Overall, 20.4% of surveyed Columbia County high school students reported using smokeless tobacco in their lifetimes and 9.2% reported using it within the past 30 days.

### Marijuana or Hashish

During the 1990s, there were major changes in trends of marijuana use throughout the United States. Results from the *Monitoring the Future* study show dramatic increases in both lifetime and past-30-day prevalence rates through the early and mid 1990s (Johnston et al., 2008). For 8<sup>th</sup> and 10<sup>th</sup> graders the past-30-day rates more than doubled during this period. Since 1996 and 1997, when marijuana use peaked, rates have declined.

**Prevalence of Use.** Of the students surveyed in Columbia County in 2008, 19.3% have used marijuana or hashish on at least one occasion in their lifetimes. This corresponds to a rate of 14.0% among middle school students and 23.9% among high school students. Current use is substantially lower. Overall, 10.3% of surveyed Columbia County students reported the use of marijuana or hashish in the past 30 days, with grade-cohort averages of 7.9% for middle school students and 12.4% for high school students.

**Statewide Comparison.** As Graph 5 shows, the prevalence of past-30-day marijuana or hashish use for 2008 in Columbia County is similar to the rate for the state of Florida as a whole. Overall, 10.3% of surveyed Columbia County students reported the use of marijuana or hashish in the past 30 days compared to 11.1% of surveyed students

statewide. Despite this overall similarity, there were larger usage rate differences in both middle school and high school, although in opposite directions (7.9% for Columbia County versus 4.4% statewide in middle school and 12.4% for Columbia County versus 16.2% statewide in high school).

**2001-2008 Trend.** In Columbia County, between 2001 and 2008, overall past-30-day marijuana use decreased 3.1 percentage points. Among middle school students, use increased 0.4 percentage points, and among high school students, use decreased 6.6 percentage points.

Between 2006 and 2008, the two most recent waves of the Columbia County survey, overall past-30-day marijuana use increased 2.7 percentage points. Among middle school students, use increased 0.4 percentage points, and among high school students, use increased 4.4 percentage points.


### Inhalants

After alcohol, tobacco and marijuana, the most commonly used drug among Florida students is inhalants. Inhalant use is measured by the survey question, "On how many occasions (if any) have you used inhalants (whippets, butane, paint thinner, or glue to sniff, etc.)?" Inhalant use is more prevalent with younger students, perhaps because it is often the easiest drug for them to obtain. The negative consequences of inhalant use can be substantial; one of them being that it is associated with the use of other illicit drugs later in life.

**Prevalence of Use.** Of the students surveyed in Columbia County in 2008, 13.3% have used inhalants on at least one occasion in their lifetimes. This

**Graph 6**

Past-30-day inhalant use, Columbia County 2001-2008 and Florida 2008





corresponds to a rate of 13.4% among middle school students and 13.3% among high school students. Current use is substantially lower. Overall, 3.4% of surveyed Columbia County students reported the use of inhalants in the past 30 days, with grade-cohort averages of 4.2% for middle school students and 2.7% for high school students.

**Statewide Comparison.** As Graph 6 shows, the prevalence of past-30-day inhalant use for 2008 in Columbia County is similar to the rate for the state of Florida as a whole. Across all surveyed grades, 3.4% of surveyed Columbia County students reported the use of inhalants in the past 30 days compared to 3.5% of surveyed students statewide. Despite this overall similarity, there was a larger usage rate difference in middle school (4.2% for Columbia County versus 5.2% statewide). The usage rate in high school was similar to the statewide rate (2.7% for Columbia County versus 2.2% statewide).

**2001-2008 Trend.** In Columbia County, between 2001 and 2008, overall past-30-day inhalant use decreased 1.2 percentage points. Among middle school students, use decreased 0.5 percentage points, and among high school students, use decreased 1.8 percentage points. Between 2006 and 2008, the two most recent waves of the Columbia County survey, overall past-30-day inhalant use increased 0.1 percentage points. Among middle school students, use increased 1.2 percentage points, and among high school students, use increased 0.8 percentage points.

### **Club Drugs**

Club drugs are a broad category of illicit substances that are classified together because their use started at dance clubs and “raves,” not because they are of a similar chemical class (like amphetamines). Their use, however, has expanded beyond these settings. For the purpose of the 2008 FYSAS, club drugs include Ecstasy, GHB, ketamine and Rohypnol. Note that this list is not meant to be exclusive, as other drugs are used at clubs and raves.

### **Club Drugs in Middle School**

**Prevalence of Use.** Of the middle school students surveyed in Columbia County in 2008, 2.1% have used club drugs on at least one occasion in their lifetimes and 1.6% have used club drugs in the past 30 days.

**Statewide Comparison.** The prevalence of past-30-day club drug use for 2008 is higher in Columbia County compared to the state of Florida as a whole. In middle school, 1.6% of surveyed Columbia

County students reported the use of club drugs in the past 30 days compared to 0.6% of surveyed students statewide.

### **Club Drugs in High School**

**Prevalence of Use.** Columbia County high school students reported lifetime prevalence-of-use rates of 4.4% for Ecstasy, 0.8% for Rohypnol, 1.3% for GHB, and 0.0% for ketamine. The prevalence of use within the past 30 days is lower, with all four drugs below 2.0%.

**Statewide Comparison.** In high school, lifetime prevalence rates for club drug use in Columbia County are similar to those found for the state of Florida as a whole. The two largest differences were for ketamine use (0.0% in Columbia County versus 0.8% in Florida) and GHB use (1.3% in Columbia County versus 0.7% in Florida). Past-30-day prevalence rates are too low to allow a meaningful comparison between the samples.

### **Over-The-Counter Drugs in Middle School**

The use of over-the-counter (OTC) drugs was measured by asking: “On how many occasions (if any) have you used drugs that can be purchased from a store without a prescription—such as cold and cough medication—in order to get high in your lifetime?” and “... in the past 30 days?”

**Prevalence of Use.** Of the middle school students surveyed in Columbia County in 2008, 5.7% have used OTC drugs on at least one occasion in their lifetimes and 3.1% have used OTC drugs in the past 30 days.

**Statewide Comparison.** The prevalence of past-30-day OTC drug use for 2008 in Columbia County is similar to the rate for the state of Florida as a whole. In middle school, 3.1% of surveyed Columbia County students reported the use of OTC drugs in the past 30 days compared to 2.2% of surveyed students statewide.

### **Prescription Drugs**

While students across the country have reported declining rates of use for many illicit drugs over the past 10 years, prescription drugs have largely bucked this trend. As a result, prevalence rates for using prescription drugs without a doctor’s orders are higher than for many illicit drugs (Johnston et al., 2008). The 2008 FYSAS includes questions that assess the use of prescription pain relievers, depressants and amphetamines. Results for these

prescription drugs are presented in Tables 3, 5, 7 and 9.

**Prevalence of Use.** Columbia County middle school and high school students reported lifetime prevalence-of-use rates for this group of drugs that range from a high of 8.9% for prescription pain relievers and 4.9% for depressants to a low of just 1.6% for amphetamines. The prevalence of use within the past 30 days is lower, with highs of 3.8% for prescription pain relievers and 2.2% for depressants. The remaining illicit drugs have past-30-day prevalence rates of less than 2.0%.

**Statewide Comparison.** Lifetime prevalence rates for prescription drug use in Columbia County are similar to those found for the state of Florida as a whole. The two largest differences were for amphetamine use (1.6% in Columbia County versus 3.7% in Florida) and depressant use (4.9% in Columbia County versus 6.0% in Florida). Past-30-day prevalence rates are too low to allow a meaningful comparison between the samples.

#### Other Illicit Drugs

The 2008 FYSAS also measured the prevalence of use of a variety of other illicit drugs among Columbia County students. This includes the use of the following: LSD or PCP, hallucinogenic mushrooms, cocaine, crack cocaine, methamphetamine, heroin and steroids. Results for these illicit drugs are

presented in Tables 3, 5, 7 and 9.

#### Other Illicit Drugs in Middle School

**Prevalence of Use.** As is typical of adolescent populations, the prevalence-of-use rates in Columbia County for these other illicit drugs are much lower than the rates for alcohol, tobacco and marijuana. Among middle school students, lifetime prevalence-of-use rates for this group of drugs range from a high of 1.9% for LSD, PCP or mushrooms to a low of 1.0% for heroin. The prevalence of use within the past 30 days goes from a high of 2.0% for LSD, PCP or mushrooms to a low of 0.0% for steroids and heroin.


**Statewide Comparison.** In middle school, lifetime prevalence rates for other illicit drug use in Columbia County are similar to those found for the state of Florida as a whole. The two largest differences were for LSD, PCP or mushrooms use (1.9% in Columbia County versus 1.5% in Florida) and steroid use (1.2% in Columbia County versus 0.8% in Florida). Past-30-day prevalence rates are too low to allow a meaningful comparison between the samples.

#### Other Illicit Drugs in High School

**Prevalence of Use.** As is typical of adolescent populations, the prevalence-of-use rates in Columbia County for these other illicit drugs are much lower than the rates for alcohol, tobacco and marijuana.

**Graph 7**

Past-30-day drug combination rates for Columbia County and Florida Statewide, 2008



Among high school students, lifetime prevalence-of-use rates for this group of drugs range from a high of 6.1% for mushrooms to a low of 0.5% for heroin. The prevalence of use within the past 30 days is lower, going from a high of 1.7% for mushrooms to a low of 0.0% for heroin.

*Statewide Comparison.* In high school, lifetime prevalence rates for other illicit drug use in Columbia County are similar to those found for the state of Florida as a whole. The two largest differences were for cocaine use (3.9% in Columbia County versus 5.5% in Florida) and LSD or PCP use (2.1% in Columbia County versus 3.0% in Florida). Past-30-day prevalence rates are too low to allow a meaningful comparison between the samples.

#### ***Drug Combination Rates***

Prevalence-of-use rates for combinations of drugs provide a helpful summary of drug use behavior. Tables 2, 4, 6 and 8 and Graph 7 provide lifetime and past-30-day prevalence rates for the use of one or more drugs from a set of illicit drugs. Illicit drugs are substances that are illegal for adults to use, so they include all drugs on the survey except alcohol, cigarettes and smokeless tobacco. Five types of drug combination rates are presented here:

**Any illicit drug** – Use of at least one illicit drug

**Any illicit drug other than marijuana** – Use of at least one illicit drug other than marijuana

**Alcohol only** – The use of alcohol and no illicit drugs

**Alcohol or any illicit drug** – Use of alcohol or at least one illicit drug

**Any illicit drug but no alcohol** – Use of at least one illicit drug, without any use of alcohol

These combination categories are created using all the illicit drug items on the current high school questionnaire: marijuana or hashish, inhalants, Ecstasy, Rohypnol, GHB, ketamine, LSD or PCP, hallucinogenic mushrooms, methamphetamine, cocaine, crack cocaine, depressants, heroin, prescription pain relievers, amphetamines and steroids. Please note that the combination categories for middle school respondents include the same illicit drugs, but as described at the beginning of this section, a reduced set of items is used to ask about these drugs.

Trend comparisons for these drug combination rates begin in 2002. This is because a number of the illicit

drugs were not included on the 2000 questionnaire. Also, OxyContin® was combined with prescription pain relievers in 2006, but this change is minor and has almost no impact on the drug combination trend lines.

#### **Any Illicit Drug**

Overall, 28.0% of surveyed Columbia County students reported at least one use of *any illicit drug* in their lifetimes, and 15.1% reported use in the past 30 days. The past-30-day prevalence rate corresponds to 14.2% among middle school students and 15.9% among high school students. As Graph 7 shows, use of *any illicit drug* in the past 30 days is similar in Columbia County and the state (15.1% for Columbia County versus 15.8% statewide).

#### **Any Illicit Drug Other than Marijuana**

The purpose of this drug combination rate is to provide prevention planners with an overall indicator of so-called “hard” drug use (Johnston et al., 2008). Overall, 21.0% of surveyed Columbia County students reported at least one use of *any illicit drug other than marijuana* in their lifetimes, and 9.2% reported use in the past 30 days. The past-30-day prevalence rate corresponds to 9.7% among middle school students and 8.8% among high school students. As Graph 7 shows, use of *any illicit drug other than marijuana* in the past 30 days is similar in Columbia County and the state (9.2% for Columbia County versus 8.9% statewide).

It is important to note that this measure—the current use of all illicit drugs other than marijuana *combined*—is less than the past-30-day prevalence of use of alcohol (24.9%), marijuana (10.3%) and cigarettes (11.6%), as well as the prevalence of binge drinking (16.2%).

#### **Alcohol Only**

Overall, 24.7% of surveyed Columbia County students reported at least one use of *alcohol only*—the use of alcohol and no illicit drugs—in their lifetimes, and 15.3% reported use in the past 30 days. The past-30-day prevalence rate corresponds to 12.9% among middle school students and 17.4% among high school students. As Graph 7 shows, use of *alcohol only* in the past 30 days is lower in Columbia County than across the state of Florida as a whole (15.3% for Columbia County versus 18.5% statewide).

#### **Alcohol or Any Illicit Drug**

*Alcohol or any illicit drug* use is a summary measure that included all drugs from the 2008 survey, with the



exception of cigarettes and smokeless tobacco. Overall, 52.4% of surveyed Columbia County students reported at least one use of *alcohol or any illicit drug* in their lifetimes, and 29.6% reported use in the past 30 days. The past-30-day prevalence rate corresponds to 26.5% among middle school students and 32.2% among high school students. As Graph 7 shows, use of *alcohol or any illicit drug* in the past 30 days is lower in Columbia County than across the state of Florida as a whole (29.6% for Columbia County versus 33.9% statewide).

### Any Illicit Drug, but No Alcohol

The final drug combination category measures the use of illicit drugs by students who are not using alcohol. As Tables 2 and 4 show, this combination is quite rare. Overall, 3.7% of surveyed Columbia County students reported having used illicit drugs in their lifetimes but never using alcohol. Current use of illicit drugs (within the past 30 days) without the accompanying use of alcohol is also rare (5.0%). The past-30-day prevalence rate corresponds to 5.8% among middle school students and 4.3% among high school students. As Graph 7 shows, use of *any illicit drug, but no alcohol* in the past 30 days is similar in Columbia County and the state (5.0% for Columbia County versus 4.4% statewide).

## Other Antisocial Behaviors


The 2008 FYSAS also measures a series of eight other problem or antisocial behaviors—that is, behaviors that run counter to established norms of good behavior. Note that information on antisocial behaviors is collected only for a prevalence period of the past 12 months. The survey measured the following antisocial behaviors: *Carrying a Handgun*, *Selling Drugs*, *Attempting to Steal a Vehicle*, *Being Arrested*, *Taking a Handgun to School*, *Getting Suspended*, *Attacking Someone with Intent to Harm* and *Being Drunk or High at School*.

Prevalence rates for these behaviors among Columbia County students, as well as comparison rates from the statewide survey, are presented in Table 10 and Graph 8. Trend comparisons to Columbia County results from the 2001, 2002, 2004 and 2006 surveys are presented in Table 16.

As Table 10 shows, the prevalence rates reported by Columbia County students differ substantially across the eight antisocial behaviors measured in the survey. Reports of *Taking a Handgun to School* (0.9%), *Attempting to Steal a Vehicle* (2.3%), and *Selling Drugs* (3.7%) are rare, while *Getting Suspended* (14.7%), *Being Drunk or High at School* (12.6%), and *Attacking Someone with Intent to Harm* (10.6%) are more common.

**Graph 8**

Comparisons of past-12-month delinquent behavior for Columbia County and Florida Statewide, 2008



.....

*Carrying a Handgun.* In Columbia County, 6.0% of students reported carrying a handgun in the past year, with rates of 6.1% and 6.0% for middle school and high school students, respectively. Male students (10.4%) were more likely than female students (1.5%) to have reported this behavior. Across the state as a whole, 5.0% of students reported carrying a handgun.

*Selling Drugs.* In Columbia County, 3.7% of students reported selling drugs in the past year, with rates of 2.8% and 4.6% for middle school and high school students, respectively. Male students (5.8%) were more likely than female students (1.8%) to have reported this behavior. Across the state as a whole, 5.5% of students reported selling drugs.

*Attempting to Steal a Vehicle.* In Columbia County, 2.3% of students reported attempting to steal a vehicle in the past year, with rates of 2.8% and 1.7% for middle school and high school students, respectively. Male students (3.0%) were more likely than female students (1.4%) to have reported this behavior. Across the state as a whole, 2.5% of students reported attempting to steal a vehicle.

*Being Arrested.* In Columbia County, 5.2% of students reported being arrested in the past year, with rates of 5.4% and 4.9% for middle school and high school students, respectively. Male students (7.3%) were more likely than female students (3.1%) to have reported this behavior. Across the state as a whole, 4.9% of students reported being arrested.

*Taking a Handgun to School.* In Columbia County, 0.9% of students reported taking a handgun to school in the past year, with rates of 1.2% and 0.7% for middle school and high school students, respectively. Male students (1.9%) were more likely than female students (0.2%) to have reported this behavior. Across the state as a whole, 1.0% of students reported taking a handgun to school.

*Getting Suspended.* In Columbia County, 14.7% of students reported getting suspended in the past year, with rates of 15.3% and 14.2% for middle school and high school students, respectively. Male students (18.1%) were more likely than female students (10.9%) to have reported this behavior. Across the state as a whole, 15.2% of students reported getting suspended.

Note, however, that the questionnaire item used to measure *Getting Suspended* does not define “suspension.” Rather, it is left to the individual respondent to define. Because suspension policies


vary substantially from county to county, comparisons to statewide results should be interpreted with caution for this item.

*Attacking Someone with Intent to Harm.* In Columbia County, 10.6% of students reported attacking someone with intent to harm in the past year, with rates of 11.1% and 10.3% for middle school and high school students, respectively. Male students (15.9%) were more likely than female students (5.6%) to have reported this behavior. Across the state as a whole, 11.8% of students reported attacking someone with intent to harm.

*Being Drunk or High at School.* In Columbia County, 12.6% of students reported being drunk or high at school in the past year, with rates of 9.0% and 15.8% for middle school and high school students, respectively. Male students (14.8%) were more likely than female students (10.7%) to have reported this behavior. Across the state as a whole, 11.6% of students reported being drunk or high at school.

**Graph  
9**

## MIDDLE SCHOOL protective factor prevalence rates for Columbia County, 2008



### ***Risk and Protective Factors***

Just as smoking is a risk factor for heart disease and getting regular exercise is a protective factor against heart disease and other health problems, there are factors that can help protect youth from, or put them at risk for, drug use and other problem behaviors.

**Protective factors**, also known as “assets,” are conditions that buffer children and youth from exposure to risk by either reducing the impact of the risks or changing the way that young people respond to risks.

**Risk factors** are conditions that increase the likelihood of a young person becoming involved in drug use, delinquency, school dropout and/or violence. For example, children living in families with poor family supervision are more likely to become involved in these problems.

Research during the past 30 years supports the view that delinquency; alcohol, tobacco and other drug use; school achievement; and other important outcomes in adolescence are associated with specific risk and protective factors in the student’s community, school and family environments, as well as with characteristics of the individual (Hawkins,

Catalano & Miller, 1992). In fact, these risk and protective factors have been shown to be more important in understanding these behaviors than ethnicity, income or family structure (Blum et al., 2000).


There is a substantial amount of research showing that adolescents’ exposure to a greater number of risk factors is associated with more drug use and delinquency. There is also evidence that exposure to a number of protective factors is associated with lower prevalence of these problem behaviors (Bry, McKeon & Pandina, 1982; Newcomb, Maddahian & Skager, 1987; Newcomb & Felix-Ortiz, 1992; Newcomb, 1995; Pollard et al., 1999).

### ***The Social Development Strategy***

The Social Development Strategy (Hawkins, Catalano & Associates, 1992) organizes these risk and protective factors into a framework that families, schools and communities can use to help children develop healthy behaviors. This strategy, which is graphically depicted in Appendix C, shows how three broad categories of protective factors—healthy beliefs and clear standards, bonding, and individual characteristics—work together to promote positive youth development and healthy behaviors (Hawkins, Arthur & Catalano, 1995). The Social Development

**Graph  
10**

## MIDDLE SCHOOL risk factor prevalence rates for Columbia County, 2008



Strategy begins with a goal of healthy behaviors for all children and youth. In order for young people to develop healthy behaviors, adults must communicate healthy beliefs and clear standards for behavior to young people (Catalano & Hawkins, 1996). Bonding (an attached, committed relationship) between a child and an adult who communicates healthy beliefs and clear standards motivates the child to follow healthy beliefs and clear standards. A child who forges a bond with an adult is less likely to threaten the relationship by violating the beliefs and standards held by the adult. Research has identified three conditions for bonding (Catalano & Hawkins, 1996):


- First, children need developmentally appropriate opportunities for meaningful involvement with a positive social group (community, family, school, etc.) or individual.

- Second, children need the emotional, cognitive, social and behavioral skills to successfully take advantage of opportunities.
- Third, children must be recognized for their involvement. Recognition sets up a reinforcing cycle in which children continue to look for opportunities and learn skills and, therefore, receive recognition.

Certain characteristics with which some children come into the world (positive social orientation, resilient temperament and high intelligence) can also help protect children from risk. For children who do not have the protective advantages of these characteristics, in order to build strong bonds to family, school and community, it is even more important for community members to:

**Graph  
11**

## HIGH SCHOOL protective factor prevalence rates for Columbia County, 2008



- make extra efforts to provide opportunities for involvement
- teach the social, emotional, and cognitive skills needed to be successful
- recognize children's efforts as well as their successes.

The developmental process outlined in this model has important implications for prevention planning. Programs that seek to change the attitudes young people hold about the pros and cons of ATOD use, for example, may produce an immediate reduction in the prevalence of problem behaviors. The effectiveness of these efforts will be limited, however, by the risk and protective factors that underlie the acquisition of healthy beliefs and clear standards. If young people have weak bonds to prosocial groups and strong bonds to antisocial groups, they will be less receptive to drug abuse prevention messages.

An alternative prevention strategy might involve targeting the risk and protective factors that operate at an earlier point in the developmental process. While programs and policies that increase the opportunities for prosocial involvement in the family,

at school and in the community may not yield an immediate reduction in the rates of ATOD use, they will encourage young people to form attachments to sources of positive social influence, thereby building the foundation for healthy behavioral choices in the future.

### Measurement

The *FYSAS* assesses 23 risk factors and 10 protective factors across four domains: Community Domain, Family Domain, School Domain, and Peer and Individual Domain. Each factor is measured by a set of survey items called a scale.

The 2008 *FYSAS* uses the same risk and protective factors scales employed in previous survey efforts. In other words, the same survey items are still used to construct each scale. (Please note that the middle school survey employs a reduced set of risk and protective factor scales. The difference between the middle school and high school questionnaires is described below.)

This year, a new method is being used to convert these scales into scores. This change is a response to requests for a risk and protective factor scoring system that is more intuitive, and therefore easier to incorporate into the prevention planning process.

For each risk and protective factor scale, the new scoring method sets a threshold above which respondents are considered to have a high level of risk or protection and below which they are considered to have a low level of risk or protection. It then becomes possible to count the number of students with high levels of risk or protection on each scale. This approach, in turn, allows risk and protective factor data to be reported in the same way as ATOD data: as prevalence rates.

Under this new system, a score of 60 for the protective factor *School Rewards for Prosocial Involvement* would indicate that 60% of surveyed students reported a high level of protection for this protective factor, while 40% reported a low level of protection. Risk factor scales are scored in the same way. For example, a score of 55 for the risk factor *Friends' Use of Drugs* would indicate that 55% of surveyed students reported a high level of risk for this

risk factor, while 45% reported a low level of risk.


Risk and protective factor scale prevalence rates for the overall sample of Columbia County middle school and high school students are presented in Tables 18 and 19 and Graphs 9 to 12.

### Calculation of Risk and Protective Factor Thresholds

The high-risk and high-protection thresholds used to calculate the risk and protective factor prevalence rates were calculated using a method recommended by Arthur et al. (2007). For risk factor scales, the high-risk threshold is the normative median—that is the scale's median value in the *Communities That Care* normative database—plus .15 times the mean absolute deviation (a measure of central tendency similar to the standard deviation). In other words, risk factor thresholds are set slightly above the normative median. For protective factor scales, the high-

**Graph  
12**

**HIGH SCHOOL risk factor prevalence rates for Columbia County, 2008**





protection threshold is the normative median minus .15 times the mean absolute deviation. In other words, protective factor thresholds are set slightly below the normative median.

It is also important to note that risk and protection thresholds are calculated separately for each grade level. For most risk factors, this means that older students must report a higher level of risk before crossing the scoring threshold and being designated as at risk. For most protective factors, this means that older students must report a lower level of protection before crossing the scoring threshold and being designated as protected.

### Comparing Risk and Protective Factor Prevalence Rates

The simplicity of the new prevalence rate scoring method will make it easier for prevention planners to analyze and compare risk and protective factor scores. However, comparisons to national risk and protective factor norms from the *Communities That Care* normative database must now be done differently.

Under the old percentile scoring system, the national median score was 50 for all risk and protective factor scales. Scores above 50 were, by definition, higher than the national median and scores below 50 were lower than the national median. Under the new method, median scores from the *Communities That Care* normative database differ for each risk and protective factor scale. These new national risk and protective factor norms are presented in Tables 18 and 19.

The risk factor scale *Early Initiation of Drug Use* provides an example. As shown in Table 52, 36% of surveyed Florida middle school and high school students reported scale scores above the high-risk threshold. In other words, 36% of surveyed Florida students are at risk due to early experimentation with drugs. Table 54 shows that across the national *Communities That Care* normative sample, 43% of survey students are at risk due to early experimentation with drugs. Florida's score of 36% is seven percentage points below the normative score.

### Normative Data

The *Communities That Care* normative database contains survey responses from over 280,000 students in grades 6 through 12. It was compiled by combining the results of selected *Communities That Care Youth Survey* efforts that were completed in 2000, 2001 and 2002. To enhance representativeness,

statistical weights were applied to adjust the sample to exactly match the population of U.S. public school students on four key demographic variables: ethnicity, sex, socioeconomic status and urbanicity. Information on the U.S. public school student population was obtained from the Common Core of Data program at the U.S. Department of Education's National Center for Education Statistics.

### Trend Analysis

Risk and protective factor scale scores generated with the new prevalence rate scoring system are not directly comparable to scores generated with the previous percentile scoring system. As a result, scores from the 2000 to 2008 FYSAS have been recalculated using the new methodology in order to support trend analysis. These results are presented in Tables 20 through 23.

### The Middle School Questionnaire

As previously noted, middle school students were given a shorter version of the FYSAS questionnaire. The following 12 risk and protective factor scales, which were deemed less critical for middle school prevention planning, are not included in the middle school survey:

- *Community Opportunities for Prosocial Involvement*
- *Family Attachment*
- *Social Skills*
- *Belief in the Moral Order*
- *Low Neighborhood Attachment*
- *Laws and Norms Favorable to Handguns*
- *Family History of Antisocial Behavior*
- *Parental Attitudes Favorable toward Antisocial Behavior*
- *Rebelliousness*
- *Friends' Delinquent Behavior*
- *Friends' Use of Drugs*
- *Sensation Seeking*

For these risk and protective factor scales, results are only presented for high school students.

### Using Your Risk and Protective Factor Data

The analysis of risk and protective factors is the most powerful tool available for understanding what promotes both positive and negative adolescent

behavior and for helping design successful prevention programs for young people. To promote positive development and prevent problem behavior, it is necessary to address the factors that predict these outcomes. By measuring these risk and protective factors, specific factors that are elevated can be prioritized in the community. This process also helps in selecting tested-effective prevention programming shown to address those elevated factors and consequently provide the greatest likelihood for success.

### **Risk and Protective Factor Prioritization**

In general, a prevention strategy that focuses on a relatively narrow set of developmental factors can be more effective than a strategy that spreads resources across a broad set of factors. Risk and protective factor data from the *FYSAS* can provide critical guidance in this prioritization process. That is, prevention planners can use the information gathered by the survey to identify youth development areas where programs, policies and practices are likely to have the greatest positive impact.

Start the prioritization process by identifying the protective factor scales with the lowest percentage of protected students and the risk factor scales with the highest percentage of at risk students. It may also be helpful to identify scales with particularly high percentages of protected students or low percentages of at risk students. These areas represent strengths that prevention planners in Columbia County may wish to build on. In addition, it is also important to compare the rates of risk and protection reported by Columbia County students to the rates reported by students in the national normative sample.

### **Lowest Protective Factor Scales**

- Across all 10 protective factor scales, middle school students in Columbia County reported the lowest level of protection for the *School Rewards for Prosocial Involvement* scale. Their score of 40% was five points lower than the statewide average of 45%. In the national normative sample, 53% reported an elevated level of protection, 13 points higher than Columbia County. Low scores on this scale indicate that students receive less praise and encouragement when they work hard and do well in school. This lack of positive feedback, in turn, may weaken the bonds students form with teachers, coaches and prosocial peers.
- High school students in Columbia County reported the lowest level of protection for the

*Community Opportunities for Prosocial Involvement* scale. Their score of 47% was eight points higher than the statewide average of 39%. In the national normative sample, 52% reported an elevated level of protection, five points higher than Columbia County. Students who reported low scores on this scale have fewer opportunities to interact closely with positive adult role models in their neighborhoods and to participate in sports, clubs and other prosocial community activities. As a result, these students are less likely to form strong community bonds that encourage the adoption of prosocial norms and values.

- Additionally, middle school students in Columbia County also reported a low level of protection for the *School Opportunities for Prosocial Involvement* scale. Their score of 48% was one point higher than the statewide average of 47%. In the national normative sample, 57% reported an elevated level of protection, nine points higher than Columbia County. Students with low scores on this scale have fewer opportunities to interact closely with teachers, get involved with special projects and activities in the classroom, and participate in sports, clubs and other school activities outside of the classroom. This lack of involvement deprives students of the opportunity to form healthy relationships with teachers and prosocial peers.
- High school students also reported low levels of protection for two other scales. The first of these was *Family Opportunities for Prosocial Involvement*. Their score of 54% was one point higher than the statewide average of 53%. In the national normative sample, 54% reported an elevated level of protection, the same as Columbia County. Low scores on this scale indicate that activities that promote family attachment—such as family recreation and involvement in family decisions—are less available to students. These prosocial activities reinforce family bonds and cause students to more easily adopt the norms projected by their families. For instance, children whose parents have high expectations for their school achievement are less likely to drop out of school.
- The second additional protective factor scale with a low score in high school was *Family Attachment*. Their score of 54% was one point lower than the statewide average of 55%. In the national normative sample, 56% reported an



elevated level of protection, two points higher than Columbia County. Students who reported low scores on this scale feel less of a bond with their parents than students with high scores. Positive bonding can act as a buffer against risk factors. A weaker bond means that children are less likely to accept guidance from parents that discourages antisocial behavior.

### **Highest Risk Factor Scales**

#### Community Domain:

- Within the Community Domain, both middle school and high school students in Columbia County reported the highest level of risk for the *Transitions and Mobility* scale. Among middle school students, 60% reported an elevated level of risk, one point lower than the statewide average of 61%. In the national normative sample, 47% reported an elevated level of risk, 13 points lower than Columbia County. Among high school students, 60% reported an elevated level of risk, four points lower than the statewide average of 64%. In the national normative sample, 46% reported an elevated level of risk, 14 points lower than Columbia County. High scores on this scale indicate that students are changing homes and schools more frequently. Dislocations of this type can inhibit the ability of young people to become involved with prosocial organizations and individuals within their school and community.

#### Family Domain:

- Within the Family Domain, both middle school and high school students in Columbia County reported the highest level of risk for the *Poor Family Management* scale. Among middle school students, 55% reported an elevated level of risk, six points higher than the statewide average of 49%. In the national normative sample, 44% reported an elevated level of risk, 11 points lower than Columbia County. Among high school students, 48% reported an elevated level of risk, one point lower than the statewide average of 49%. In the national normative sample, 45% reported an elevated level of risk, three points lower than Columbia County. Students with high scores on this scale live in families in which child supervision is a lower priority. Parents in these families place less emphasis on making sure homework is completed on time, monitoring children's activities outside of the home, and setting clear

rules about alcohol and drug use. Delinquent behaviors such as drug use, skipping school and carrying a weapon are also less likely to be noticed and punished.

#### School Domain:

- Within the School Domain, both middle school and high school students in Columbia County reported the highest level of risk for the *Lack of Commitment to School* scale. Among middle school students, 55% reported an elevated level of risk, equaling the statewide average. In the national normative sample, 47% reported an elevated level of risk, eight points lower than Columbia County. Among high school students, 49% reported an elevated level of risk, two points higher than the statewide average of 47%. In the national normative sample, 46% reported an elevated level of risk, three points lower than Columbia County. Students with high scores on this scale have negative feelings about school, and are less likely to report that school work is meaningful or important for their future. Young people who have lost this commitment to school are at higher risk for a variety of problem behaviors.

#### Peer and Individual Domain:

- Within the Peer and Individual Domain, both middle school and high school students in Columbia County reported the highest level of risk for the *Favorable Attitudes toward Antisocial Behavior* scale. Among middle school students, 51% reported an elevated level of risk, three points higher than the statewide average of 48%. In the national normative sample, 40% reported an elevated level of risk, 11 points lower than Columbia County. Among high school students, 44% reported an elevated level of risk, three points lower than the statewide average of 47%. In the national normative sample, 46% reported an elevated level of risk, two points higher than Columbia County. A high score on this scale indicates that fewer students express disapproval for fighting, skipping school and other forms of antisocial behavior. During the elementary school years, children usually express anticrime and prosocial attitudes and have difficulty imagining why people commit crimes or drop out of school. However, in middle school, as others they know begin to participate in such activities, their attitudes often shift toward greater acceptance of

these behaviors. This acceptance places them at higher risk for antisocial behaviors.

### ***Strengths to Build on***

In addition to specifying problem areas, the prioritization process also benefits from identifying the scales for which students reported the highest levels of protection and the lowest levels of risk. These areas represent strengths that Columbia County may wish to build on.

#### Highest Protective Factor Scales:

- Across all 10 protective factor scales, middle school students in Columbia County reported the highest level of protection for the *Family Opportunities for Prosocial Involvement* scale. Their score of 60% was four points higher than the statewide average of 56%. In the national normative sample, 59% reported an elevated level of protection, one point lower than Columbia County. High scores on this scale indicate that activities that promote family attachment—such as family recreation and involvement in family decisions—are available to students. These prosocial activities reinforce family bonds and cause students to more easily adopt the norms projected by their families. For instance, children whose parents have high expectations for their school achievement are less likely to drop out of school.
- High school students in Columbia County reported the highest level of protection for the *Religiosity* scale. Their score of 65% was four points higher than the statewide average of 61%. In the national normative sample, 62% reported an elevated level of protection, three points lower than Columbia County. Students who reported high scores on this scale attend religious services and activities more frequently. As a result, they are more likely to benefit from relationships with prosocial adults and peers, opportunities for prosocial activities, and the teaching of prosocial values that are often part of religious involvement.
- Additionally, both middle school and high school students in Columbia County reported a high level of protection for the *Community Rewards for Prosocial Involvement* scale. Among middle school students, 55% reported an elevated level of protection, four points higher than the statewide average of 51%. In the national normative sample, 56% reported an

elevated level of protection, one point higher than Columbia County. Among high school students, 64% reported an elevated level of protection, three points higher than the statewide average of 61%. In the national normative sample, 63% reported an elevated level of protection, one point lower than Columbia County. Students who reported high scores on this scale receive encouragement and praise from neighbors and other members of their communities. With this type of support, young people may be more likely to accept the guidance available from the positive role models in their communities.

- High school students also reported a high level of protection for one other scale, *Belief in the Moral Order*. Their score of 64% was five points higher than the statewide average of 59%. In the national normative sample, 62% reported an elevated level of protection, two points lower than Columbia County. High scores on this scale indicate that students are more likely to accept commonly held beliefs about what constitutes appropriate and inappropriate behavior. When students accept basic social values they may be less likely to engage in ATOD use and other forms of delinquent behavior.

#### Lowest Risk Factor Scales:

- Across all 23 risk factor scales, middle school students in Columbia County reported the lowest level of risk for the *Parental Attitudes Favorable toward ATOD Use* scale. Their score of 28% was six points higher than the statewide average of 22%. In the national normative sample, 23% reported an elevated level of protection, five points lower than Columbia County. Students with low scores on this scale have parents who strenuously disapprove of youth ATOD use. Parental attitudes is one of the strongest predictors of youth ATOD use.
- High school students in Columbia County reported the lowest level of risk for the *Family Conflict* scale. Their score of 29% was eight points lower than the statewide average of 37%. In the national normative sample, 37% reported an elevated level of protection, eight points higher than Columbia County. Students with low scores on this scale live in families where serious arguments are less common. Bonding between family members, especially between children and their parents or guardians, is a key

component in the development of positive social norms. Low levels of family conflict promote the development of these bonds, and decrease the likelihood that young people will engage in illegal drug use and other forms of delinquent behavior.

- Additionally, middle school students in Columbia County also reported a low level of risk for the *Perceived Availability of Handguns* scale. Their score of 35% was eight points higher than the statewide average of 27%. In the national normative sample, 25% reported an elevated level of protection, 10 points lower than Columbia County. A low score on this scale indicates that it is difficult for students to get a handgun.
- High school students also reported low levels of risk for two other scales. The first of these was *Friends' Use of Drugs*. Their score of 31% was eight points lower than the statewide average of 39%. In the national normative sample, 47% reported an elevated level of protection, 16 points higher than Columbia County. Students with low scores on this scale indicated that few, if any, of their best friends use alcohol, tobacco or other drugs. Young people who do not associate with drug using peers are less likely to use drugs themselves. Research has shown peer influence to be one of the strongest predictors of ATOD use—stronger, in many cases, than the influence of parents or other family members.
- The second additional risk factor scale with a low score in high school was *Laws and Norms Favorable to Handguns*. Their score of 31% was eight points higher than the statewide average of 23%. In the national normative sample, 23% reported an elevated level of protection, eight points lower than Columbia County. Students with low scores on this scale believe that police are likely to catch young people who carry handguns. When young people believe that the laws and norms concerning firearms are strictly enforced, they are less likely to engage in dangerous behavior.

### Further Considerations

In addition to identifying the highest risk factor scales and lowest protective factor scales, the prevention prioritization process may include several supplemental steps, such as:

- Compare county-level results to state-level results. Risk and protective factor scale scores from the statewide *FYSAS* are presented in Tables 18 and 19. A comparison to statewide results may reveal additional strengths and weaknesses in Columbia County's risk and protective factor profile. For example, a risk factor scale that is not the most elevated within its domain may be designated as a target for prevention programming because it is notably higher in Columbia County than across the state as a whole.
- Review the prevalence of ATOD use and other antisocial behaviors in your community. A high rate of alcohol use, for example, may dictate a different prevention strategy than a high rate of youth violence. The table on the second page in Appendix C provides a resource for this analysis by showing the behavioral outcomes that have been linked, in multiple longitudinal studies, to each risk factor.
- Use archival data to fill the gaps in the *FYSAS* data, and to support findings in the survey. For example, Teen Pregnancy and School Drop-Out are problem behaviors not measured by the survey that may influence prevention planning. Archival data are information sources that have already been collected and/or documented at the local, state or national level. They can include records that are kept by governmental and other agencies, and records that are normally kept as part of the operation of an institution or organization.
- Consider which risk and protective factors the community can realistically tackle at this time. Some factors may be too big, or there may be other efforts already underway in the community to address them. If your community does not have extensive financial or human resources, then it may be appropriate to narrow the list down to one or two priority factors.
- Consider political, social and economic factors in the community. What is best for the community? Which risk and protective factors would policy makers find acceptable to address at this time?

### Choosing Effective Prevention Strategies

After completing the prioritization process and identifying key risk and protective factors for focused prevention efforts, the next step for communities is to

select research-based, proven-effective programs that target these problem areas.

A major breakthrough in the field of positive youth development in the past two decades has been the development and testing of programs, policies and practices that are shown to work to reduce adolescent drug use, violence, risky sexual behavior and school failure. State and national agencies have become increasingly interested in and committed to programs, policies and practices that have been rigorously tested for effectiveness.

Prevention strategies identified as “tested, effective” are those that have been tested in well-controlled trials comparing schools, families, young people or communities that received the strategy with those that did not. Results of those trials showed that those who received the strategies were better off than those that did not, in terms of lower risk, greater protection and better behavioral outcomes.

A good first step in the strategy selection process is to review published lists of tested, effective prevention resources. A number of organizations have constructed lists that link research-based programs with the risk and protective factors they have been shown to effectively address. Additional information on the four lists presented below is available in Appendix D of this report.

- The *Communities That Care* Prevention Strategies Guide
- The U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration’s (SAMHSA) Model Programs list
- The University of Colorado at Boulder’s Blueprints for Violence Prevention initiative, sponsored by the Office of Juvenile Justice and Delinquency Prevention (OJJDP)
- The Western Center for the Application of Prevention Technologies (CAPT) list of Best Practices and Promising Practices

In addition to selecting research-based, proven-effective programs to target areas of low protection and high risk, communities should also consider the impact of environmental factors and public policies. For example, a strategy to combat a high level of *Perceived Availability of Drugs* might incorporate changes to local laws or provide resources to strengthen the enforcement of existing laws.

## ***Special Topics***

Several analyses were conducted to investigate alcohol, tobacco and other drug (ATOD) use results. These include early initiation of ATOD use and attitudes toward ATOD use (perceived risk of harm, personal disapproval and peer approval).

### ***Early Initiation of ATOD Use***

Students were asked to report on when they began using alcohol, cigarettes and marijuana. Early initiation for these drugs is of special importance, since they are often precursors to the use of harder drugs, such as methamphetamine and cocaine. The question related to cigarettes is “How old were you when you first smoked a cigarette, even just a puff?” The question about marijuana is “How old were you when you first smoked marijuana?” Two questions about alcohol were asked, one asking when the student first “had more than a sip or two of beer, wine or hard liquor (for example, vodka, whiskey or gin)” and one asking the student when he or she “began drinking alcoholic beverages regularly, that is, at least once or twice a month.”

Tables 13 and 17 present the percentage of high school students, age 14 years or older, who started using alcohol, cigarettes or marijuana at age 13 or younger. This percentage is the early initiation rate.

Columbia County high school students reported the highest rate of early ATOD initiation for “more than a sip or two” of alcohol (32.5%), followed by cigarette use (25.7%), marijuana use (11.6%) and drinking at least once a month (5.5%).

### ***Perceived Risk of Harm***

Perception of risk is an important determinant in the decision-making process young people go through when deciding whether or not to use alcohol, tobacco or other drugs. Evidence also suggests that the perceptions of the risks and benefits associated with drug use sometimes serve as a leading indicator of future drug use patterns in a community (Bachman, Johnston, O’Malley & Humphrey, 1986). Tables 14 and 17 present prevalence rates for surveyed Columbia County students assigning “great risk” of harm to four drug use behaviors: near daily use of alcohol, daily use of cigarettes, regular use of marijuana, and trying marijuana once or twice.

Surveyed Columbia County students assigned the highest risk of harm to regular use of marijuana (62.4%), followed by daily use of cigarettes (59.8%),



near daily use of alcohol (44.5%) and trying marijuana once or twice (39.6%).

Daily Use of Alcohol. In Columbia County, 44.5% of students reported that having one or more drinks nearly every day would pose a “great risk” of harm. This is up 10.3 percentage points from 2001. Middle school students reported a rate of 43.5% and high school students reported a rate of 45.4%. Across the state as a whole, 41.9% of students reported that near daily use of alcohol would pose a “great risk” of harm.

Daily Use of Cigarettes. In Columbia County, 59.8% of students reported that smoking a pack or more of cigarettes every day would pose a “great risk” of harm. This is down 2.6 percentage points from 2001. Middle school students reported a rate of 60.2% and high school students reported a rate of 59.5%. Across the state as a whole, 67.6% of students reported that near daily use of cigarettes would pose a “great risk” of harm.

Regular Use of Marijuana. In Columbia County, 62.4% of students reported that smoking marijuana regularly would pose a “great risk” of harm. Students reported the same rate in 2001. Middle school students reported a rate of 70.1% and high school students reported a rate of 55.9%. Across the state as a whole, 59.8% of students reported that smoking marijuana regularly would pose a “great risk” of harm.

Trying Marijuana Once or Twice. In Columbia County, 39.6% of students reported that trying marijuana once or twice would pose a “great risk” of harm. This is up 2.4 percentage points from 2001. Middle school students reported a rate of 48.7% and high school students reported a rate of 31.9%. Across the state as a whole, 32.5% of students reported trying marijuana once or twice would pose a “great risk” of harm.

### ***Personal Disapproval***

In addition to perceptions of risk, personal approval or disapproval of drugs has been linked to the prevalence of ATOD use (Bachman, Johnston & O’Malley, 1996). Personal disapproval was measured by asking students how wrong it would be for someone their age to drink alcohol regularly, smoke cigarettes, smoke marijuana, or use other illicit drugs (“LSD, cocaine, amphetamines or another illegal drug”). The rates presented in Tables 14 and 17 represent the percentages of students who thought it would be “wrong” or “very wrong” to use each drug.

Surveyed Columbia County students were most likely to disapprove of other illicit drug use (95.3%), followed by smoking marijuana (80.8%), smoking cigarettes (78.3%) and drinking alcohol regularly (68.6%).

Smoking Cigarettes. In Columbia County, 78.3% of students reported that they think it would be “wrong” or “very wrong” for someone their age to smoke cigarettes. This is up 4.2 percentage points from 2001. Middle school students reported a rate of 85.0% and high school students reported a rate of 72.6%. Across the state as a whole, 80.5% of students reported disapproval of smoking cigarettes.

Drinking Alcohol Regularly. In Columbia County, 68.6% of students reported that they think it would be “wrong” or “very wrong” for someone their age to drink alcohol regularly. This is up 5.3 percentage points from 2001. Middle school students reported a rate of 78.9% and high school students reported a rate of 59.8%. Across the state as a whole, 65.4% of students reported disapproval of drinking alcohol regularly.

Smoking Marijuana. In Columbia County, 80.8% of students reported that they think it would be “wrong” or “very wrong” for someone their age to smoke marijuana. This is up 1.7 percentage points from 2001. Middle school students reported a rate of 85.5% and high school students reported a rate of 76.8%. Across the state as a whole, 80.2% of students reported disapproval of smoking marijuana.

Using Other Illicit Drugs. In Columbia County, 95.3% of students reported that they think it would be “wrong” or “very wrong” for someone their age to use other illicit drugs. This is up 0.8 percentage points from 2001. Middle school students reported a rate of 95.3% and high school students reported a rate of 95.2%. Across the state as a whole, 94.9% of students reported disapproval of using other illicit drugs.

### ***Peer Approval***

In addition to perceived risk of harm and disapproval, expectations of how one’s peer group might react have an impact on whether or not young people choose to use drugs. The data presented in Tables 14 and 17 show the percentage of students who said that there is a “pretty good” or “very good” chance that they would be seen as cool if they smoked cigarettes, drank alcohol regularly or smoked marijuana.

Drinking Alcohol Regularly. In Columbia County, 13.6% of students reported that there is a “pretty

good” or a “very good” chance that they would be seen as cool if they drank alcohol regularly. This is up 1.5 percentage points from 2001. Middle school students reported a rate of 13.5% and high school students reported a rate of 13.7%. Across the state as a whole, 12.0% of students reported peer approval of drinking alcohol regularly.

**Smoking Cigarettes.** In Columbia County, 5.7% of students reported that there is a “pretty good” or a “very good” chance that they would be seen as cool if they smoked cigarettes. This is down 1.7 percentage points from 2001. Middle school students reported a rate of 8.6% and high school students reported a rate of 3.3%. Across the state as a whole, 5.8% of students reported peer approval of smoking cigarettes.

**Smoking Marijuana.** In Columbia County, 10.4% of students reported that there is a “pretty good” or a “very good” chance that they would be seen as cool if they smoked marijuana. This is down 3.1 percentage points from 2001. Middle school students reported a rate of 11.1% and high school students reported a rate of 9.7%. Across the state as a whole, 11.3% of students reported peer approval of smoking marijuana.

#### ***Extracurricular Activities***

In 2006 a new item was added to the *FYSAS* questionnaire that measures participation in five extracurricular activities: school sports, organized sports outside of school, school band, school clubs, and community clubs. Results for these items are presented in Table 15. Participation in these activities help students build stronger ties to their school and community. Through these connections students are also more likely to develop attachments to prosocial peers and to positive adult role models. Since these bonds encourage students to engage in developmentally positive activity, they serve as a

buffer against ATOD use and other antisocial behaviors. Please note that this measure is similar to two of the protective factor scales discussed earlier in this report: *Community Opportunities for Prosocial Involvement* and *School Opportunities for Prosocial Involvement*.

**School Sports.** In Columbia County, 43.2% of students reported participation in school sports. Middle school students participated at a rate of 45.5% and high school students participated at a rate of 41.1%. Across the state as a whole, the rate of participation was 37.0%.

**Organized Sports Outside of School.** In Columbia County, 28.9% of students reported participation in organized sports outside of school. Middle school students participated at a rate of 34.7% and high school students participated at a rate of 24.1%. Across the state as a whole, the rate of participation was 33.7%.


**School Band.** In Columbia County, 9.2% of students reported participation in school band. Middle school students participated at a rate of 15.3% and high school students participated at a rate of 4.1%. Across the state as a whole, the rate of participation was 10.0%.

**School Clubs.** In Columbia County, 35.6% of students reported participation in school clubs. Middle school students participated at a rate of 25.4% and high school students participated at a rate of 44.3%. Across the state as a whole, the rate of participation was 26.4%.

**Community Clubs.** In Columbia County, 6.1% of students reported participation in community clubs. Middle school students participated at a rate of 6.6% and high school students participated at a rate of 5.6%. Across the state as a whole, the rate of participation was 12.4%.

**Graph  
13**

### Bullying-related behaviors among Columbia County MIDDLE SCHOOL students, 2008



#### ***Bullying Behavior***

In 2008 a new item set was added to the *FYSAS* middle school questionnaire that assesses student involvement with bullying. The new items include (1) worry or fear due to bullying, (2) skipping school because of being bullied, (3) being physically bullied (kicking, shoving, stealing, etc.), (4) being verbally bullied (taunting, teasing, name-calling, etc.), (5) being cyber bullied (mean emails, mean text messages, etc.), (6) physically bullying others, (7) verbally bullying others, and (8) cyber bullying others. Table 12 and Graph 13 present prevalence rates for these behaviors.

***Worry or Fear.*** In Columbia County, 33.2% of middle school students reported that bullying causes them to be “somewhat” or “a whole lot” worried or fearful. Across the state as a whole, 30.1% reported being worried or fearful.

***Skipping School.*** In Columbia County, 3.8% of middle school students reported skipping school because someone was bullying them. Across the state as a whole, 2.9% reported skipping school because of bullying.

***Was Physically Bullied.*** In Columbia County, 19.9% of middle school students reported experiencing

“somewhat” or “a whole lot” of physical bullying in the past 30 days. Across the state as a whole, 20.7% reported experiencing physical bullying.

***Was Verbally Bullied.*** In Columbia County, 40.0% of middle school students reported experiencing “somewhat” or “a whole lot” of verbal bullying in the past 30 days. Across the state as a whole, 41.3% reported experiencing verbal bullying.

***Was Cyber Bullied.*** In Columbia County, 6.5% of middle school students reported experiencing “somewhat” or “a whole lot” of cyber bullying in the past 30 days. Across the state as a whole, 8.2% reported experiencing cyber bullying.

***Physically Bullied Others.*** In Columbia County, 12.1% of middle school students reported physically bullying others “somewhat” or “a whole lot” in the past 30 days. Across the state as a whole, 15.8% reported physically bullying others.

***Verbally Bullied Others.*** In Columbia County, 23.4% of middle school students reported verbally bullying others “somewhat” or “a whole lot” in the past 30 days. Across the state as a whole, 28.0% reported verbally bullying others.



Cyber Bullied Others. In Columbia County, 4.6% of middle school students reported cyber bullying others “somewhat” or “a whole lot” in the past 30 days. Across the state as a whole, 6.4% reported cyber bullying others.





# Appendix A

## Detailed Tables

---

**Table 1. Major demographic characteristics of surveyed Columbia County youth and Florida Statewide youth**

	Columbia County			Florida Statewide	
	N	%		N	%
<b>Sex</b>					
Female	434	53.1		43,913	48.0
Male	361	44.1		45,413	49.6
<b>Race/Ethnic group</b>					
African American	142	17.4		16,647	18.2
American Indian	15	1.8		1,011	1.1
Asian	6	0.7		1,994	2.2
Hispanic/Latino	36	4.4		20,767	22.7
Native Hawaiian/Pacific Islander	6	0.7		490	0.5
Other/Multiple	76	9.3		12,821	14.0
White, non-Hispanic	531	64.9		37,000	40.4
<b>Age</b>					
10	1	0.1		98	0.1
11	37	4.5		3,294	3.6
12	122	14.9		10,971	12.0
13	110	13.4		13,299	14.5
14	145	17.7		14,098	15.4
15	145	17.7		14,339	15.7
16	100	12.2		13,913	15.2
17	94	11.5		12,824	14.0
18	49	6.0		7,552	8.3
19 or older	10	1.2		718	0.8
<b>Grade</b>					
6th	177	21.6		13,265	14.5
7th	98	12.0		13,552	14.8
8th	149	18.2		12,869	14.1
9th	154	18.8		14,738	16.1
10th	82	10.0		13,593	14.9
11th	100	12.2		12,297	13.4
12th	58	7.1		11,157	12.2
Overall Middle School	424	51.8		39,686	43.4
Overall High School	394	48.2		51,785	56.6
<b>Total</b>	<b>818</b>	<b>100.0</b>		<b>91,471</b>	<b>100.0</b>

Note: Some categories do not sum to 100% of the total due to missing values (e.g., not all survey questions were answered). In addition, rounding can produce totals that do not equal 100%. "N" represents the number of valid cases. In this table, county data are unweighted while statewide data are weighted.

**Table 2. Percentages of Columbia County youth and Florida Statewide youth who reported having used various drugs in their lifetimes**

	Columbia County							Florida Statewide						
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
Alcohol	40.4	56.5	50.3	48.4	38.4	56.0	48.9	36.3	66.2	54.9	51.5	37.4	65.6	53.2
Cigarettes	26.2	35.0	27.5	34.1	24.0	36.1	30.8	17.4	34.4	27.4	26.4	17.1	34.3	27.0
Smokeless Tobacco	--	20.4	8.9	34.9	--	20.8	--	--	12.5	6.9	18.1	--	12.0	--
Marijuana or Hashish	14.0	23.9	18.7	20.1	12.3	24.2	19.3	8.5	30.8	20.0	22.1	8.4	30.5	21.1
Inhalants	13.4	13.3	13.2	13.6	13.6	11.7	13.3	13.2	10.1	12.9	10.0	13.0	10.5	11.4
Over-The-Counter Drugs	5.7	--	6.1	5.7	4.8	--	--	4.9	--	5.9	3.9	4.6	--	--
Any illicit drug	24.0	31.6	27.8	28.9	22.2	31.6	28.0	21.2	37.5	30.7	30.1	21.1	37.4	30.4
Any illicit drug other than marijuana	18.8	22.9	21.4	21.0	18.1	22.3	21.0	17.5	23.1	21.7	19.6	17.4	23.0	20.7
Alcohol only	21.6	27.5	26.4	23.6	21.2	27.0	24.7	20.8	31.7	28.0	25.9	21.8	31.3	26.9
Alcohol or any illicit drug	45.4	58.6	53.9	52.0	43.2	58.1	52.4	41.9	68.9	58.5	55.7	42.7	68.4	57.1
Any illicit drug, but no alcohol	5.1	2.5	4.1	3.5	4.9	2.4	3.7	5.8	3.0	3.9	4.6	5.6	3.1	4.2

Note: The first set of data rows show results for alcohol, tobacco, marijuana, inhalants and over-the-counter drugs. The second set of data rows show results for various combinations of drugs. The symbol "--" indicates that data are not available.

**Table 3. Percentages of Columbia County youth and Florida Statewide youth who reported having used various drugs in their lifetimes**

	Columbia County							Florida Statewide						
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
Club Drugs <sup>1</sup>	2.1	--	1.9	2.3	2.0	--	--	1.6	--	1.4	1.8	1.4	--	--
LSD, PCP or Mushrooms	1.9	--	1.5	2.5	1.9	--	--	1.5	--	1.1	1.9	1.4	--	--
Cocaine or Crack Cocaine	1.6	--	1.5	1.7	1.7	--	--	1.8	--	1.7	1.9	1.7	--	--
Ecstasy	--	4.4	4.2	5.0	--	5.0	--	--	4.9	4.9	5.1	--	4.7	--
Rohypnol	--	0.8	0.4	1.4	--	0.6	--	--	1.2	1.0	1.4	--	1.2	--
GHB	--	1.3	1.2	1.4	--	1.7	--	--	0.7	0.4	0.9	--	0.6	--
Ketamine	--	0.0	0.0	0.0	--	0.0	--	--	0.8	0.6	0.9	--	0.7	--
LSD or PCP	--	2.1	2.1	2.2	--	1.9	--	--	3.0	2.3	3.7	--	2.9	--
Hallucinogenic Mushrooms	--	6.1	4.5	7.8	--	6.2	--	--	5.3	3.9	6.8	--	5.1	--
Cocaine	--	3.9	4.5	3.5	--	3.6	--	--	5.5	5.5	5.6	--	5.4	--
Crack Cocaine	--	1.1	1.6	0.5	--	0.9	--	--	1.8	1.8	1.7	--	1.7	--
Methamphetamine	1.3	1.8	2.5	0.7	1.3	1.6	1.6	1.2	1.5	1.3	1.4	1.2	1.5	1.4
Heroin	1.0	0.5	0.5	1.0	1.0	0.4	0.7	0.8	1.0	0.8	1.1	0.8	1.0	0.9
Depressants	3.0	6.7	5.5	4.0	2.7	7.6	4.9	2.4	8.7	6.5	5.4	2.5	8.5	6.0
Prescription Pain Relievers	6.0	11.4	9.6	8.1	4.6	12.1	8.9	4.9	10.4	8.3	7.6	4.8	10.3	8.0
Prescription Amphetamines	1.8	1.4	1.8	1.1	1.8	1.7	1.6	1.6	5.3	3.8	3.5	1.6	5.0	3.7
Steroids	1.2	1.5	0.7	2.2	1.3	1.9	1.4	0.8	1.2	0.6	1.4	0.8	1.2	1.0

Note: The first set of data rows show results for items that are on the middle school questionnaire. The second set of data rows show results for items that are on the high school questionnaire. The third set of data rows show results for items that are on both questionnaires. The symbol "--" indicates that data are not available.

<sup>1</sup> Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs.

**Table 4. Percentages of Columbia County youth and Florida Statewide youth who reported having used various drugs in the past 30 days**

	Columbia County							Florida Statewide						
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
Alcohol	21.0	28.3	25.5	24.2	18.9	28.2	24.9	17.3	39.5	30.6	29.0	17.8	38.6	29.8
Binge Drinking	11.4	20.5	16.6	15.5	10.8	20.1	16.2	6.2	21.5	14.0	15.6	6.4	20.5	14.8
Cigarettes	8.8	14.0	10.1	12.9	7.0	15.1	11.6	4.7	12.6	8.8	9.4	4.5	12.2	9.1
Smokeless Tobacco	--	9.2	2.4	17.2	--	10.0	--	--	5.3	2.4	8.3	--	5.0	--
Marijuana or Hashish	7.9	12.4	9.4	11.6	6.7	13.5	10.3	4.4	16.2	9.8	12.3	4.3	16.1	11.1
Inhalants	4.2	2.7	3.3	3.5	4.0	2.6	3.4	5.2	2.2	4.1	2.8	4.9	2.4	3.5
Over-The-Counter Drugs	3.1	--	3.1	3.3	2.2	--	--	2.2	--	2.6	1.8	2.1	--	--
Any illicit drug	14.2	15.9	14.6	15.7	11.6	18.3	15.1	10.6	19.9	15.2	16.3	10.3	19.8	15.8
Any illicit drug other than marijuana	9.7	8.8	8.9	9.4	7.9	10.5	9.2	8.0	9.7	9.1	8.7	7.8	9.5	8.9
Alcohol only	12.9	17.4	15.8	15.0	12.3	16.9	15.3	11.4	24.0	19.7	17.4	12.0	23.5	18.5
Alcohol or any illicit drug	26.5	32.2	29.7	29.6	23.2	34.0	29.6	21.7	43.3	34.6	33.2	22.0	42.7	33.9
Any illicit drug, but no alcohol	5.8	4.3	4.6	5.7	4.5	6.4	5.0	4.7	4.2	4.2	4.5	4.5	4.4	4.4

Note: The first set of data rows show results for alcohol, tobacco, marijuana, inhalants and over-the-counter drugs. The second set of data rows show results for various combinations of drugs. The symbol "--" indicates that data are not available.

**Table 5. Percentages of Columbia County youth and Florida Statewide youth who reported having used various drugs in the past 30 days**

	Columbia County							Florida Statewide						
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
Club Drugs <sup>1</sup>	1.6	--	0.8	2.4	1.5	--	--	0.6	--	0.5	0.7	0.5	--	--
LSD, PCP or Mushrooms	2.0	--	0.8	3.3	1.6	--	--	0.6	--	0.3	0.8	0.5	--	--
Cocaine or Crack Cocaine	1.1	--	0.8	1.5	0.8	--	--	0.6	--	0.5	0.7	0.5	--	--
Ecstasy	--	1.7	1.6	1.9	--	1.6	--	--	1.5	1.2	1.9	--	1.4	--
Rohypnol	--	0.0	0.0	0.0	--	0.0	--	--	0.4	0.3	0.6	--	0.4	--
GHB	--	0.8	0.8	0.8	--	1.0	--	--	0.3	0.1	0.4	--	0.2	--
Ketamine	--	0.0	0.0	0.0	--	0.0	--	--	0.2	0.1	0.3	--	0.2	--
LSD or PCP	--	0.7	0.3	1.2	--	1.0	--	--	1.1	0.7	1.5	--	1.1	--
Hallucinogenic Mushrooms	--	1.7	1.2	2.5	--	2.0	--	--	1.5	0.9	2.0	--	1.3	--
Cocaine	--	1.3	1.5	1.3	--	1.2	--	--	1.6	1.4	1.7	--	1.5	--
Crack Cocaine	--	0.2	0.0	0.5	--	0.0	--	--	0.5	0.4	0.6	--	0.5	--
Methamphetamine	1.1	0.4	1.2	0.4	1.2	0.5	0.8	0.6	0.5	0.4	0.6	0.5	0.5	0.5
Heroin	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.4	0.3	0.2	0.5	0.3	0.3	0.3
Depressants	1.6	2.7	2.5	1.7	1.0	4.0	2.2	1.0	3.0	2.4	2.0	1.0	3.0	2.1
Prescription Pain Relievers	3.7	3.8	3.4	4.4	2.5	4.4	3.8	2.3	3.9	3.2	3.2	2.3	3.8	3.2
Prescription Amphetamines	1.2	0.2	0.9	0.4	1.2	0.2	0.7	0.6	1.6	1.0	1.3	0.7	1.5	1.2
Steroids	0.0	0.6	0.3	0.4	0.0	0.8	0.3	0.3	0.5	0.2	0.7	0.3	0.5	0.4

Note: The first set of data rows show results for items that are on the middle school questionnaire. The second set of data rows show results for items that are on the high school questionnaire. The third set of data rows show results for items that are on both questionnaires. The symbol "--" indicates that data are not available.

<sup>1</sup> Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs.

**Table 6. Lifetime trend in alcohol, tobacco and other drug use for Columbia County youth, 2001, 2002, 2004, 2006 and 2008**

	2001			2002			2004			2006			2008		
	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>
Alcohol	39.0	69.8	54.4	40.3	68.9	54.8	43.7	66.1	55.6	37.3	74.6	57.4	40.4	56.5	48.9
Cigarettes	34.3	57.7	46.2	35.6	50.0	42.9	33.1	50.7	42.3	27.3	45.0	37.2	26.2	35.0	30.8
Smokeless Tobacco	17.2	22.9	20.1	19.4	24.0	21.7	20.2	26.5	23.5	14.9	24.3	20.5	--	20.4	--
Marijuana or Hashish	13.8	33.9	24.0	13.5	33.9	24.0	13.1	34.6	24.6	12.7	28.5	21.9	14.0	23.9	19.3
Inhalants	12.2	18.3	15.3	11.5	12.2	12.0	15.1	17.3	16.4	7.6	10.9	10.2	13.4	13.3	13.3
Over-The-Counter Drugs	--	--	--	--	--	--	--	--	--	--	--	--	5.7	--	--
Any illicit drug	--	--	--	24.1	40.2	32.4	25.0	43.1	34.6	19.2	35.2	28.3	24.0	31.6	28.0
Any illicit drug other than marijuana	--	--	--	19.0	23.4	21.3	20.3	30.6	25.8	12.9	30.2	22.8	18.8	22.9	21.0
Alcohol only	--	--	--	22.4	32.0	27.3	23.6	26.8	25.2	21.4	40.8	31.4	21.6	27.5	24.7
Alcohol or any illicit drug	--	--	--	46.1	72.0	59.3	47.7	69.8	59.3	40.7	75.7	59.5	45.4	58.6	52.4
Any illicit drug, but no alcohol	--	--	--	6.4	2.9	4.8	4.9	3.9	4.3	3.7	1.6	2.5	5.1	2.5	3.7

Note: The first set of data rows show results for alcohol, tobacco, marijuana, inhalants and over-the-counter drugs. The second set of data rows show results for various combinations of drugs. Results for combinations of drugs are not presented for 2001 because new ATOD items were added in 2002. The symbol "--" indicates that data are not available.



**Table 7. Lifetime trend in alcohol, tobacco and other drug use for Columbia County youth, 2001, 2002, 2004, 2006 and 2008**

	2001			2002			2004			2006			2008		
	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>
Club Drugs <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	2.1	--	--
LSD, PCP or Mushrooms	--	--	--	--	--	--	--	--	--	--	--	--	1.9	--	--
Cocaine or Crack Cocaine	--	--	--	--	--	--	--	--	--	--	--	--	1.6	--	--
Ecstasy	--	--	--	4.0	8.2	6.1	2.0	5.7	4.0	1.3	11.1	7.5	--	4.4	--
Rohypnol	--	--	--	1.1	1.6	1.3	0.8	1.9	1.4	1.0	0.0	0.4	--	0.8	--
GHB <sup>2</sup>	--	--	--	3.0	0.2	1.6	1.2	0.4	0.8	0.9	0.0	0.4	--	1.3	--
Ketamine	--	--	--	2.0	1.3	1.7	0.6	1.1	0.9	0.7	0.0	0.3	--	0.0	--
LSD or PCP <sup>3</sup>	2.2	10.8	6.5	1.5	2.7	2.2	1.3	1.8	1.5	1.3	0.0	0.6	--	2.1	--
Hallucinogenic Mushrooms	--	--	--	6.5	7.7	7.1	4.8	9.6	7.4	2.6	10.5	6.8	--	6.1	--
Cocaine	1.4	7.3	4.4	1.7	4.3	3.0	1.7	6.6	4.3	1.4	4.8	4.1	--	3.9	--
Crack Cocaine	2.0	2.4	2.2	1.5	0.7	1.1	1.1	2.9	2.0	1.3	1.2	2.1	--	1.1	--
Methamphetamine	1.9	9.3	5.7	2.1	2.0	2.0	1.7	3.5	2.7	1.5	0.0	1.6	1.3	1.8	1.6
Heroin	1.9	2.6	2.3	1.1	0.8	0.9	0.6	0.9	0.8	0.8	0.6	0.7	1.0	0.5	0.7
Depressants <sup>4</sup>	3.5	11.3	7.4	3.9	8.1	6.0	3.9	12.1	8.3	3.2	14.2	10.0	3.0	6.7	4.9
Prescription Pain Relievers <sup>5</sup>	--	--	--	5.4	11.7	8.6	8.1	16.2	12.4	3.4	20.1	13.3	6.0	11.4	8.9
Prescription Amphetamines	--	--	--	3.3	4.2	3.8	3.3	4.7	4.1	3.0	2.2	3.5	1.8	1.4	1.6
Steroids	3.5	3.1	3.3	2.7	0.8	1.7	2.2	1.4	1.7	1.3	0.6	1.8	1.2	1.5	1.4

Note: The first set of data rows show results for items that are on the middle school questionnaire. The second set of data rows show results for items that are on the high school questionnaire. The third set of data rows show results for items that are on both questionnaires. The symbol "--" indicates that data are not available.

<sup>1</sup> Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs.

<sup>2</sup> In 2006, the question for GHB was changed to include a more up-to-date set of slang or street names for the drug.

<sup>3</sup> Measured as "LSD or other psychedelics" in the 2000 survey, and as "LSD or PCP" in the 2001, 2004, 2006 and 2008 surveys.

<sup>4</sup> In 2002, the prescription drug Xanax<sup>®</sup> was added to the list of examples given in the depressants question.

<sup>5</sup> In 2006, OxyContin<sup>®</sup> was removed as an individual item and added to the list of examples included in the prescription pain relievers item.

**Table 8. Past-30-day trend in alcohol, tobacco and other drug use for Columbia County youth, 2001, 2002, 2004, 2006 and 2008**

	2001			2002			2004			2006			2008		
	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>
Alcohol	19.5	46.2	33.0	18.5	39.6	29.2	21.5	39.3	30.9	17.0	38.1	28.9	21.0	28.3	24.9
Binge Drinking	8.0	24.8	16.4	10.7	24.3	17.6	13.5	24.7	19.4	9.0	20.7	16.0	11.4	20.5	16.2
Cigarettes	10.0	25.7	18.0	12.3	17.4	14.8	11.4	22.1	17.0	8.5	10.7	10.6	8.8	14.0	11.6
Smokeless Tobacco	7.2	12.4	9.8	5.5	9.8	7.7	7.8	11.7	9.8	8.2	10.6	10.2	--	9.2	--
Marijuana or Hashish	7.5	19.0	13.4	6.5	17.0	12.0	6.7	12.6	9.8	7.5	8.0	7.6	7.9	12.4	10.3
Inhalants	4.7	4.5	4.6	3.7	2.5	3.3	5.3	3.4	4.3	3.0	1.9	3.3	4.2	2.7	3.4
Over-The-Counter Drugs	--	--	--	--	--	--	--	--	--	--	--	--	3.1	--	--
Any illicit drug	--	--	--	13.4	18.6	16.3	12.3	16.4	14.5	11.4	12.2	12.6	14.2	15.9	15.1
Any illicit drug other than marijuana	--	--	--	10.2	7.0	8.8	8.5	8.2	8.3	6.4	8.5	8.4	9.7	8.8	9.2
Alcohol only	--	--	--	12.0	25.5	18.8	13.2	26.6	20.3	10.6	27.4	19.5	12.9	17.4	15.3
Alcohol or any illicit drug	--	--	--	24.9	43.3	34.4	24.8	41.9	33.8	22.0	40.5	32.4	26.5	32.2	29.6
Any illicit drug, but no alcohol	--	--	--	6.9	3.7	5.5	3.6	2.9	3.2	5.2	2.7	3.8	5.8	4.3	5.0

Note: The first set of data rows show results for alcohol, tobacco, marijuana, inhalants and over-the-counter drugs. The second set of data rows show results for various combinations of drugs. Results for combinations of drugs are not presented for 2001 because new ATOD items were added in 2002. The symbol "--" indicates that data are not available.

**Table 9. Past-30-day trend in alcohol, tobacco and other drug use for Columbia County youth, 2001, 2002, 2004, 2006 and 2008**

	2001			2002			2004			2006			2008		
	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>
Club Drugs <sup>1</sup>	--	--	--	--	--	--	--	--	--	--	--	--	1.6	--	--
LSD, PCP or Mushrooms	--	--	--	--	--	--	--	--	--	--	--	--	2.0	--	--
Cocaine or Crack Cocaine	--	--	--	--	--	--	--	--	--	--	--	--	1.1	--	--
Ecstasy	--	--	--	1.6	2.9	2.2	0.5	1.4	1.0	0.8	0.0	1.3	--	1.7	--
Rohypnol	--	--	--	0.4	0.6	0.5	0.0	0.5	0.2	0.5	0.0	0.2	--	0.0	--
GHB <sup>2</sup>	--	--	--	2.6	0.2	1.5	0.5	0.2	0.4	0.0	0.0	0.0	--	0.8	--
Ketamine	--	--	--	0.2	0.4	0.3	0.3	0.0	0.1	0.0	0.0	0.0	--	0.0	--
LSD or PCP <sup>3</sup>	1.3	0.0	0.6	0.7	0.7	0.7	0.5	0.3	0.4	0.5	0.0	0.2	--	0.7	--
Hallucinogenic Mushrooms	--	--	--	2.5	1.8	2.1	0.9	0.8	0.9	0.8	0.0	0.4	--	1.7	--
Cocaine	1.0	0.4	0.7	0.4	1.0	0.7	0.3	2.0	1.2	0.7	4.3	3.6	--	1.3	--
Crack Cocaine	0.8	0.0	0.4	0.4	0.3	0.4	0.9	0.3	0.6	0.7	0.0	1.3	--	0.2	--
Methamphetamine	1.0	3.3	2.2	1.2	0.6	0.9	1.0	0.3	0.6	0.2	0.0	0.1	1.1	0.4	0.8
Heroin	0.6	0.0	0.3	0.2	0.4	0.3	0.3	0.0	0.1	0.3	0.0	0.1	0.0	0.0	0.0
Depressants <sup>4</sup>	1.9	1.3	1.6	2.2	2.6	2.5	1.4	3.6	2.6	0.8	3.7	3.3	1.6	2.7	2.2
Prescription Pain Relievers <sup>5</sup>	--	--	--	2.3	5.2	3.9	2.7	4.1	3.4	1.0	2.5	2.8	3.7	3.8	3.8
Prescription Amphetamines	--	--	--	1.4	0.3	0.8	1.7	0.3	1.0	0.6	0.6	1.5	1.2	0.2	0.7
Steroids	1.3	2.3	1.8	0.5	0.5	0.5	1.3	0.0	0.6	0.3	0.0	1.1	0.0	0.6	0.3

Note: The first set of data rows show results for items that are on the middle school questionnaire. The second set of data rows show results for items that are on the high school questionnaire. The third set of data rows show results for items that are on both questionnaires. The symbol "--" indicates that data are not available.

<sup>1</sup> Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs.

<sup>2</sup> In 2006, the question for GHB was changed to include a more up-to-date set of slang or street names for the drug.

<sup>3</sup> Measured as "LSD or other psychedelics" in the 2000 survey, and as "LSD or PCP" in the 2001, 2004, 2006 and 2008 surveys.

<sup>4</sup> In 2002, the prescription drug Xanax<sup>®</sup> was added to the list of examples given in the depressants question.

<sup>5</sup> In 2006, OxyContin<sup>®</sup> was removed as an individual item and added to the list of examples included in the prescription pain relievers item.

**Table 10. Percentages of Columbia County youth and Florida Statewide youth who reported engaging in delinquent behavior within the past 12 months**

	Columbia County							Florida Statewide						
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
Carrying a handgun	6.1	6.0	1.5	10.4	4.7	6.9	6.0	4.3	5.6	2.1	8.0	3.9	5.8	5.0
Selling drugs	2.8	4.6	1.8	5.8	2.4	5.2	3.7	2.7	7.6	3.2	7.8	2.5	7.9	5.5
Attempting to steal a vehicle	2.8	1.7	1.4	3.0	2.1	2.7	2.3	2.4	2.6	1.7	3.3	2.2	2.9	2.5
Being arrested	5.4	4.9	3.1	7.3	3.5	6.1	5.2	4.0	5.6	3.3	6.4	3.6	6.2	4.9
Taking a handgun to school	1.2	0.7	0.2	1.9	1.0	0.6	0.9	0.8	1.1	0.4	1.5	0.8	1.2	1.0
Getting suspended	15.3	14.2	10.9	18.1	13.2	16.4	14.7	16.2	14.4	11.5	18.9	14.9	16.1	15.2
Attacking someone with intent to harm	11.1	10.3	5.6	15.9	10.3	9.6	10.6	11.4	12.1	9.9	13.7	10.9	12.9	11.8
Being drunk or high at school	9.0	15.8	10.7	14.8	8.2	16.7	12.6	6.6	15.5	10.9	12.3	6.4	15.8	11.6

**Table 11. Percentages of Columbia County youth and Florida Statewide youth who reported gambling and arguing about gambling in the past 12 months**

	Columbia County							Florida Statewide						
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
Gambling	51.7	52.6	41.1	63.8	50.6	54.0	52.2	57.0	54.0	44.7	65.6	56.8	54.3	55.3
Arguing about gambling	14.6	12.0	8.4	17.9	13.6	12.9	13.2	17.4	14.4	11.5	19.7	17.1	14.8	15.7

**Table 12. Percentages of Columbia County youth and Florida Statewide middle school youth who reported involvement in bullying behavior**

	Columbia County							Florida Statewide						
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
Bullying caused worry	33.2	--	37.1	29.9	34.3	--	--	30.1	--	36.0	24.6	30.7	--	--
Skipped school because of bullying	3.8	--	4.0	3.7	3.7	--	--	2.9	--	3.5	2.3	2.8	--	--
Was kicked or shoved in past 30 days	19.9	--	16.8	22.8	20.4	--	--	20.7	--	17.3	23.9	21.1	--	--
Was taunted or teased in past 30 days	40.0	--	35.8	43.3	41.0	--	--	41.3	--	42.3	40.4	41.8	--	--
Was victim of cyber bullying in past 30 days	6.5	--	8.5	5.0	6.2	--	--	8.2	--	11.2	5.2	8.2	--	--
Physically bullied others in past 30 days	12.1	--	8.3	15.0	10.8	--	--	15.8	--	13.3	18.1	15.5	--	--
Verbally bullied others in past 30 days	23.4	--	21.0	24.9	24.0	--	--	28.0	--	27.8	28.2	27.8	--	--
Cyber bullied others in past 30 days	4.6	--	5.4	4.2	4.8	--	--	6.4	--	7.9	4.8	6.2	--	--

The symbol "--" indicates that data are not available.

**Table 13. Percentages of Columbia County youth and Florida Statewide high school youth who started using alcohol at age 13 or younger**

	Columbia County							Florida Statewide						
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
More than a sip of alcohol	--	32.5	30.8	35.2	--	33.5	--	--	32.3	31.0	33.9	--	32.4	--
Drinking at least once a month	--	5.5	5.4	5.3	--	6.8	--	--	5.9	5.5	6.4	--	5.9	--
Cigarettes	--	25.7	22.6	29.7	--	26.7	--	--	19.9	19.6	20.3	--	20.2	--
Marijuana	--	11.6	10.2	13.8	--	12.7	--	--	10.6	8.8	12.4	--	10.7	--

Note: The symbol "--" indicates that data are not available.

**Table 14. Percentages of Columbia County youth and Florida Statewide youth who reported a perceived risk of harm, personal disapproval and peer approval**

	Columbia County							Florida Statewide						
	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total	Middle School	High School	Female	Male	Ages 10-14	Ages 15-17	Total
<b><i>Perceive great risk of harm if...</i></b>														
One or more drinks every day	43.5	45.4	51.2	38.4	44.9	45.1	44.5	43.2	40.8	46.2	37.6	43.4	40.7	41.9
Smoke a pack or more every day	60.2	59.5	67.0	54.0	62.6	59.8	59.8	67.0	68.0	70.6	64.7	68.3	67.3	67.6
Smoke marijuana regularly	70.1	55.9	67.4	58.3	71.2	57.0	62.4	71.2	51.0	63.8	56.0	71.7	50.7	59.8
Try marijuana once or twice	48.7	31.9	42.9	36.5	48.9	33.0	39.6	42.6	24.7	34.3	30.8	42.0	24.6	32.5
<b><i>Think it would be wrong for someone their age to...</i></b>														
Smoke cigarettes	85.0	72.6	79.6	77.6	87.0	73.3	78.3	90.5	72.9	80.7	80.3	90.4	74.9	80.5
Drink alcohol regularly	78.9	59.8	68.7	68.2	79.6	61.9	68.6	81.2	53.5	65.7	65.4	80.2	54.6	65.4
Smoke marijuana	85.5	76.8	81.5	80.6	87.6	76.3	80.8	91.1	71.9	82.2	78.4	91.0	72.3	80.2
Use other illicit drugs	95.3	95.2	95.7	95.1	96.0	94.8	95.3	96.8	93.5	95.6	94.2	96.8	93.4	94.9
<b><i>Good chance of being seen as cool if...</i></b>														
Drink alcohol regularly	13.5	13.7	13.4	13.9	12.6	15.8	13.6	8.1	14.9	12.5	11.5	8.4	14.6	12.0
Smoke cigarettes	8.6	3.3	5.1	6.5	7.4	4.9	5.7	6.2	5.6	5.7	6.0	6.1	5.6	5.8
Smoke marijuana	11.1	9.7	8.8	11.6	9.9	10.7	10.4	9.2	12.9	11.1	11.5	9.1	13.4	11.3

**Table 15. Percentages of Columbia County youth and Florida Statewide youth who reported participation in extracurricular activities**

	Columbia County							Florida Statewide						
	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Female</i>	<i>Male</i>	<i>Ages 10-14</i>	<i>Ages 15-17</i>	<i>Total</i>
School Sports	45.5	41.1	39.9	46.1	46.9	43.7	43.2	35.7	37.9	33.2	40.4	35.5	39.2	37.0
Organized Sports Outside of School	34.7	24.1	29.5	28.6	34.6	24.5	28.9	43.9	26.0	29.5	38.0	43.1	26.5	33.7
School Band	15.3	4.1	6.9	11.9	15.2	5.0	9.2	13.5	7.3	9.6	10.4	13.3	7.4	10.0
School Club(s)	25.4	44.3	45.2	24.9	28.4	41.3	35.6	18.5	32.5	35.0	18.1	19.6	31.1	26.4
Community Club(s)	6.6	5.6	8.0	4.3	6.9	5.4	6.1	11.0	13.5	16.4	8.7	11.2	13.1	12.4

**Table 16. Trends in delinquent behaviors for Columbia County youth, 2001, 2002, 2004, 2006 and 2008**

	2001			2002			2004			2006			2008		
	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>	<i>Middle School</i>	<i>High School</i>	<i>Total</i>
Carrying a handgun	3.9	9.9	6.9	2.7	3.9	3.3	3.4	3.2	3.3	5.3	3.7	4.3	6.1	6.0	6.0
Selling drugs	3.0	10.7	6.8	2.3	7.6	5.0	2.3	6.9	4.8	2.8	2.8	3.8	2.8	4.6	3.7
Attempting to steal a vehicle	2.5	4.2	3.3	2.5	3.4	2.9	1.5	1.8	1.6	2.5	2.5	2.4	2.8	1.7	2.3
Being arrested	5.3	7.2	6.2	3.9	6.1	5.0	4.9	6.8	5.9	4.0	0.4	3.0	5.4	4.9	5.2
Taking a handgun to school	0.8	1.3	1.1	0.7	1.8	1.2	0.8	1.0	0.9	1.2	0.0	0.5	1.2	0.7	0.9
Getting suspended	14.1	17.0	15.5	19.0	13.4	16.2	19.2	18.7	18.9	19.2	3.6	12.0	15.3	14.2	14.7
Attacking someone with intent to harm	11.4	18.4	14.9	11.3	12.2	11.9	13.6	13.2	13.4	9.8	5.9	8.5	11.1	10.3	10.6
Being drunk or high at school	8.3	23.7	15.9	5.8	18.8	12.5	10.1	18.3	14.6	8.5	21.6	16.6	9.0	15.8	12.6



**Table 17. Trends in early ATOD use and attitudes toward substance use for Columbia County youth, 2001, 2002, 2004, 2006 and 2008**

	2001			2002			2004			2006			2008		
	Middle School	High School	Total	Middle School	High School	Total	Middle School	High School	Total	Middle School	High School	Total	Middle School	High School	Total
<b><i>Early ATOD use (age 13 or younger)</i></b>															
More than a sip or two of alcohol	--	44.3	--	--	32.9	--	--	33.8	--	--	43.9	--	--	32.5	--
Drinking at least once a month	--	7.9	--	--	6.3	--	--	6.1	--	--	3.6	--	--	5.5	--
Cigarettes	--	39.7	--	--	33.0	--	--	41.7	--	--	26.0	--	--	25.7	--
Marijuana	--	17.8	--	--	14.0	--	--	19.1	--	--	4.9	--	--	11.6	--
<b><i>Perceive great risk of harm if...</i></b>															
One or more drinks every day	38.3	30.5	34.2	39.7	34.6	36.9	39.3	29.8	34.2	40.3	31.0	35.0	43.5	45.4	44.5
Smoke a pack or more every day	64.6	60.2	62.4	59.8	64.8	62.4	62.0	59.7	60.7	59.9	74.1	66.5	60.2	59.5	59.8
Smoke marijuana regularly	70.2	55.1	62.4	67.8	55.2	61.3	69.7	52.4	60.7	68.5	63.2	64.7	70.1	55.9	62.4
Try marijuana once or twice	41.2	33.2	37.2	39.3	29.5	34.2	41.0	26.5	33.4	43.2	33.6	37.4	48.7	31.9	39.6
<b><i>Think it wrong if...</i></b>															
Smoke cigarettes	84.8	63.9	74.1	85.7	67.6	76.4	84.0	66.4	74.7	87.8	72.0	78.9	85.0	72.6	78.3
Drink alcohol regularly	79.1	48.0	63.3	78.4	53.7	65.7	76.2	49.5	62.2	79.3	55.0	66.3	78.9	59.8	68.6
Smoke marijuana	86.7	71.8	79.1	89.0	74.4	81.4	89.4	74.5	81.5	89.4	86.5	87.2	85.5	76.8	80.8
Use other illicit drugs	96.5	92.6	94.5	97.9	95.9	96.8	97.9	94.6	96.0	97.9	96.0	96.1	95.3	95.2	95.3
<b><i>Seen as cool if...</i></b>															
Drink alcohol regularly	10.4	13.6	12.1	10.7	16.3	13.5	9.1	15.4	12.4	8.3	12.2	11.1	13.5	13.7	13.6
Smoke cigarettes	9.4	5.4	7.4	8.5	5.8	7.2	5.8	5.7	5.7	4.1	4.0	4.9	8.6	3.3	5.7
Smoke marijuana	14.4	12.3	13.5	11.3	10.1	10.7	8.1	11.6	9.9	8.1	7.0	8.3	11.1	9.7	10.4

Note: The symbol "--" indicates that data are not available.

**Table 18. Protective factor prevalence rates for Columbia County, Florida Statewide and the national normative database, 2008**

Domain	Scale	Columbia County		Florida Statewide		National Norms	
		<i>Middle School</i>	<i>High School</i>	<i>Middle School</i>	<i>High School</i>	<i>Middle School</i>	<i>High School</i>
<b>Community</b>	Community Opportunities for Prosocial Involvement	--	47	--	39	--	52
	Community Rewards for Prosocial Involvement	55	64	51	61	56	63
<b>Family</b>	Family Attachment	--	54	--	55	--	56
	Family Opportunities for Prosocial Involvement	60	54	56	53	59	54
	Family Rewards for Prosocial Involvement	52	57	50	54	54	55
<b>School</b>	School Opportunities for Prosocial Involvement	48	58	47	59	57	60
	School Rewards for Prosocial Involvement	40	55	45	56	53	58
<b>Peer and Individual</b>	Religiosity	54	65	51	61	56	62
	Social Skills	--	62	--	62	--	57
	Belief in the Moral Order	--	64	--	59	--	62
<b>Average Prevalence Rate</b>		<b>52</b>	<b>57</b>	<b>50</b>	<b>56</b>	<b>56</b>	<b>57</b>

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values. The symbol "--" indicates that data are not available.

**Table 19. Risk factor prevalence rates for Columbia County, Florida Statewide and the national normative database, 2008**

Domain	Scale	Columbia County		Florida Statewide		National Norms	
		Middle School	High School	Middle School	High School	Middle School	High School
<b>Community</b>	Low Neighborhood Attachment	--	53	--	49	--	44
	Community Disorganization	38	41	48	49	47	47
	Transitions and Mobility	60	60	61	64	47	46
	Laws and Norms Favorable to Drug Use	52	34	44	35	42	42
	Laws and Norms Favorable to Handguns	--	31	--	23	--	23
	Perceived Availability of Drugs	52	38	49	40	45	45
	Perceived Availability of Handguns	35	47	27	41	25	42
<b>Family</b>	Poor Family Management	55	48	49	49	44	45
	Family Conflict	37	29	43	37	42	37
	Family History of Antisocial Behavior	--	38	--	43	--	45
	Parental Attitudes Favorable toward ATOD Use	28	34	22	38	23	41
	Parental Attitudes Favorable toward Antisocial Behavior	--	41	--	45	--	48
<b>School</b>	Poor Academic Performance	50	48	45	44	45	48
	Lack of Commitment to School	55	49	55	47	47	46
<b>Peer and Individual</b>	Rebelliousness	--	36	--	41	--	40
	Friends' Delinquent Behavior	--	39	--	44	--	41
	Friends' Use of Drugs	--	31	--	39	--	47
	Peer Rewards for Antisocial Behavior	45	36	42	42	40	46
	Favorable Attitudes toward Antisocial Behavior	51	44	48	47	40	46
	Favorable Attitudes toward ATOD Use	43	38	40	40	39	45
	Low Perceived Risks of Drug Use	42	41	41	44	40	46
	Early Initiation of Drug Use	42	36	37	35	41	46
	Sensation Seeking	--	39	--	44	--	45
<b>Average Prevalence Rate</b>		<b>46</b>	<b>41</b>	<b>43</b>	<b>43</b>	<b>40</b>	<b>44</b>

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values. The symbol "--" indicates that data are not available.

**Table 20. Protective factor prevalence rate trends among middle school students for Columbia County, 2001, 2002, 2004, 2006 and 2008**

Domain	Scale	Columbia County				
		2001	2002	2004	2006	2008
<b>Community</b>	Community Opportunities for Prosocial Involvement	60	53	--	50	--
	Community Rewards for Prosocial Involvement	58	53	54	50	55
<b>Family</b>	Family Attachment	54	57	52	55	--
	Family Opportunities for Prosocial Involvement	54	56	56	58	60
	Family Rewards for Prosocial Involvement	56	56	51	51	52
<b>School</b>	School Opportunities for Prosocial Involvement	53	42	48	54	48
	School Rewards for Prosocial Involvement	45	42	48	46	40
<b>Peer and Individual</b>	Religiosity	57	61	58	54	54
	Social Skills	51	52	46	43	--
	Belief in the Moral Order	50	44	50	51	--
<b>Average Prevalence Rate</b>		<b>54</b>	<b>52</b>	<b>53</b>	<b>51</b>	<b>52</b>

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values. The symbol "--" indicates that data are not available.

**Table 21. Risk factor prevalence rate trends among middle school students for Columbia County, 2001, 2002, 2004, 2006 and 2008**

Domain	Scale	Columbia County				
		2001	2002	2004	2006	2008
<b>Community</b>	Low Neighborhood Attachment	49	51	45	48	--
	Community Disorganization	43	43	43	41	38
	Transitions and Mobility	74	63	64	58	60
	Laws and Norms Favorable to Drug Use	50	49	48	48	52
	Laws and Norms Favorable to Handguns	51	49	49	52	--
	Perceived Availability of Drugs	51	45	49	38	52
	Perceived Availability of Handguns	37	31	32	26	35
<b>Family</b>	Poor Family Management	49	46	52	53	55
	Family Conflict	43	43	39	32	37
	Family History of Antisocial Behavior	56	56	57	47	--
	Parental Attitudes Favorable toward ATOD Use	20	22	27	22	28
	Parental Attitudes Favorable toward Antisocial Behavior	34	33	36	37	--
<b>School</b>	Poor Academic Performance	47	56	60	53	50
	Lack of Commitment to School	47	53	50	50	55
<b>Peer and Individual</b>	Rebelliousness	47	46	45	43	--
	Friends' Delinquent Behavior	52	56	53	48	--
	Friends' Use of Drugs	48	47	47	41	--
	Peer Rewards for Antisocial Behavior	46	44	40	37	45
	Favorable Attitudes toward Antisocial Behavior	38	41	45	42	51
	Favorable Attitudes toward ATOD Use	46	49	47	40	43
	Low Perceived Risks of Drug Use	46	45	47	46	42
	Early Initiation of Drug Use	49	55	50	42	42
	Sensation Seeking	45	46	54	46	--
<b>Average Prevalence Rate</b>		<b>47</b>	<b>46</b>	<b>47</b>	<b>43</b>	<b>46</b>

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values. The symbol "--" indicates that data are not available.

**Table 22. Protective factor prevalence rate trends among high school students for Columbia County, 2001, 2002, 2004, 2006 and 2008**

Domain	Scale	Columbia County				
		2001	2002	2004	2006	2008
<b>Community</b>	Community Opportunities for Prosocial Involvement	53	57	--	40	47
	Community Rewards for Prosocial Involvement	68	68	60	64	64
<b>Family</b>	Family Attachment	57	57	53	48	54
	Family Opportunities for Prosocial Involvement	57	56	54	44	54
	Family Rewards for Prosocial Involvement	56	58	53	45	57
<b>School</b>	School Opportunities for Prosocial Involvement	49	56	52	52	58
	School Rewards for Prosocial Involvement	38	48	48	51	55
<b>Peer and Individual</b>	Religiosity	67	72	63	63	65
	Social Skills	56	60	57	57	62
	Belief in the Moral Order	61	62	61	67	64
<b>Average Prevalence Rate</b>		<b>56</b>	<b>59</b>	<b>55</b>	<b>52</b>	<b>57</b>

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values. The symbol "--" indicates that data are not available.

**Table 23. Risk factor prevalence rate trends among high school students for Columbia County, 2001, 2002, 2004, 2006 and 2008**

Domain	Scale	Columbia County				
		2001	2002	2004	2006	2008
<b>Community</b>	Low Neighborhood Attachment	42	48	49	42	53
	Community Disorganization	38	41	45	31	41
	Transitions and Mobility	67	63	64	58	60
	Laws and Norms Favorable to Drug Use	40	40	43	33	34
	Laws and Norms Favorable to Handguns	31	35	33	38	31
	Perceived Availability of Drugs	49	48	44	50	38
	Perceived Availability of Handguns	59	50	49	62	47
<b>Family</b>	Poor Family Management	51	52	49	59	48
	Family Conflict	38	32	36	42	29
	Family History of Antisocial Behavior	53	49	49	39	38
	Parental Attitudes Favorable toward ATOD Use	43	37	46	43	34
	Parental Attitudes Favorable toward Antisocial Behavior	32	40	44	54	41
<b>School</b>	Poor Academic Performance	51	44	52	45	48
	Lack of Commitment to School	56	45	53	65	49
<b>Peer and Individual</b>	Rebelliousness	37	39	41	41	36
	Friends' Delinquent Behavior	51	40	44	34	39
	Friends' Use of Drugs	55	44	43	33	31
	Peer Rewards for Antisocial Behavior	38	42	45	43	36
	Favorable Attitudes toward Antisocial Behavior	44	40	45	40	44
	Favorable Attitudes toward ATOD Use	42	43	45	36	38
	Low Perceived Risks of Drug Use	45	44	47	34	41
	Early Initiation of Drug Use	53	44	49	45	36
	Sensation Seeking	48	44	46	41	39
<b>Average Prevalence Rate</b>		<b>47</b>	<b>44</b>	<b>47</b>	<b>45</b>	<b>41</b>

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.





# Appendix B

## References


- Arthur, M. W., Briney, J. S., Hawkins, J. D., Abbott, R. D., Brooke-Weiss, B. L. & Catalano, R. F. (2007). Measuring risk and protection in communities using the Communities That Care Youth Survey. *Evaluation and Program Planning, 30*, 197-211.
- Arthur, M. W., Hawkins, J. D., Pollard, J. A., Catalano, R. F. & Baglioni, A. J. (2002). Measuring risk and protective factors for substance use, delinquency, and other adolescent problem behaviors: The communities that care youth survey. *Evaluation Review, 26*, 575-601.
- Bachman, J. G., Johnston, L. D. & O'Malley, P. M. (1996). *The Monitoring the Future project after twenty-two years: Design and procedures*. (Monitoring the Future Occasional Paper No. 38.) Ann Arbor, MI: Institute for Social Research.
- Bachman, J. G., Johnston, L. D., O'Malley, P. M. & Humphrey, R. H. (1986). *Changes in marijuana use linked to changes in perceived risks and disapproval*. (Monitoring the Future Occasional Paper No. 19.) Ann Arbor, MI: Institute for Social Research.
- Blum, R. W., Beuhring, T., Shew, M. L., Bearinger, L. H., Sieving, R. E. & Resnick, M. D. (2000). The effects of race/ethnicity, income, and family structure on adolescent risk behaviors. *American Journal of Public Health, 90*, 1879-1884.
- Bracht, N. & Kingsbury, L. (1990). Community organization principles in health promotion: A five-state model. In N. Bracht (Ed.), *Health promotion at the community level* (pp. 66-88). Beverly Hills, CA: Sage.
- Bry, B. H., McKeon, P. & Pandina, R. J. (1982). Extent of drug use as a function of number of risk factors. *Journal of Abnormal Psychology, 91*, 273-279.
- Catalano, R. F. & Hawkins, J. D. (1996). The social development model: A theory of antisocial behavior. In J. D. Hawkins (Ed.), *Delinquency and crime: Current theories* (pp. 149-197). New York, NY: Cambridge University Press.
- Hawkins, J. D., Arthur, M. W. & Catalano, R. F. (1995). Preventing substance abuse. In M. Tonry & D. Farrington (Eds.), *Building a safer society: Strategic approaches to crime prevention* (Vol. 19, pp. 343-427, Crime and justice: A review of research). Chicago, IL: University of Chicago Press.
- Hawkins, J. D., Catalano, R. F. & Associates. (1992). *Communities that care: Action for drug abuse prevention* (1<sup>st</sup> ed.). San Francisco: Jossey-Bass.
- Hawkins, J. D., Catalano, R. F. & Miller, J. Y. (1992). Risk and protective factors for alcohol and other drug problems in adolescence and early adulthood: Implications for substance abuse prevention. *Psychological Bulletin, 112*, 64-105.
- Johnston, L. D., O'Malley, P. M., Bachman, J. G. & Schulenberg, J. E. (2008). *Monitoring the Future national survey results on drug use, 1975-2007. Volume I: Secondary school students* (NIH Publication No. 08-6418A). Bethesda, MD: National Institute on Drug Abuse.

- .....
- Newcomb, M. D. (1995). Identifying high-risk youth: Prevalence and patterns of adolescent drug abuse. In E. Rahdert & D. Czechowicz (Eds.), *Adolescent drug abuse: Clinical assessment and therapeutic interventions* (NIDA Research Monograph, 156). Washington, DC: U.S. Department of Health and Human Services.
- Newcomb, M. D. & Felix-Ortiz, M. (1992). Multiple protective and risk factors for drug use and abuse: Cross-sectional and prospective findings. *Journal of Personality and Social Psychology*, 51, 564-577.
- Newcomb, M. D., Maddahian, E. & Skager, R. (1987). Substance abuse and psychosocial risk factors among teenagers: Associations with sex, age, ethnicity, and type of school. *American Journal of Drug and Alcohol Abuse*, 13, 413-433.
- Pollard, J. A., Hawkins, J. D. & Arthur, M. W. (1999). Risk and protection: Are both necessary to understand diverse behavioral outcomes in adolescence? *Social Work Research*, 23, 145-158.

# Appendix C

## The Social Development Strategy

.....



## Communities That Care®

### Risk Factors

### Adolescent Problem Behaviors

Community	Substance Abuse	Delinquency	Teen Pregnancy	School Drop-Out	Violence
Availability of drugs	●				●
Availability of firearms		●			●
Community laws and norms favorable toward drug use, firearms and crime	●	●			●
Media portrayals of violence					●
Transitions and mobility	●	●		●	
Low neighborhood attachment and community disorganization	●	●			●
Extreme economic deprivation	●	●	●	●	●
<b>Family</b>					
Family history of the problem behavior	●	●	●	●	●
Family management problems	●	●	●	●	●
Family conflict	●	●	●	●	●
Favorable parental attitudes and involvement in the problem behavior	●	●			●
<b>School</b>					
Academic failure beginning in late elementary school	●	●	●	●	●
Lack of commitment to school	●	●	●	●	●
<b>Peer and Individual</b>					
Early and persistent antisocial behavior	●	●	●	●	●
Rebelliousness	●	●		●	
Friends who engage in the problem behavior	●	●	●	●	●
Gang involvement	●	●			●
Favorable attitudes toward the problem behavior	●	●	●	●	
Early initiation of the problem behavior	●	●	●	●	●
Constitutional factors	●	●			●

# Appendix D

## Other Resources

---

### Web Sites

Office of National Drug Control Policy [www.whitehousedrugpolicy.gov](http://www.whitehousedrugpolicy.gov)

National Clearinghouse for Alcohol and Drug Information [www.health.org/index.htm](http://www.health.org/index.htm)

Substance Abuse and Mental Health Services Administration (SAMHSA) [www.samhsa.gov](http://www.samhsa.gov)

Monitoring the Future [www.monitoringthefuture.org](http://www.monitoringthefuture.org)

National Institute on Drug Abuse (NIDA) [www.nida.nih.gov](http://www.nida.nih.gov) and [www.drugabuse.gov](http://www.drugabuse.gov)

National Institute on Alcohol Abuse and Alcoholism (NIAAA) [www.niaaa.nih.gov](http://www.niaaa.nih.gov)

Social Development Research Group <http://depts.washington.edu/sdrg>

### Prevention Program Guides

Center for Substance Abuse Prevention, Western Center for the Application of Prevention Technologies. (2006). *Building a successful prevention program: list of all practices*. [Data file]. Available at the University of Nevada Reno's Web site, <http://casat.unr.edu/bestpractices/alpha-list.php>.

Center for the Study and Prevention of Violence, Institute of Behavioral Science. (2006). *Blueprints for Violence Prevention*. [Data file]. Available from the University of Colorado Boulder's Web site, [www.colorado.edu/cspv/blueprints](http://www.colorado.edu/cspv/blueprints).

U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration (SAMHSA). (2006). *Communities That Care Prevention Strategies Guide*. [Datafile]. Available from the SAMHSA Web site, <http://preventionplatform.samhsa.gov>.

U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration (SAMHSA). (2006). *Model Programs list*. [Data file]. Available from the SAMHSA Web site, <http://modelprograms.samhsa.gov>.

### Prevention Planning

Hawkins, J. D., Catalano, R. F., & Associates. (1992). *Communities that care: Action for drug abuse prevention* (1<sup>st</sup> ed.). San Francisco: Jossey-Bass.