

2018 Florida Youth Substance Abuse Survey

Big Bend Community Based Care Data Tables

Table 1. Major demographic characteristics of surveyed Big Bend Community Based Care and Florida Statewide, 2018

| | Big Bend Community Based Care | | | Florida Statewide | |
|----------------------------------|-------------------------------|--------------|--|-------------------|--------------|
| | N | % | | N | % |
| Sex | | | | | |
| Female | 5,971 | 50.6 | | 26,340 | 48.2 |
| Male | 5,656 | 47.9 | | 27,468 | 50.3 |
| Race/Ethnic group | | | | | |
| African American | 2,101 | 17.8 | | 555 | 1.0 |
| American Indian | 345 | 2.9 | | 659 | 1.2 |
| Asian | 206 | 1.7 | | 12,088 | 22.1 |
| Hispanic/Latino | 665 | 5.6 | | 11,242 | 20.6 |
| Native Hawaiian/Pacific Islander | 37 | 0.3 | | 79 | 0.1 |
| Other/Multiple | 2,048 | 17.3 | | 6,899 | 12.6 |
| White, non-Hispanic | 6,280 | 53.2 | | 22,618 | 41.4 |
| Age | | | | | |
| 10 | 6 | 0.1 | | 50 | 0.1 |
| 11 | 863 | 7.3 | | 3,339 | 6.1 |
| 12 | 1,990 | 16.9 | | 7,363 | 13.5 |
| 13 | 1,965 | 16.6 | | 7,738 | 14.2 |
| 14 | 1,920 | 16.3 | | 7,864 | 14.4 |
| 15 | 1,699 | 14.4 | | 7,982 | 14.6 |
| 16 | 1,465 | 12.4 | | 7,926 | 14.5 |
| 17 | 1,144 | 9.7 | | 7,725 | 14.1 |
| 18 | 630 | 5.3 | | 3,990 | 7.3 |
| 19 or older | 92 | 0.8 | | 495 | 0.9 |
| Grade | | | | | |
| 6th | 2,302 | 19.5 | | 8,050 | 14.7 |
| 7th | 1,954 | 16.5 | | 7,706 | 14.1 |
| 8th | 2,046 | 17.3 | | 7,715 | 14.1 |
| 9th | 1,676 | 14.2 | | 8,024 | 14.7 |
| 10th | 1,592 | 13.5 | | 7,925 | 14.5 |
| 11th | 1,204 | 10.2 | | 7,775 | 14.2 |
| 12th | 1,033 | 8.7 | | 7,417 | 13.6 |
| Overall Middle School | 6,302 | 53.4 | | 23,470 | 43.0 |
| Overall High School | 5,505 | 46.6 | | 31,141 | 57.0 |
| Total | 11,807 | 100.0 | | 54,611 | 100.0 |

Note: Some categories do not sum to 100% of the total due to missing values (e.g., not all survey questions were answered). In addition, rounding can produce totals that do not equal 100%. "N" represents the number of valid cases. In this table, county data are unweighted while statewide data are weighted.

Table 2. Percentages of Big Bend Community Based Care and Florida Statewide youth who reported having used various drugs in their lifetimes, 2018

| | Big Bend Community Based Care | | | | | | | Florida Statewide | | | | | | |
|---------------------------------------|-------------------------------|--------------------|---------------|-------------|-------------------|-------------------|--------------|----------------------|--------------------|---------------|-------------|-------------------|-------------------|--------------|
| | <i>Middle School</i> | <i>High School</i> | <i>Female</i> | <i>Male</i> | <i>Ages 10-14</i> | <i>Ages 15-17</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Female</i> | <i>Male</i> | <i>Ages 10-14</i> | <i>Ages 15-17</i> | <i>Total</i> |
| Alcohol | 22.1 | 47.6 | 38.0 | 34.6 | 23.6 | 47.5 | 36.4 | 20.8 | 48.0 | 38.9 | 34.1 | 22.9 | 48.1 | 36.5 |
| Blacking Out | -- | 16.4 | 17.0 | 15.9 | -- | 16.8 | -- | -- | 14.2 | 15.0 | 13.5 | -- | 14.0 | -- |
| Cigarettes | 10.0 | 20.2 | 15.3 | 15.9 | 9.7 | 19.8 | 15.6 | 6.7 | 14.7 | 11.1 | 11.4 | 6.7 | 14.7 | 11.3 |
| Vaporizer / E-Cigarette | 15.3 | 38.4 | 28.1 | 28.2 | 16.4 | 39.0 | 28.1 | 15.1 | 36.0 | 26.5 | 27.6 | 16.7 | 36.4 | 27.1 |
| Marijuana or Hashish | 7.8 | 29.9 | 21.1 | 19.0 | 8.3 | 30.8 | 20.0 | 7.6 | 29.7 | 21.0 | 19.4 | 8.7 | 29.8 | 20.2 |
| Synthetic Marijuana | -- | 3.4 | 3.9 | 2.9 | -- | 3.8 | -- | -- | 3.5 | 3.7 | 3.3 | -- | 3.5 | -- |
| Inhalants | 8.0 | 5.2 | 7.3 | 5.5 | 7.8 | 5.5 | 6.4 | 7.8 | 4.3 | 6.7 | 4.9 | 7.6 | 4.3 | 5.8 |
| Flakka | -- | 0.9 | 0.8 | 1.1 | -- | 1.1 | -- | -- | 0.8 | 0.7 | 0.8 | -- | 0.7 | -- |
| Club Drugs | 0.6 | 2.4 | 1.5 | 1.6 | 0.7 | 2.3 | 1.6 | 0.7 | 1.8 | 1.2 | 1.4 | 0.7 | 1.7 | 1.3 |
| LSD, PCP or Mushrooms | 0.7 | 4.4 | 2.2 | 3.3 | 0.8 | 4.4 | 2.7 | 1.0 | 4.0 | 2.3 | 3.1 | 1.1 | 3.9 | 2.7 |
| Methamphetamine | 0.5 | 1.0 | 0.7 | 0.8 | 0.4 | 1.1 | 0.8 | 0.7 | 0.7 | 0.6 | 0.8 | 0.7 | 0.6 | 0.7 |
| Cocaine or Crack Cocaine | 0.7 | 2.5 | 1.8 | 1.6 | 0.7 | 2.4 | 1.7 | 0.8 | 1.9 | 1.3 | 1.6 | 0.8 | 1.7 | 1.4 |
| Heroin | 0.3 | 0.5 | 0.4 | 0.4 | 0.3 | 0.6 | 0.4 | 0.4 | 0.3 | 0.3 | 0.4 | 0.4 | 0.3 | 0.4 |
| Prescription Depressants | 2.9 | 8.0 | 5.8 | 5.4 | 2.9 | 8.0 | 5.7 | 2.3 | 5.9 | 4.4 | 4.3 | 2.5 | 5.9 | 4.4 |
| Prescription Pain Relievers | 2.6 | 6.4 | 5.1 | 4.3 | 3.0 | 6.1 | 4.7 | 2.6 | 5.0 | 4.4 | 3.6 | 2.8 | 4.9 | 4.0 |
| Prescription Amphetamines | 1.5 | 5.7 | 4.1 | 3.6 | 1.6 | 5.4 | 3.8 | 1.2 | 3.6 | 2.6 | 2.6 | 1.4 | 3.4 | 2.6 |
| Steroids (without a doctor's order) | 0.9 | 0.5 | 0.6 | 0.8 | 0.9 | 0.5 | 0.7 | 0.7 | 0.4 | 0.5 | 0.6 | 0.7 | 0.4 | 0.5 |
| Over-the-Counter Drugs | 3.5 | 6.2 | 5.0 | 4.8 | 3.5 | 6.8 | 5.0 | 3.2 | 4.9 | 4.3 | 4.1 | 3.4 | 4.9 | 4.2 |
| Needle to Inject Illegal Drugs | -- | 0.8 | 0.7 | 0.8 | -- | 0.7 | -- | -- | 0.6 | 0.5 | 0.8 | -- | 0.6 | -- |
| Any illicit drug | 17.1 | 36.2 | 29.5 | 25.8 | 17.6 | 37.4 | 27.7 | 16.5 | 34.8 | 28.7 | 25.2 | 17.4 | 34.9 | 26.9 |
| Any illicit drug other than marijuana | 13.2 | 19.1 | 17.3 | 15.6 | 13.3 | 19.7 | 16.5 | 12.6 | 16.2 | 15.5 | 13.7 | 12.8 | 16.2 | 14.6 |
| Alcohol only | 11.4 | 18.1 | 15.3 | 15.0 | 12.5 | 17.0 | 15.2 | 11.3 | 19.8 | 16.8 | 15.6 | 12.4 | 19.7 | 16.2 |
| Alcohol or any illicit drug | 28.1 | 54.2 | 44.4 | 40.6 | 29.7 | 54.3 | 42.6 | 27.5 | 54.4 | 45.2 | 40.5 | 29.5 | 54.5 | 42.9 |
| Any illicit drug, but no alcohol | 6.3 | 6.7 | 6.8 | 6.2 | 6.3 | 6.9 | 6.5 | 6.9 | 6.6 | 6.7 | 6.7 | 6.9 | 6.7 | 6.7 |

Note: The first 19 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. For "Blacking Out," respondents were asked on how many occasions in their lifetime they woke up after a night of drinking and did not remember the things they did or the places they went. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "--" indicates that data are not available.

Table 3. Percentages of Big Bend Community Based Care and Florida Statewide youth who reported having used various drugs in the past 30 days, 2018

| | Big Bend Community Based Care | | | | | | | Florida Statewide | | | | | | |
|---------------------------------------|-------------------------------|--------------------|---------------|-------------|-------------------|-------------------|--------------|----------------------|--------------------|---------------|-------------|-------------------|-------------------|--------------|
| | <i>Middle School</i> | <i>High School</i> | <i>Female</i> | <i>Male</i> | <i>Ages 10-14</i> | <i>Ages 15-17</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Female</i> | <i>Male</i> | <i>Ages 10-14</i> | <i>Ages 15-17</i> | <i>Total</i> |
| Alcohol | 8.5 | 23.5 | 18.3 | 15.4 | 9.1 | 22.9 | 16.9 | 7.3 | 21.2 | 16.8 | 13.8 | 8.1 | 20.9 | 15.3 |
| Binge Drinking | 2.8 | 11.5 | 7.2 | 8.0 | 2.9 | 11.1 | 7.6 | 3.1 | 9.6 | 6.7 | 6.9 | 3.2 | 9.4 | 6.8 |
| Cigarettes | 1.7 | 5.6 | 3.7 | 4.0 | 1.7 | 5.7 | 3.9 | 1.2 | 3.5 | 2.6 | 2.4 | 1.2 | 3.3 | 2.5 |
| Vaporizer / E-Cigarette | 6.0 | 21.9 | 14.8 | 14.9 | 6.9 | 22.1 | 14.8 | 6.4 | 19.2 | 13.5 | 14.0 | 7.5 | 19.1 | 13.7 |
| Marijuana or Hashish | 3.5 | 15.1 | 10.4 | 9.5 | 3.8 | 15.4 | 10.0 | 3.7 | 16.3 | 11.0 | 10.7 | 4.2 | 16.2 | 10.9 |
| Synthetic Marijuana | -- | 1.2 | 1.1 | 1.1 | -- | 1.4 | -- | -- | 1.1 | 1.1 | 1.1 | -- | 1.1 | -- |
| Inhalants | 3.0 | 1.3 | 2.3 | 1.8 | 2.8 | 1.5 | 2.1 | 2.8 | 1.1 | 2.1 | 1.6 | 2.6 | 1.1 | 1.8 |
| Flakka | -- | 0.4 | 0.2 | 0.6 | -- | 0.5 | -- | -- | 0.4 | 0.3 | 0.4 | -- | 0.3 | -- |
| Club Drugs | 0.3 | 0.7 | 0.4 | 0.6 | 0.2 | 0.8 | 0.5 | 0.2 | 0.5 | 0.3 | 0.5 | 0.2 | 0.5 | 0.4 |
| LSD, PCP or Mushrooms | 0.4 | 1.3 | 0.5 | 1.3 | 0.4 | 1.5 | 0.9 | 0.4 | 1.1 | 0.7 | 0.9 | 0.4 | 1.2 | 0.8 |
| Methamphetamine | 0.2 | 0.4 | 0.2 | 0.4 | 0.2 | 0.4 | 0.3 | 0.4 | 0.4 | 0.3 | 0.5 | 0.4 | 0.4 | 0.4 |
| Cocaine or Crack Cocaine | 0.3 | 0.7 | 0.6 | 0.5 | 0.3 | 0.6 | 0.5 | 0.3 | 0.5 | 0.4 | 0.5 | 0.3 | 0.4 | 0.4 |
| Heroin | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.2 | 0.1 | 0.1 | 0.1 |
| Prescription Depressants | 0.9 | 2.4 | 1.7 | 1.7 | 0.9 | 2.4 | 1.7 | 0.9 | 1.6 | 1.4 | 1.2 | 0.9 | 1.6 | 1.3 |
| Prescription Pain Relievers | 0.9 | 1.9 | 1.7 | 1.3 | 1.0 | 1.8 | 1.5 | 1.0 | 1.4 | 1.5 | 0.9 | 1.0 | 1.5 | 1.2 |
| Prescription Amphetamines | 0.6 | 1.6 | 1.1 | 1.3 | 0.6 | 1.7 | 1.2 | 0.5 | 1.0 | 0.8 | 0.7 | 0.5 | 1.0 | 0.8 |
| Steroids (without a doctor's order) | 0.1 | 0.1 | 0.1 | 0.1 | 0.2 | 0.1 | 0.1 | 0.2 | 0.1 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 |
| Over-the-Counter Drugs | 1.5 | 2.2 | 1.7 | 2.0 | 1.5 | 2.5 | 1.9 | 1.4 | 1.8 | 1.7 | 1.5 | 1.5 | 1.8 | 1.6 |
| Any illicit drug | 7.8 | 18.4 | 14.6 | 12.8 | 8.0 | 18.9 | 13.7 | 7.9 | 19.0 | 15.0 | 13.4 | 8.3 | 19.1 | 14.3 |
| Any illicit drug other than marijuana | 5.7 | 7.3 | 6.8 | 6.3 | 5.7 | 7.7 | 6.6 | 5.5 | 6.0 | 6.5 | 5.1 | 5.5 | 6.1 | 5.8 |
| Alcohol only | 5.3 | 12.7 | 10.5 | 8.3 | 5.8 | 11.7 | 9.4 | 4.3 | 11.1 | 9.0 | 7.4 | 4.8 | 10.9 | 8.2 |
| Alcohol or any illicit drug | 12.9 | 30.3 | 24.5 | 20.6 | 13.5 | 29.9 | 22.6 | 12.1 | 29.5 | 23.7 | 20.4 | 12.9 | 29.4 | 22.0 |
| Any illicit drug, but no alcohol | 4.6 | 7.1 | 6.5 | 5.4 | 4.6 | 7.3 | 6.0 | 4.9 | 8.6 | 7.2 | 6.9 | 5.1 | 8.8 | 7.1 |

Note: The first 18 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. Binge drinking is defined as having had five or more alcoholic drinks in a row in the past two weeks. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "--" indicates that data are not available.

Table 4. Lifetime trend in alcohol, tobacco and other drug use for Big Bend Community Based Care, 2008 to 2018

| | 2008 | | | 2010 | | | 2012 | | | 2014 | | | 2016 | | | 2018 | | |
|---------------------------------------|----------------------|--------------------|--------------|----------------------|--------------------|--------------|----------------------|--------------------|--------------|----------------------|--------------------|--------------|----------------------|--------------------|--------------|----------------------|--------------------|--------------|
| | <i>Middle School</i> | <i>High School</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Total</i> |
| Alcohol | 38.6 | 65.3 | 53.6 | 35.4 | 62.5 | 50.6 | 28.3 | 59.6 | 45.7 | 23.4 | 54.3 | 40.6 | 22.1 | 52.5 | 39.3 | 22.1 | 47.6 | 36.4 |
| Blacking Out | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 20.7 | -- | -- | 19.0 | -- | -- | 16.4 | -- |
| Cigarettes | 25.3 | 41.0 | 34.1 | 21.6 | 37.4 | 30.5 | 16.3 | 33.4 | 25.8 | 12.5 | 29.3 | 21.8 | 10.4 | 26.3 | 19.3 | 10.0 | 20.2 | 15.6 |
| Vaporizer / E-Cigarette | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 13.8 | 37.8 | 27.2 | 15.3 | 38.4 | 28.1 |
| Marijuana or Hashish | 12.0 | 33.5 | 24.2 | 11.2 | 34.8 | 24.5 | 8.0 | 33.9 | 22.5 | 8.1 | 32.5 | 21.6 | 7.1 | 32.0 | 21.0 | 7.8 | 29.9 | 20.0 |
| Synthetic Marijuana | -- | -- | -- | -- | -- | -- | -- | 16.0 | -- | -- | 9.3 | -- | -- | 6.7 | -- | -- | 3.4 | -- |
| Inhalants | 13.5 | 10.9 | 12.1 | 13.6 | 9.0 | 11.0 | 10.7 | 7.8 | 9.0 | 8.2 | 5.8 | 6.8 | 6.8 | 4.9 | 5.8 | 8.0 | 5.2 | 6.4 |
| Flakka | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 1.0 | -- | -- | 0.9 | -- |
| Club Drugs | 2.0 | -- | -- | 1.7 | 4.9 | 3.5 | 1.1 | 4.0 | 2.7 | 0.8 | 4.2 | 2.7 | 0.9 | 3.5 | 2.4 | 0.6 | 2.4 | 1.6 |
| LSD, PCP or Mushrooms | 2.3 | | -- | 2.3 | 5.6 | 4.2 | 1.3 | 5.2 | 3.5 | 1.1 | 5.7 | 3.7 | 1.0 | 5.8 | 3.7 | 0.7 | 4.4 | 2.7 |
| Methamphetamine | 1.1 | 1.4 | 1.3 | 1.2 | 1.6 | 1.4 | 1.0 | 1.4 | 1.2 | 0.7 | 1.3 | 1.0 | 0.6 | 1.0 | 0.9 | 0.5 | 1.0 | 0.8 |
| Cocaine or Crack | 1.9 | -- | -- | 1.9 | 3.8 | 3.0 | 1.0 | 3.9 | 2.7 | 0.8 | 2.7 | 1.9 | 0.9 | 3.3 | 2.2 | 0.7 | 2.5 | 1.7 |
| Heroin | 0.9 | 0.9 | 0.9 | 1.0 | 1.2 | 1.1 | 0.7 | 0.9 | 0.8 | 0.5 | 0.9 | 0.7 | 0.4 | 0.4 | 0.4 | 0.3 | 0.5 | 0.4 |
| Prescription Depressants | 3.6 | 10.7 | 7.6 | 3.2 | 10.0 | 7.0 | 2.5 | 8.2 | 5.7 | 2.0 | 8.3 | 5.5 | 2.7 | 9.8 | 6.7 | 2.9 | 8.0 | 5.7 |
| Prescription Pain Relievers | 6.6 | 13.1 | 10.2 | 5.4 | 12.2 | 9.3 | 4.0 | 10.8 | 7.8 | 3.3 | 9.8 | 6.9 | 3.5 | 8.3 | 6.2 | 2.6 | 6.4 | 4.7 |
| Prescription Amphetamines | 3.1 | 8.9 | 6.4 | 2.4 | 9.6 | 6.5 | 2.0 | 8.1 | 5.4 | 1.3 | 7.7 | 4.9 | 1.8 | 7.4 | 4.9 | 1.5 | 5.7 | 3.8 |
| Steroids (without a doctor's order) | 1.3 | 1.5 | 1.4 | 1.0 | 1.2 | 1.1 | 0.6 | 1.2 | 0.9 | 0.6 | 0.8 | 0.7 | 0.6 | 0.7 | 0.7 | 0.9 | 0.5 | 0.7 |
| Over-the-Counter Drugs | 5.7 | -- | -- | 5.8 | 9.4 | 7.8 | 4.2 | 7.6 | 6.1 | 3.6 | 6.5 | 5.2 | 3.2 | 6.5 | 5.0 | 3.5 | 6.2 | 5.0 |
| Needle to Inject Drugs | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 0.9 | -- | -- | 0.8 | -- |
| Any illicit drug | 24.4 | 41.2 | 33.8 | 25.0 | 41.8 | 34.5 | 19.5 | 41.8 | 31.9 | 17.7 | 39.6 | 29.8 | 16.0 | 37.3 | 27.9 | 17.1 | 36.2 | 27.7 |
| Any illicit drug other than marijuana | 19.3 | 26.9 | 23.6 | 20.4 | 26.2 | 23.7 | 16.0 | 24.1 | 20.6 | 13.6 | 22.8 | 18.7 | 12.6 | 21.5 | 17.6 | 13.2 | 19.1 | 16.5 |
| Alcohol only | 20.5 | 27.8 | 24.6 | 17.7 | 25.1 | 21.8 | 15.2 | 22.5 | 19.3 | 12.7 | 20.3 | 16.9 | 12.4 | 20.4 | 16.9 | 11.4 | 18.1 | 15.2 |
| Alcohol or any illicit drug | 44.9 | 68.9 | 58.4 | 42.7 | 66.8 | 56.2 | 34.5 | 64.1 | 51.0 | 30.3 | 59.6 | 46.6 | 28.0 | 57.6 | 44.5 | 28.1 | 54.2 | 42.6 |
| Any illicit drug, but no alcohol | 6.5 | 3.8 | 4.9 | 7.4 | 4.4 | 5.7 | 6.4 | 4.6 | 5.4 | 7.0 | 5.4 | 6.1 | 6.5 | 5.3 | 5.8 | 6.3 | 6.7 | 6.5 |

Note: The first 19 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. For "Blacking Out," respondents were asked on how many occasions in their lifetime they woke up after a night of drinking and did not remember the things they did or the places they went. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "--" indicates that data are not available.

Table 5. Past-30-day trend in alcohol, tobacco and other drug use for Big Bend Community Based Care, 2008 to 2018

| | 2008 | | | 2010 | | | 2012 | | | 2014 | | | 2016 | | | 2018 | | |
|---------------------------------------|----------------------|--------------------|--------------|----------------------|--------------------|--------------|----------------------|--------------------|--------------|----------------------|--------------------|--------------|----------------------|--------------------|--------------|----------------------|--------------------|--------------|
| | <i>Middle School</i> | <i>High School</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Total</i> |
| Alcohol | 18.9 | 39.6 | 30.5 | 16.5 | 37.6 | 28.4 | 11.5 | 34.1 | 24.1 | 9.9 | 29.2 | 20.6 | 8.2 | 27.6 | 19.2 | 8.5 | 23.5 | 16.9 |
| Binge Drinking | 7.8 | 22.1 | 15.8 | 7.2 | 20.8 | 14.9 | 4.5 | 17.8 | 11.9 | 3.8 | 15.1 | 10.0 | 2.9 | 13.5 | 8.9 | 2.8 | 11.5 | 7.6 |
| Cigarettes | 7.3 | 16.8 | 12.7 | 6.1 | 15.5 | 11.4 | 4.4 | 12.9 | 9.1 | 3.2 | 9.0 | 6.4 | 2.4 | 7.5 | 5.2 | 1.7 | 5.6 | 3.9 |
| Vaporizer / E-Cigarette | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 4.7 | 14.9 | 10.4 | 6.0 | 21.9 | 14.8 |
| Marijuana or Hashish | 6.3 | 17.7 | 12.8 | 5.9 | 19.2 | 13.4 | 3.8 | 17.6 | 11.6 | 4.1 | 16.8 | 11.1 | 3.2 | 16.0 | 10.4 | 3.5 | 15.1 | 10.0 |
| Synthetic Marijuana | -- | -- | -- | -- | -- | -- | -- | 4.8 | -- | -- | 1.6 | -- | -- | 1.3 | -- | -- | 1.2 | -- |
| Inhalants | 5.5 | 2.5 | 3.8 | 5.6 | 2.4 | 3.8 | 3.9 | 2.0 | 2.8 | 2.8 | 1.5 | 2.1 | 2.3 | 1.1 | 1.6 | 3.0 | 1.3 | 2.1 |
| Flakka | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 0.4 | -- | -- | 0.4 | -- |
| Club Drugs | 0.8 | -- | -- | 0.5 | 1.4 | 1.0 | 0.3 | 1.3 | 0.9 | 0.4 | 1.1 | 0.8 | 0.3 | 0.9 | 0.6 | 0.3 | 0.7 | 0.5 |
| LSD, PCP or Mushrooms | 1.0 | -- | -- | 0.9 | 1.3 | 1.1 | 0.5 | 1.4 | 1.0 | 0.5 | 1.5 | 1.1 | 0.3 | 1.4 | 0.9 | 0.4 | 1.3 | 0.9 |
| Methamphetamine | 0.6 | 0.4 | 0.5 | 0.7 | 0.6 | 0.7 | 0.3 | 0.6 | 0.5 | 0.4 | 0.5 | 0.5 | 0.3 | 0.4 | 0.4 | 0.2 | 0.4 | 0.3 |
| Cocaine or Crack | 0.6 | -- | -- | 0.8 | 0.9 | 0.9 | 0.4 | 1.0 | 0.7 | 0.4 | 0.9 | 0.7 | 0.3 | 1.0 | 0.7 | 0.3 | 0.7 | 0.5 |
| Heroin | 0.3 | 0.2 | 0.3 | 0.4 | 0.4 | 0.4 | 0.2 | 0.5 | 0.4 | 0.2 | 0.2 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| Prescription Depressants | 1.5 | 4.1 | 3.0 | 1.5 | 3.4 | 2.6 | 0.9 | 3.2 | 2.2 | 0.7 | 2.9 | 1.9 | 1.0 | 3.8 | 2.6 | 0.9 | 2.4 | 1.7 |
| Prescription Pain Relievers | 3.0 | 4.6 | 3.9 | 2.9 | 5.1 | 4.1 | 1.6 | 3.7 | 2.8 | 1.6 | 3.7 | 2.8 | 1.5 | 3.0 | 2.4 | 0.9 | 1.9 | 1.5 |
| Prescription Amphetamines | 1.2 | 2.9 | 2.1 | 1.2 | 3.4 | 2.5 | 0.8 | 2.8 | 2.0 | 0.7 | 2.6 | 1.7 | 0.6 | 2.3 | 1.5 | 0.6 | 1.6 | 1.2 |
| Steroids (without a doctor's order) | 0.6 | 0.5 | 0.5 | 0.4 | 0.5 | 0.5 | 0.2 | 0.6 | 0.4 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.1 | 0.1 | 0.1 |
| Over-the-Counter Drugs | 2.5 | -- | -- | 2.9 | 3.6 | 3.3 | 1.7 | 2.9 | 2.4 | 1.9 | 3.0 | 2.5 | 1.5 | 2.2 | 1.9 | 1.5 | 2.2 | 1.9 |
| Any illicit drug | 12.7 | 21.6 | 17.8 | 13.4 | 23.9 | 19.4 | 9.0 | 22.9 | 16.8 | 8.6 | 21.6 | 15.8 | 7.3 | 19.3 | 14.0 | 7.8 | 18.4 | 13.7 |
| Any illicit drug other than marijuana | 9.2 | 10.6 | 10.0 | 10.3 | 11.8 | 11.2 | 7.0 | 10.9 | 9.2 | 6.2 | 10.0 | 8.3 | 5.4 | 8.5 | 7.1 | 5.7 | 7.3 | 6.6 |
| Alcohol only | 11.4 | 22.9 | 17.9 | 9.5 | 20.0 | 15.4 | 7.3 | 17.5 | 13.0 | 6.0 | 15.1 | 11.0 | 5.4 | 15.1 | 10.9 | 5.3 | 12.7 | 9.4 |
| Alcohol or any illicit drug | 23.9 | 44.1 | 35.3 | 22.6 | 43.5 | 34.4 | 16.1 | 40.0 | 29.4 | 14.4 | 36.3 | 26.5 | 12.4 | 34.1 | 24.5 | 12.9 | 30.3 | 22.6 |
| Any illicit drug, but no alcohol | 5.3 | 4.8 | 5.0 | 6.3 | 6.1 | 6.2 | 4.9 | 6.0 | 5.5 | 4.6 | 7.4 | 6.1 | 4.5 | 6.7 | 5.7 | 4.6 | 7.1 | 6.0 |

Note: The first 18 data rows show results for alcohol, cigarettes, and other drugs. The last five data rows show results for various combinations of drugs. Binge drinking is defined as having had five or more alcoholic drinks in a row in the past two weeks. Ecstasy, Rohypnol, GHB and ketamine are provided as examples in the question about club drugs. The symbol "--" indicates that data are not available.

Table 6. Percentages of Big Bend Community Based Care and Florida Statewide youth who reported early ATOD use, perceived risk of harm, personal disapproval, parental disapproval and peer disapproval, 2018

| | Big Bend Community Based Care | | | | | | | Florida Statewide | | | | | | |
|--|-------------------------------|-------------|--------|------|------------|------------|-------|-------------------|-------------|--------|------|------------|------------|-------|
| | Middle School | High School | Female | Male | Ages 10-14 | Ages 15-17 | Total | Middle School | High School | Female | Male | Ages 10-14 | Ages 15-17 | Total |
| Early ATOD use | | | | | | | | | | | | | | |
| More than a sip of alcohol | -- | 17.9 | 16.9 | 18.9 | -- | 17.4 | -- | -- | 17.6 | 17.1 | 18.1 | -- | 17.1 | -- |
| Drinking at least once a month | -- | 3.0 | 2.4 | 3.4 | -- | 2.7 | -- | -- | 2.5 | 2.4 | 2.5 | -- | 2.5 | -- |
| Smoke cigarettes | -- | 10.0 | 9.1 | 10.8 | -- | 10.2 | -- | -- | 7.4 | 6.9 | 7.9 | -- | 7.6 | -- |
| Smoke marijuana | -- | 9.8 | 9.2 | 10.5 | -- | 9.7 | -- | -- | 9.3 | 8.0 | 10.6 | -- | 9.1 | -- |
| Great risk of harm | | | | | | | | | | | | | | |
| 1 or 2 drinks nearly every day | 46.7 | 40.2 | 47.0 | 39.3 | 46.5 | 40.0 | 43.1 | 49.3 | 43.3 | 49.1 | 42.8 | 48.7 | 43.6 | 45.9 |
| 5+ drinks once or twice a week | 61.0 | 51.8 | 60.5 | 51.5 | 60.9 | 51.7 | 55.9 | 62.1 | 55.0 | 61.3 | 54.9 | 61.9 | 55.3 | 58.0 |
| 1+ packs of cigarettes per day | 65.1 | 63.1 | 65.1 | 62.8 | 65.0 | 63.2 | 64.0 | 66.8 | 67.1 | 67.6 | 66.6 | 67.3 | 67.2 | 67.0 |
| Marijuana once or twice a week | 49.8 | 24.5 | 37.0 | 34.7 | 48.5 | 24.3 | 35.8 | 49.1 | 23.3 | 35.2 | 33.7 | 47.1 | 23.1 | 34.4 |
| Try marijuana once or twice | 32.7 | 15.6 | 23.4 | 23.1 | 31.2 | 15.8 | 23.3 | 33.2 | 15.8 | 22.8 | 23.6 | 31.5 | 15.7 | 23.2 |
| Use prescription drugs | 67.6 | 64.3 | 67.8 | 63.9 | 67.8 | 63.7 | 65.8 | 68.2 | 67.1 | 69.5 | 65.9 | 68.2 | 67.0 | 67.6 |
| Think it would be wrong to... | | | | | | | | | | | | | | |
| Smoke cigarettes | 94.6 | 85.1 | 89.9 | 88.7 | 94.5 | 86.3 | 89.3 | 95.6 | 89.8 | 92.6 | 92.1 | 95.6 | 90.8 | 92.3 |
| Drink alcohol regularly | 88.0 | 65.2 | 75.6 | 75.0 | 86.6 | 65.5 | 75.3 | 88.4 | 67.2 | 75.9 | 76.7 | 87.1 | 67.7 | 76.3 |
| Smoke marijuana | 88.5 | 60.9 | 73.0 | 73.3 | 87.2 | 61.0 | 73.1 | 88.5 | 60.9 | 72.9 | 72.8 | 86.7 | 61.1 | 72.8 |
| Smoke synthetic marijuana | 92.4 | 88.4 | 90.8 | 89.7 | 92.0 | 88.5 | 90.2 | 92.3 | 88.3 | 90.0 | 90.1 | 91.9 | 88.4 | 90.0 |
| Use other illicit drugs | 97.2 | 94.0 | 96.6 | 94.2 | 96.9 | 94.2 | 95.4 | 97.3 | 94.3 | 96.2 | 95.1 | 97.2 | 94.6 | 95.6 |
| Parents think it would be wrong to... | | | | | | | | | | | | | | |
| 1 or 2 drinks nearly every day | 97.5 | 93.4 | 95.4 | 95.0 | 97.4 | 94.0 | 95.2 | 97.6 | 94.2 | 95.7 | 95.5 | 97.3 | 94.4 | 95.6 |
| Smoke cigarettes | 98.5 | 96.1 | 97.3 | 97.0 | 98.3 | 96.9 | 97.2 | 98.8 | 97.4 | 98.1 | 97.9 | 98.8 | 97.7 | 98.0 |
| Smoke marijuana | 96.0 | 86.4 | 90.6 | 90.8 | 95.6 | 86.8 | 90.7 | 96.4 | 87.9 | 91.8 | 91.2 | 96.0 | 88.2 | 91.5 |
| Use prescription drugs | 98.1 | 96.2 | 96.8 | 97.3 | 97.8 | 96.2 | 97.1 | 98.1 | 97.1 | 97.3 | 97.8 | 98.0 | 97.1 | 97.5 |
| Friends think it would be wrong to... | | | | | | | | | | | | | | |
| 1 or 2 drinks nearly every day | 90.9 | 78.0 | 84.9 | 82.7 | 89.9 | 77.9 | 83.7 | 90.9 | 81.2 | 87.1 | 83.7 | 90.0 | 81.3 | 85.3 |
| Smoke tobacco | 93.6 | 82.7 | 88.2 | 87.0 | 92.7 | 84.1 | 87.6 | 94.4 | 88.7 | 92.1 | 90.2 | 94.1 | 89.3 | 91.1 |
| Smoke marijuana | 87.3 | 58.4 | 70.6 | 72.0 | 85.3 | 58.4 | 71.3 | 87.0 | 59.1 | 71.5 | 70.6 | 84.7 | 59.3 | 71.0 |
| Use prescription drugs | 94.6 | 88.8 | 92.4 | 90.5 | 94.1 | 88.9 | 91.4 | 95.0 | 91.8 | 94.1 | 92.3 | 94.7 | 91.9 | 93.2 |

Note: Early ATOD use is defined as the percentage of high school students who started using at age 13 or younger. The symbol "--" indicates that data are not available.

Table 7. Trends in early ATOD use and attitudes toward substance use for Big Bend Community Based Care, 2008 to 2018

| | 2008 | | | 2010 | | | 2012 | | | 2014 | | | 2016 | | | 2018 | | |
|--------------------------------------|---------------|-------------|-------|---------------|-------------|-------|---------------|-------------|-------|---------------|-------------|-------|---------------|-------------|-------|---------------|-------------|-------|
| | Middle School | High School | Total | Middle School | High School | Total | Middle School | High School | Total | Middle School | High School | Total | Middle School | High School | Total | Middle School | High School | Total |
| <i>Early ATOD use</i> | | | | | | | | | | | | | | | | | | |
| Try alcohol | -- | 31.1 | -- | -- | 26.5 | -- | -- | 26.0 | -- | -- | 21.9 | -- | -- | 20.8 | -- | -- | 17.9 | -- |
| Drink monthly | -- | 6.2 | -- | -- | 5.4 | -- | -- | 5.8 | -- | -- | 4.4 | -- | -- | 3.9 | -- | -- | 3.0 | -- |
| Smoke cigarettes | -- | 24.1 | -- | -- | 20.8 | -- | -- | 18.1 | -- | -- | 15.6 | -- | -- | 13.2 | -- | -- | 10.0 | -- |
| Smoke marijuana | -- | 11.8 | -- | -- | 12.4 | -- | -- | 11.8 | -- | -- | 11.4 | -- | -- | 11.4 | -- | -- | 9.8 | -- |
| <i>Great risk of harm</i> | | | | | | | | | | | | | | | | | | |
| Daily alcohol use | 43.2 | 40.1 | 41.5 | 42.4 | 39.9 | 41.0 | 43.6 | 36.6 | 39.7 | 44.3 | 37.2 | 40.4 | 44.9 | 36.6 | 40.2 | 46.7 | 40.2 | 43.1 |
| 5+ drinks daily | -- | -- | -- | -- | -- | -- | -- | -- | -- | 57.8 | 49.4 | 53.1 | 59.2 | 48.3 | 53.1 | 61.0 | 51.8 | 55.9 |
| Daily cigarette use | 67.4 | 66.2 | 66.7 | 67.4 | 66.4 | 66.8 | 67.3 | 67.1 | 67.2 | 67.5 | 68.8 | 68.3 | 68.4 | 65.2 | 66.6 | 65.1 | 63.1 | 64.0 |
| Marijuana weekly* | -- | -- | -- | -- | -- | -- | -- | -- | -- | 55.3 | 29.4 | 40.9 | 53.9 | 26.4 | 38.5 | 49.8 | 24.5 | 35.8 |
| Try marijuana | 42.6 | 24.2 | 32.2 | 40.7 | 23.3 | 30.9 | 40.0 | 20.7 | 29.2 | 39.5 | 18.0 | 27.5 | 37.5 | 17.7 | 26.4 | 32.7 | 15.6 | 23.3 |
| Prescription drugs | -- | -- | -- | -- | -- | -- | -- | 64.3 | 64.3 | 72.2 | 67.4 | 69.5 | 70.4 | 63.8 | 66.7 | 67.6 | 64.3 | 65.8 |
| <i>Think it wrong</i> | | | | | | | | | | | | | | | | | | |
| Smoke cigarettes | 86.9 | 68.9 | 76.7 | 88.1 | 70.2 | 78.1 | 92.3 | 77.0 | 83.8 | 93.6 | 80.7 | 86.5 | 95.0 | 82.5 | 88.0 | 94.6 | 85.1 | 89.3 |
| Drink regularly | 78.9 | 54.7 | 65.2 | 79.7 | 56.1 | 66.4 | 86.1 | 59.2 | 71.1 | 88.4 | 61.7 | 73.6 | 88.8 | 63.1 | 74.4 | 88.0 | 65.2 | 75.3 |
| Smoke marijuana | 88.8 | 70.8 | 78.5 | 88.0 | 67.1 | 76.2 | 90.8 | 68.5 | 78.4 | 89.5 | 65.4 | 76.2 | 90.2 | 63.7 | 75.3 | 88.5 | 60.9 | 73.1 |
| Smoke synthetic marijuana | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 94.5 | 88.7 | 91.3 | 92.4 | 88.4 | 90.2 |
| Use other illicit drugs | 96.2 | 93.6 | 94.7 | 96.0 | 92.6 | 94.1 | 97.4 | 93.9 | 95.4 | 97.9 | 93.2 | 95.3 | 97.9 | 92.5 | 94.9 | 97.2 | 94.0 | 95.4 |
| <i>Parents think it wrong</i> | | | | | | | | | | | | | | | | | | |
| Daily alcohol use* | -- | -- | -- | -- | -- | -- | -- | -- | -- | 97.3 | 92.2 | 94.5 | 97.6 | 92.8 | 94.9 | 97.5 | 93.4 | 95.2 |
| Smoke cigarettes | 95.9 | 90.3 | 92.8 | 96.2 | 90.2 | 92.8 | 97.5 | 92.2 | 94.5 | 98.4 | 94.3 | 96.1 | 98.1 | 95.2 | 96.5 | 98.5 | 96.1 | 97.2 |
| Smoke marijuana | 96.7 | 91.6 | 93.9 | 96.2 | 90.6 | 93.1 | 97.5 | 89.7 | 93.2 | 96.5 | 89.8 | 92.8 | 96.4 | 88.7 | 92.1 | 96.0 | 86.4 | 90.7 |
| Prescription drugs | -- | -- | -- | -- | -- | -- | -- | -- | -- | 98.2 | 96.2 | 97.1 | 98.4 | 96.6 | 97.4 | 98.1 | 96.2 | 97.1 |
| <i>Friends think it wrong</i> | | | | | | | | | | | | | | | | | | |
| Daily alcohol use | -- | -- | -- | -- | -- | -- | -- | -- | -- | 89.6 | 73.3 | 80.5 | 90.6 | 75.9 | 82.4 | 90.9 | 78.0 | 83.7 |
| Smoke tobacco | -- | -- | -- | -- | -- | -- | -- | -- | -- | 92.5 | 79.1 | 85.1 | 93.3 | 82.1 | 87.1 | 93.6 | 82.7 | 87.6 |
| Smoke marijuana | -- | -- | -- | -- | -- | -- | -- | -- | -- | 88.3 | 62.3 | 73.9 | 88.9 | 62.1 | 73.9 | 87.3 | 58.4 | 71.3 |
| Prescription drugs | -- | -- | -- | -- | -- | -- | -- | -- | -- | 96.0 | 88.3 | 91.7 | 95.8 | 87.8 | 91.3 | 94.6 | 88.8 | 91.4 |

Note: Early ATOD use is defined as the percentage of high school students who started using at age 13 or younger. The symbol "--" indicates that data are not available.

* In 2014, the description of marijuana use in the risk of harm question was changed from "regularly" to "once or twice a week." Also, in 2014, the description of alcohol use in the parent disapproval question was changed from "regularly" to "nearly every day." For these items, care should be exercised when comparing 2014 data to previous years.

Table 8. Among high school drinkers, usual source of alcohol within the past 30 days, Big Bend Community Based Care and Florida Statewide youth, 2018

| | Big Bend Community Based Care | | | | | | | Florida Statewide | | | | | | |
|--------------------------------------|-------------------------------|--------------------|---------------|-------------|-------------------|-------------------|--------------|----------------------|--------------------|---------------|-------------|-------------------|-------------------|--------------|
| | <i>Middle School</i> | <i>High School</i> | <i>Female</i> | <i>Male</i> | <i>Ages 10-14</i> | <i>Ages 15-17</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Female</i> | <i>Male</i> | <i>Ages 10-14</i> | <i>Ages 15-17</i> | <i>Total</i> |
| Bought in a store | -- | 4.3 | 2.2 | 7.2 | -- | 3.9 | -- | -- | 7.9 | 4.9 | 11.3 | -- | 7.0 | -- |
| Bought in a restaurant, bar, or club | -- | 1.6 | 1.8 | 1.4 | -- | 1.4 | -- | -- | 1.7 | 2.1 | 1.2 | -- | 1.6 | -- |
| Bought at a public event | -- | 0.4 | 0.0 | 0.8 | -- | 0.5 | -- | -- | 1.1 | 1.2 | 1.1 | -- | 1.1 | -- |
| Someone bought it for me | -- | 21.3 | 21.8 | 21.0 | -- | 20.7 | -- | -- | 15.2 | 15.4 | 15.2 | -- | 15.1 | -- |
| Someone gave it to me | -- | 39.9 | 44.2 | 34.0 | -- | 39.9 | -- | -- | 41.9 | 45.7 | 37.4 | -- | 42.3 | -- |
| Took it from a store | -- | 0.9 | 0.4 | 1.7 | -- | 1.1 | -- | -- | 0.6 | 0.4 | 0.7 | -- | 0.5 | -- |
| Took it from a family member | -- | 15.2 | 15.3 | 14.5 | -- | 16.4 | -- | -- | 13.8 | 16.1 | 10.9 | -- | 14.6 | -- |
| Some other way | -- | 16.5 | 14.3 | 19.4 | -- | 16.1 | -- | -- | 17.8 | 14.3 | 22.2 | -- | 17.9 | -- |

Note: The symbol "--" indicates that data are not available.

Table 9. Among high school drinkers, usual drinking location within the past 30 days, Big Bend Community Based Care and Florida Statewide youth, 2018

| | Big Bend Community Based Care | | | | | | | Florida Statewide | | | | | | |
|--------------------------|-------------------------------|--------------------|---------------|-------------|-------------------|-------------------|--------------|----------------------|--------------------|---------------|-------------|-------------------|-------------------|--------------|
| | <i>Middle School</i> | <i>High School</i> | <i>Female</i> | <i>Male</i> | <i>Ages 10-14</i> | <i>Ages 15-17</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Female</i> | <i>Male</i> | <i>Ages 10-14</i> | <i>Ages 15-17</i> | <i>Total</i> |
| My home | -- | 38.9 | 37.5 | 40.2 | -- | 39.0 | -- | -- | 41.0 | 41.9 | 39.5 | -- | 41.5 | -- |
| Another person's home | -- | 41.9 | 45.3 | 38.2 | -- | 42.2 | -- | -- | 38.3 | 39.0 | 37.8 | -- | 38.2 | -- |
| Car or other vehicle | -- | 1.7 | 1.2 | 2.4 | -- | 2.0 | -- | -- | 1.5 | 1.7 | 1.3 | -- | 1.6 | -- |
| Restaurant, bar, or club | -- | 2.8 | 3.7 | 1.6 | -- | 2.9 | -- | -- | 3.3 | 4.4 | 1.6 | -- | 2.7 | -- |
| Public place | -- | 3.2 | 2.1 | 4.4 | -- | 3.6 | -- | -- | 3.6 | 2.8 | 4.5 | -- | 3.5 | -- |
| Public event | -- | 1.5 | 1.4 | 1.6 | -- | 1.4 | -- | -- | 1.7 | 1.5 | 1.9 | -- | 1.6 | -- |
| School property | -- | 0.9 | 0.5 | 1.3 | -- | 0.7 | -- | -- | 1.2 | 0.8 | 1.8 | -- | 1.3 | -- |
| Some other place | -- | 9.1 | 8.3 | 10.3 | -- | 8.2 | -- | -- | 9.4 | 7.8 | 11.6 | -- | 9.6 | -- |

Note: The symbol "--" indicates that data are not available.

Table 10. Among high school drinkers, number of drinks usually consumed per day, on the days students drank in the past 30 days, Big Bend Community Based Care and Florida Statewide youth, 2018

| | Big Bend Community Based Care | | | | | | | Florida Statewide | | | | | | |
|-----------|-------------------------------|--------------------|---------------|-------------|-------------------|-------------------|--------------|----------------------|--------------------|---------------|-------------|-------------------|-------------------|--------------|
| | <i>Middle School</i> | <i>High School</i> | <i>Female</i> | <i>Male</i> | <i>Ages 10-14</i> | <i>Ages 15-17</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Female</i> | <i>Male</i> | <i>Ages 10-14</i> | <i>Ages 15-17</i> | <i>Total</i> |
| 1 | -- | 27.8 | 30.7 | 23.9 | -- | 27.1 | -- | -- | 30.0 | 33.1 | 26.0 | -- | 30.5 | -- |
| 2 | -- | 23.4 | 25.8 | 20.4 | -- | 24.9 | -- | -- | 23.5 | 24.0 | 22.7 | -- | 23.8 | -- |
| 3 | -- | 14.9 | 17.1 | 12.2 | -- | 16.2 | -- | -- | 14.9 | 15.9 | 13.9 | -- | 14.7 | -- |
| 4 | -- | 11.1 | 10.4 | 12.3 | -- | 10.2 | -- | -- | 10.7 | 10.9 | 10.5 | -- | 10.6 | -- |
| 5 or more | -- | 22.7 | 16.0 | 31.2 | -- | 21.6 | -- | -- | 20.9 | 16.1 | 26.8 | -- | 20.4 | -- |

Note: The symbol "--" indicates that data are not available.

.....

Table 11. Percentages of Big Bend Community Based Care and Florida Statewide youth who reported drinking alcohol, smoking marijuana, or using another drug to get high before or during school in the past 12 months, 2018

| | Big Bend Community Based Care | | | | | | | Florida Statewide | | | | | | |
|-------------------------------|-------------------------------|--------------------|---------------|-------------|-------------------|-------------------|--------------|----------------------|--------------------|---------------|-------------|-------------------|-------------------|--------------|
| | <i>Middle School</i> | <i>High School</i> | <i>Female</i> | <i>Male</i> | <i>Ages 10-14</i> | <i>Ages 15-17</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Female</i> | <i>Male</i> | <i>Ages 10-14</i> | <i>Ages 15-17</i> | <i>Total</i> |
| Drinking alcohol | 3.1 | 7.7 | 5.8 | 5.5 | 3.4 | 7.8 | 5.7 | 3.7 | 6.7 | 5.9 | 4.8 | 3.8 | 7.0 | 5.4 |
| Smoking marijuana | 3.4 | 11.7 | 8.2 | 7.7 | 3.7 | 11.8 | 8.0 | 3.7 | 13.0 | 8.7 | 9.2 | 4.1 | 13.3 | 9.0 |
| Used another drug to get high | 1.6 | 4.6 | 3.4 | 3.1 | 2.0 | 4.1 | 3.3 | 1.7 | 3.7 | 2.6 | 3.1 | 1.9 | 3.7 | 2.9 |

Table 12. Percentages of Big Bend Community Based Care and Florida Statewide youth who reported engaging in delinquent behavior within the past 12 months, 2018

| | Big Bend Community Based Care | | | | | | | Florida Statewide | | | | | | |
|---------------------------------------|-------------------------------|--------------------|---------------|-------------|-------------------|-------------------|--------------|----------------------|--------------------|---------------|-------------|-------------------|-------------------|--------------|
| | <i>Middle School</i> | <i>High School</i> | <i>Female</i> | <i>Male</i> | <i>Ages 10-14</i> | <i>Ages 15-17</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Female</i> | <i>Male</i> | <i>Ages 10-14</i> | <i>Ages 15-17</i> | <i>Total</i> |
| Carrying a handgun | 7.0 | 7.7 | 3.2 | 11.4 | 7.0 | 7.8 | 7.4 | 5.7 | 6.2 | 3.2 | 8.7 | 5.6 | 6.4 | 6.0 |
| Selling drugs | 1.5 | 5.2 | 2.5 | 4.5 | 1.5 | 5.3 | 3.5 | 1.5 | 5.7 | 2.7 | 5.1 | 1.8 | 5.6 | 3.9 |
| Attempting to steal a vehicle | 1.1 | 1.6 | 0.9 | 1.8 | 1.1 | 1.7 | 1.4 | 1.4 | 1.6 | 1.2 | 1.8 | 1.4 | 1.6 | 1.5 |
| Being arrested | 2.1 | 2.9 | 1.6 | 3.5 | 2.1 | 3.1 | 2.5 | 1.7 | 2.6 | 1.7 | 2.8 | 1.8 | 2.7 | 2.3 |
| Taking a handgun to school | 0.4 | 1.0 | 0.3 | 1.1 | 0.4 | 1.0 | 0.7 | 0.4 | 0.8 | 0.3 | 0.9 | 0.5 | 0.8 | 0.6 |
| Getting suspended | 10.1 | 7.8 | 7.1 | 10.6 | 9.6 | 8.3 | 8.8 | 10.7 | 8.7 | 7.3 | 11.7 | 10.4 | 8.9 | 9.5 |
| Attacking someone with intent to harm | 8.1 | 6.0 | 6.0 | 7.8 | 7.9 | 6.2 | 6.9 | 7.2 | 5.9 | 5.6 | 7.3 | 7.0 | 6.0 | 6.5 |

Table 13. Trends in delinquent behaviors for Big Bend Community Based Care, 2008 to 2018

| | 2008 | | | 2010 | | | 2012 | | | 2014 | | | 2016 | | | 2018 | | |
|---------------------------------------|----------------------|--------------------|--------------|----------------------|--------------------|--------------|----------------------|--------------------|--------------|----------------------|--------------------|--------------|----------------------|--------------------|--------------|----------------------|--------------------|--------------|
| | <i>Middle School</i> | <i>High School</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Total</i> |
| Carrying a handgun | 5.2 | 5.5 | 5.4 | 6.0 | 5.7 | 5.8 | 5.8 | 4.9 | 5.3 | 6.4 | 6.7 | 6.5 | 7.1 | 6.7 | 6.9 | 7.0 | 7.7 | 7.4 |
| Selling drugs | 2.8 | 8.5 | 6.0 | 3.5 | 9.1 | 6.7 | 2.0 | 8.2 | 5.4 | 2.0 | 6.7 | 4.6 | 1.5 | 6.3 | 4.2 | 1.5 | 5.2 | 3.5 |
| Attempting to steal a vehicle | 2.5 | 2.8 | 2.7 | 2.1 | 2.2 | 2.1 | 1.2 | 2.6 | 2.0 | 1.1 | 1.8 | 1.5 | 0.6 | 2.0 | 1.4 | 1.1 | 1.6 | 1.4 |
| Being arrested | 4.1 | 6.5 | 5.5 | 3.7 | 4.9 | 4.4 | 2.8 | 4.3 | 3.6 | 2.3 | 3.0 | 2.7 | 1.9 | 3.0 | 2.5 | 2.1 | 2.9 | 2.5 |
| Taking a handgun to school | 0.7 | 1.1 | 0.9 | 0.9 | 1.1 | 1.0 | 0.4 | 0.9 | 0.7 | 0.4 | 0.8 | 0.6 | 0.3 | 0.8 | 0.6 | 0.4 | 1.0 | 0.7 |
| Getting suspended | 16.9 | 15.2 | 15.9 | 15.7 | 13.1 | 14.2 | 13.0 | 11.0 | 11.9 | 11.8 | 9.3 | 10.4 | 10.9 | 9.3 | 10.1 | 10.1 | 7.8 | 8.8 |
| Attacking someone with intent to harm | 11.9 | 13.0 | 12.5 | 11.9 | 10.7 | 11.2 | 7.6 | 8.2 | 7.9 | 6.8 | 6.8 | 6.8 | 6.8 | 6.8 | 6.8 | 8.1 | 6.0 | 6.9 |

Table 14. Percentages of Big Bend Community Based Care and Florida Statewide youth who reported symptoms of depression, 2018

| | Big Bend Community Based Care | | | | | | | Florida Statewide | | | | | | |
|--|-------------------------------|--------------------|---------------|-------------|-------------------|-------------------|--------------|----------------------|--------------------|---------------|-------------|-------------------|-------------------|--------------|
| | <i>Middle School</i> | <i>High School</i> | <i>Female</i> | <i>Male</i> | <i>Ages 10-14</i> | <i>Ages 15-17</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Female</i> | <i>Male</i> | <i>Ages 10-14</i> | <i>Ages 15-17</i> | <i>Total</i> |
| Sometimes I think that life is not worth it | 26.7 | 32.0 | 38.7 | 20.7 | 26.7 | 33.7 | 29.6 | 26.5 | 29.5 | 35.9 | 20.5 | 26.9 | 30.1 | 28.2 |
| At times I think I am no good at all | 39.8 | 44.1 | 52.8 | 31.9 | 40.4 | 44.9 | 42.2 | 38.2 | 40.8 | 49.9 | 29.6 | 38.8 | 41.6 | 39.7 |
| All in all, I am inclined to think that I am a failure | 24.8 | 28.2 | 34.2 | 19.3 | 25.0 | 29.9 | 26.7 | 24.5 | 26.3 | 32.1 | 18.8 | 24.8 | 27.3 | 25.5 |
| In the past year, have you felt depressed or sad MOST days, even if you felt OK sometimes? | 41.6 | 46.0 | 54.6 | 33.8 | 41.5 | 46.8 | 44.1 | 40.6 | 44.1 | 53.1 | 32.2 | 40.5 | 45.3 | 42.6 |

Table 15. Percentages of Big Bend Community Based Care and Florida Statewide youth who reported involvement in bullying behavior, 2018

| | Big Bend Community Based Care | | | | | | | Florida Statewide | | | | | | |
|-------------------------------------|-------------------------------|--------------------|---------------|-------------|-------------------|-------------------|--------------|----------------------|--------------------|---------------|-------------|-------------------|-------------------|--------------|
| | <i>Middle School</i> | <i>High School</i> | <i>Female</i> | <i>Male</i> | <i>Ages 10-14</i> | <i>Ages 15-17</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Female</i> | <i>Male</i> | <i>Ages 10-14</i> | <i>Ages 15-17</i> | <i>Total</i> |
| Skipped school because of bullying | 9.2 | 9.1 | 13.6 | 4.8 | 8.9 | 9.5 | 9.1 | 7.7 | 8.4 | 11.7 | 4.5 | 7.7 | 8.5 | 8.1 |
| Was ever kicked or shoved | 34.6 | 22.3 | 27.3 | 27.9 | 33.1 | 23.2 | 27.7 | 33.3 | 20.3 | 24.7 | 26.8 | 32.3 | 20.3 | 25.8 |
| Was ever taunted or teased | 54.8 | 45.0 | 54.4 | 44.4 | 54.7 | 46.2 | 49.2 | 54.1 | 42.7 | 52.1 | 43.0 | 53.4 | 43.3 | 47.4 |
| Was ever a victim of cyber bullying | 22.2 | 26.8 | 33.2 | 16.7 | 22.6 | 27.8 | 24.7 | 20.7 | 24.5 | 30.3 | 15.8 | 21.1 | 25.0 | 22.8 |
| Ever physically bullied others | 17.1 | 10.7 | 10.5 | 16.4 | 16.4 | 10.9 | 13.5 | 17.4 | 10.5 | 10.9 | 15.9 | 16.7 | 10.4 | 13.4 |
| Ever verbally bullied others | 27.4 | 22.3 | 23.9 | 24.9 | 27.3 | 22.3 | 24.5 | 28.1 | 22.2 | 23.9 | 25.5 | 27.8 | 22.5 | 24.7 |
| Ever cyber bullied others | 9.3 | 11.5 | 12.1 | 8.7 | 9.2 | 11.7 | 10.5 | 9.3 | 10.7 | 11.6 | 8.6 | 9.5 | 10.9 | 10.1 |

Table 16. Percentages of Big Bend Community Based Care and Florida Statewide youth who reported gang membership, 2018

| | Big Bend Community Based Care | | | | | | | Florida Statewide | | | | | | |
|-----------------------------------|-------------------------------|--------------------|---------------|-------------|-------------------|-------------------|--------------|----------------------|--------------------|---------------|-------------|-------------------|-------------------|--------------|
| | <i>Middle School</i> | <i>High School</i> | <i>Female</i> | <i>Male</i> | <i>Ages 10-14</i> | <i>Ages 15-17</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Female</i> | <i>Male</i> | <i>Ages 10-14</i> | <i>Ages 15-17</i> | <i>Total</i> |
| Have you ever belonged to a gang? | 4.8 | 4.3 | 2.8 | 6.3 | 4.5 | 4.6 | 4.5 | 3.8 | 3.7 | 2.4 | 5.0 | 3.6 | 3.8 | 3.7 |
| Did that gang have a name? | 31.7 | 20.3 | 18.9 | 28.1 | 29.8 | 21.7 | 24.5 | 25.5 | 16.2 | 14.4 | 23.1 | 24.0 | 17.0 | 19.4 |
| Are you a gang member now? | -- | 2.6 | 1.5 | 3.8 | -- | 2.7 | -- | -- | 2.3 | 1.4 | 3.3 | -- | 2.3 | -- |

Note: The prevalence rates for “Did that gang have a name?” exclude students who reported that they have never belonged to a gang. The symbol “--” indicates that data are not available.

Table 17. Percentages of Big Bend Community Based Care and Florida Statewide youth who “agree” or “strongly agree” with statements indicating impulsiveness or a lack of self-control, 2018

| | Big Bend Community Based Care | | | | | | | Florida Statewide | | | | | | |
|---|-------------------------------|--------------------|---------------|-------------|-------------------|-------------------|--------------|----------------------|--------------------|---------------|-------------|-------------------|-------------------|--------------|
| | <i>Middle School</i> | <i>High School</i> | <i>Female</i> | <i>Male</i> | <i>Ages 10-14</i> | <i>Ages 15-17</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Female</i> | <i>Male</i> | <i>Ages 10-14</i> | <i>Ages 15-17</i> | <i>Total</i> |
| I often do what brings me pleasure now | 27.5 | 25.5 | 26.4 | 26.4 | 26.6 | 26.3 | 26.3 | 29.3 | 28.3 | 29.0 | 28.3 | 28.9 | 28.8 | 28.7 |
| I'm more concerned with the short run | 22.8 | 19.8 | 21.2 | 20.9 | 21.7 | 20.2 | 21.1 | 25.6 | 21.6 | 24.2 | 22.3 | 25.1 | 22.4 | 23.3 |
| Getting in trouble is exciting | 23.9 | 28.5 | 26.4 | 26.4 | 24.3 | 28.8 | 26.5 | 23.3 | 27.7 | 25.6 | 25.9 | 24.1 | 27.9 | 25.8 |
| Excitement is more important than security | 24.6 | 24.0 | 21.5 | 27.0 | 23.8 | 24.8 | 24.3 | 23.5 | 23.5 | 21.4 | 25.5 | 23.5 | 23.5 | 23.5 |
| People better stay away from me when I'm really angry | 38.0 | 33.5 | 36.6 | 34.2 | 37.3 | 34.1 | 35.5 | 36.0 | 32.5 | 35.8 | 32.1 | 35.7 | 33.0 | 34.0 |
| I get upset when I have a disagreement with someone | 48.3 | 39.1 | 49.1 | 37.0 | 47.9 | 39.2 | 43.1 | 45.0 | 38.0 | 47.4 | 34.6 | 45.0 | 38.3 | 41.0 |

Note: The symbol “--” indicates that data are not available.

Table 18. Percentages of Big Bend Community Based Care and Florida Statewide youth who reported participation in extracurricular activities, 2018

| | Big Bend Community Based Care | | | | | | | Florida Statewide | | | | | | |
|------------------------------------|-------------------------------|--------------------|---------------|-------------|-------------------|-------------------|--------------|----------------------|--------------------|---------------|-------------|-------------------|-------------------|--------------|
| | <i>Middle School</i> | <i>High School</i> | <i>Female</i> | <i>Male</i> | <i>Ages 10-14</i> | <i>Ages 15-17</i> | <i>Total</i> | <i>Middle School</i> | <i>High School</i> | <i>Female</i> | <i>Male</i> | <i>Ages 10-14</i> | <i>Ages 15-17</i> | <i>Total</i> |
| School Sports | 42.2 | 41.4 | 38.0 | 45.3 | 42.6 | 42.3 | 41.8 | 36.0 | 39.7 | 34.7 | 41.4 | 36.8 | 40.0 | 38.1 |
| Organized Sports Outside of School | 37.1 | 18.4 | 25.8 | 27.5 | 35.4 | 18.7 | 26.7 | 38.6 | 19.8 | 25.7 | 30.1 | 37.2 | 19.7 | 27.9 |
| School Band | 19.2 | 9.8 | 14.8 | 13.3 | 18.3 | 10.1 | 14.0 | 17.3 | 8.1 | 12.5 | 11.6 | 16.6 | 8.0 | 12.0 |
| School Club(s) | 25.5 | 33.9 | 38.7 | 22.0 | 26.3 | 34.1 | 30.1 | 22.2 | 30.2 | 34.6 | 19.4 | 22.8 | 30.4 | 26.8 |
| Community Club(s) | 8.1 | 11.6 | 12.7 | 7.5 | 8.3 | 11.4 | 10.1 | 8.5 | 11.7 | 13.3 | 7.5 | 8.4 | 11.6 | 10.3 |

Table 19. Percentages of high school students who reported *riding in* a vehicle within the past 30 days driven by someone who had been drinking alcohol or using marijuana, or *driving* a vehicle within the past 30 days after drinking alcohol or using marijuana, among Big Bend Community Based Care and Florida Statewide youth, 2018

| | Big Bend Community Based Care | | | | | | | Florida Statewide | | | | | | |
|-----------------------------------|-------------------------------|-------------|--------|------|------------|------------|-------|-------------------|-------------|--------|------|------------|------------|-------|
| | Middle School | High School | Female | Male | Ages 10-14 | Ages 15-17 | Total | Middle School | High School | Female | Male | Ages 10-14 | Ages 15-17 | Total |
| <i>Riding in a vehicle</i> | | | | | | | | | | | | | | |
| Alcohol | -- | 16.0 | 17.8 | 13.9 | -- | 15.7 | -- | -- | 14.3 | 15.1 | 13.4 | -- | 14.3 | -- |
| Marijuana | -- | 21.8 | 24.2 | 19.4 | -- | 22.5 | -- | -- | 22.9 | 24.4 | 21.5 | -- | 22.8 | -- |
| <i>Driving a vehicle</i> | | | | | | | | | | | | | | |
| Alcohol | -- | 4.9 | 4.5 | 5.3 | -- | 4.6 | -- | -- | 4.4 | 3.8 | 4.9 | -- | 4.3 | -- |
| Marijuana | -- | 9.6 | 9.5 | 9.7 | -- | 9.3 | -- | -- | 9.5 | 8.7 | 10.1 | -- | 9.2 | -- |

Note: The symbol "--" indicates that data are not available.

Table 20. Trends for high school students *riding in* a vehicle within the past 30 days driven by someone who had been drinking alcohol or using marijuana, or *driving* a vehicle within the past 30 days after drinking alcohol or using marijuana, for Big Bend Community Based Care, 2012 and 2018

| | 2008 | | | 2010 | | | 2012 | | | 2014 | | | 2016 | | | 2018 | | |
|-----------------------------------|---------------|-------------|-------|---------------|-------------|-------|---------------|-------------|-------|---------------|-------------|-------|---------------|-------------|-------|---------------|-------------|-------|
| | Middle School | High School | Total | Middle School | High School | Total | Middle School | High School | Total | Middle School | High School | Total | Middle School | High School | Total | Middle School | High School | Total |
| <i>Riding in a vehicle</i> | | | | | | | | | | | | | | | | | | |
| Alcohol | -- | -- | -- | -- | -- | -- | -- | 23.1 | -- | -- | 19.2 | -- | -- | 17.7 | -- | -- | 16.0 | -- |
| Marijuana | -- | -- | -- | -- | -- | -- | -- | 24.4 | -- | -- | 22.0 | -- | -- | 23.1 | -- | -- | 21.8 | -- |
| <i>Driving a vehicle</i> | | | | | | | | | | | | | | | | | | |
| Alcohol | -- | -- | -- | -- | -- | -- | -- | 9.7 | -- | -- | 7.4 | -- | -- | 6.6 | -- | -- | 4.9 | -- |
| Marijuana | -- | -- | -- | -- | -- | -- | -- | 11.4 | -- | -- | 11.2 | -- | -- | 10.6 | -- | -- | 9.6 | -- |

Note: The symbol "--" indicates that data are not available.

Table 21. Protective factor prevalence rates for Big Bend Community Based Care, Florida Statewide youth and the national normative database, 2018

| Domain | Scale | Big Bend Community Based Care | | Florida Statewide | | National Norms | |
|--------------------------------|--|-------------------------------|--------------------|----------------------|--------------------|----------------------|--------------------|
| | | <i>Middle School</i> | <i>High School</i> | <i>Middle School</i> | <i>High School</i> | <i>Middle School</i> | <i>High School</i> |
| Family | Family Opportunities for Prosocial Involvement | 58 | 59 | 58 | 57 | 59 | 54 |
| | Family Rewards for Prosocial Involvement | 51 | 53 | 50 | 51 | 54 | 55 |
| School | School Opportunities for Prosocial Involvement | 56 | 66 | 54 | 64 | 57 | 60 |
| | School Rewards for Prosocial Involvement | 48 | 59 | 45 | 55 | 53 | 58 |
| Peer and Individual | Religiosity | 52 | 58 | 46 | 54 | 56 | 62 |
| Average Prevalence Rate | | 53 | 59 | 51 | 56 | 56 | 58 |

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.

Table 22. Risk factor prevalence rates for Big Bend Community Based Care, Florida Statewide youth and the national normative database, 2018

| Domain | Scale | Big Bend Community Based Care | | Florida Statewide | | National Norms | |
|--------------------------------|--|-------------------------------|--------------------|----------------------|--------------------|----------------------|--------------------|
| | | <i>Middle School</i> | <i>High School</i> | <i>Middle School</i> | <i>High School</i> | <i>Middle School</i> | <i>High School</i> |
| Community | Community Disorganization | 36 | 37 | 38 | 40 | 47 | 47 |
| | Transitions and Mobility | 59 | 60 | 59 | 61 | 47 | 46 |
| | Laws and Norms Favorable to Drug Use | 38 | 29 | 38 | 32 | 42 | 42 |
| | Perceived Availability of Drugs | 33 | 25 | 35 | 24 | 45 | 45 |
| | Perceived Availability of Handguns | 29 | 39 | 24 | 34 | 25 | 42 |
| Family | Poor Family Management | 41 | 34 | 43 | 37 | 44 | 45 |
| | Family Conflict | 39 | 33 | 39 | 34 | 42 | 37 |
| School | Poor Academic Performance | 44 | 42 | 43 | 43 | 45 | 48 |
| | Lack of Commitment to School | 58 | 55 | 60 | 57 | 47 | 46 |
| Peer and Individual | Favorable Attitudes toward Antisocial Behavior | 42 | 37 | 43 | 36 | 40 | 46 |
| | Favorable Attitudes toward ATOD Use | 35 | 38 | 35 | 34 | 39 | 45 |
| | Early Initiation of Drug Use | 25 | 21 | 24 | 19 | 41 | 46 |
| Average Prevalence Rate | | 40 | 38 | 40 | 38 | 40 | 45 |

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.

Table 23. Protective factor prevalence rate trends among middle school students for Big Bend Community Based Care, 2008 to 2018

| Domain | | Big Bend Community Based Care | | | | | |
|--------------------------------|--|-------------------------------|-----------|-----------|-----------|-----------|-----------|
| Scale | | 2008 | 2010 | 2012 | 2014 | 2016 | 2018 |
| Family | Family Opportunities for Prosocial Involvement | 55 | 55 | 59 | 63 | 62 | 58 |
| | Family Rewards for Prosocial Involvement | 51 | 52 | 55 | 59 | 58 | 51 |
| School | School Opportunities for Prosocial Involvement | 46 | 47 | 54 | 57 | 57 | 56 |
| | School Rewards for Prosocial Involvement | 45 | 47 | 53 | 54 | 50 | 48 |
| Peer and Individual | Religiosity | 58 | 58 | 58 | 57 | 59 | 52 |
| Average Prevalence Rate | | 51 | 52 | 56 | 58 | 57 | 53 |

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.

Table 24. Risk factor prevalence rate trends among middle school students for Big Bend Community Based Care, 2008 to 2018

| Domain | | Big Bend Community Based Care | | | | | |
|-------------------------|--|-------------------------------|------|------|------|------|------|
| Scale | | 2008 | 2010 | 2012 | 2014 | 2016 | 2018 |
| Community | Community Disorganization | 46 | 46 | 44 | 42 | 38 | 36 |
| | Transitions and Mobility | 60 | 60 | 58 | 57 | 59 | 59 |
| | Laws and Norms Favorable to Drug Use | 47 | 46 | 38 | 38 | 37 | 38 |
| | Perceived Availability of Drugs | 51 | 48 | 39 | 38 | 38 | 33 |
| | Perceived Availability of Handguns | 33 | 31 | 29 | 31 | 31 | 29 |
| Family | Poor Family Management | 49 | 48 | 40 | 38 | 37 | 41 |
| | Family Conflict | 43 | 40 | 39 | 36 | 36 | 39 |
| School | Poor Academic Performance | 44 | 43 | 42 | 42 | 43 | 44 |
| | Lack of Commitment to School | 58 | 54 | 48 | 48 | 52 | 58 |
| Peer and Individual | Favorable Attitudes toward Antisocial Behavior | 48 | 48 | 40 | 37 | 39 | 42 |
| | Favorable Attitudes toward ATOD Use | 43 | 43 | 34 | 32 | 31 | 35 |
| | Early Initiation of Drug Use | 42 | 39 | 29 | 25 | 23 | 25 |
| Average Prevalence Rate | | 47 | 46 | 40 | 39 | 39 | 40 |

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.

Table 25. Protective factor prevalence rate trends among high school students for Big Bend Community Based Care, 2008 to 2018

| Domain | | Scale | | | | | |
|--------------------------------|--|-------------------------------|-----------|-----------|-----------|-----------|-----------|
| | | Big Bend Community Based Care | | | | | |
| | | 2008 | 2010 | 2012 | 2014 | 2016 | 2018 |
| Family | Family Opportunities for Prosocial Involvement | 54 | 56 | 57 | 59 | 61 | 59 |
| | Family Rewards for Prosocial Involvement | 55 | 55 | 55 | 58 | 56 | 53 |
| School | School Opportunities for Prosocial Involvement | 58 | 60 | 61 | 61 | 64 | 66 |
| | School Rewards for Prosocial Involvement | 57 | 61 | 61 | 61 | 59 | 59 |
| Peer and Individual | Religiosity | 68 | 67 | 67 | 66 | 64 | 58 |
| Average Prevalence Rate | | 58 | 60 | 60 | 61 | 61 | 59 |

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.

Table 26. Risk factor prevalence rate trends among high school students for Big Bend Community Based Care, 2008 to 2018

| Domain | | Big Bend Community Based Care | | | | | |
|--------------------------------|--|-------------------------------|-----------|-----------|-----------|-----------|-----------|
| Scale | | 2008 | 2010 | 2012 | 2014 | 2016 | 2018 |
| Community | Community Disorganization | 42 | 42 | 42 | 41 | 43 | 37 |
| | Transitions and Mobility | 64 | 62 | 59 | 59 | 63 | 60 |
| | Laws and Norms Favorable to Drug Use | 36 | 37 | 34 | 31 | 31 | 29 |
| | Perceived Availability of Drugs | 42 | 38 | 32 | 30 | 28 | 25 |
| | Perceived Availability of Handguns | 46 | 43 | 42 | 42 | 41 | 39 |
| Family | Poor Family Management | 48 | 44 | 40 | 38 | 36 | 34 |
| | Family Conflict | 36 | 34 | 33 | 31 | 33 | 33 |
| School | Poor Academic Performance | 44 | 45 | 44 | 40 | 44 | 42 |
| | Lack of Commitment to School | 49 | 51 | 46 | 49 | 53 | 55 |
| Peer and Individual | Favorable Attitudes toward Antisocial Behavior | 46 | 41 | 37 | 35 | 36 | 37 |
| | Favorable Attitudes toward ATOD Use | 43 | 42 | 39 | 39 | 38 | 38 |
| | Early Initiation of Drug Use | 38 | 34 | 32 | 28 | 26 | 21 |
| Average Prevalence Rate | | 45 | 43 | 40 | 39 | 39 | 38 |

Note: Because risk is associated with negative behavioral outcomes, it is better to have lower risk factor scale scores, not higher. Conversely, because protective factors are associated with better student behavioral outcomes, it is better to have protective factor scale scores with high values.